

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C07K 5/062, 5/065, 5/078, A61K 38/05		A1	(11) International Publication Number: WO 99/06431
			(43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/US98/15313 (22) International Filing Date: 30 July 1998 (30.07.98) (30) Priority Data: 08/920,394                      31 July 1997 (31.07.97)                      US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US                                      08/920,394 (CON) Filed on                              31 July 1997 (31.07.97) (71) Applicants (for all designated States except US): ATHENA NEUROSCIENCES, INC. [US/US]; 800 Gateway Boulevard, South San Francisco, CA 94080 (US). AMERICAN HOME PRODUCTS CORPORATION [US/US]; 5 Giraldi Farms, Madison, NJ 07940 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): THORSETT, Eugene, D. [US/US]; 571 Buena Vista, Moss Beach, CA 94038 (US). SEMKO, Christopher, M. [CA/US]; 2361 Carpenter Court, Fremont, CA 94539 (US). SARANTAKIS, Dimitrios [US/US]; 262 Sentinel Avenue, Newtown, PA 18940 (US). PLEISS, Michael, A. [US/US]; 848 Stella Court, Sunnyvale,		CA 94087 (US). LOMBARDO, Louis, John [US/US]; 412 Southwoods Road, Belle Mead, NJ 08502 (US). KREFT, Anthony [US/US]; 43 Barley Court, Langhorne, PA 19047 (US). KONRAD, Andrei, W. [US/US]; 95 Cervantes #105, San Francisco, CA 94123 (US). GRANT, Francine, S. [FR/US]; 3735 Sacramento Street, San Francisco, CA 94118 (US). DRESSEN, Darren, B. [US/US]; 3110 Casa de Campo #2, San Mateo, CA 94403 (US). DAPPEN, Michael, S. [US/US]; 640 Topaz Street, Redwood City, CA 94061 (US). BAUDY, Reinhardt, Bernhard [DE/US]; 5281 Harrington Court, Doylestown, PA 18901 (US). ASHWELL, Susan [GB/US]; 1015 Aspen Drive, Plainsboro, NJ 08536 (US). (74) Agents: SWISS, Gerald, F. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: SUBSTITUTED PHENYLALANINE TYPE COMPOUNDS WHICH INHIBIT LEUKOCYTE ADHESION MEDIATED BY VLA-4			
(57) Abstract			
Disclosed are compounds which bind VLA-4. Certain of these compounds also inhibit leukocyte adhesion and, in particular, leukocyte adhesion mediated by VLA-4. Such compounds are useful in the treatment of inflammatory diseases in a mammalian patient, e.g., human, such as asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes, inflammatory bowel disease, rheumatoid arthritis, tissue transplantation, tumor metastasis and myocardial ischemia. The compounds are also be administered for the treatment of inflammatory brain diseases such as multiple sclerosis.			

**SUBSTITUTED PHENYLALANINE TYPE COMPOUNDS WHICH  
INHIBIT LEUKOCYTE ADHESION MEDIATED BY VLA-4**

Cross-Reference to Related Application

This application claims the benefit of U.S. Provisional Application No. 60/\_\_\_\_\_, which was converted pursuant to 37 C.F.R. § 1.53(c)(2)(i) from U.S. Patent Application No. 08/920,394, filed July 31, 1997.

Field of the Invention

This invention relates to compounds which inhibit leukocyte adhesion and, in particular, leukocyte adhesion mediated by VLA-4.

5

References

The following publications, patents and patent applications are cited in this application as superscript numbers:

- |    |   |   |
|----|---|---|
| 10 | 1 | Hemler and Takada, <i>European Patent Application Publication No. 330,506</i> , published August 30, 1989 |
|    | 2 | Elices, et al., <i>Cell</i> , <u>60</u> :577-584 (1990)   |
| 15 | 3 | Springer, <i>Nature</i> , <u>346</u> :425-434 (1990)  |
|    | 4 | Osborn, <i>Cell</i> , <u>62</u> :3-6 (1990)   |
|    | 5 | Vedder, et al., <i>Surgery</i> , <u>106</u> :509 (1989)   |
| 20 | 6 | Pretolani, et al., <i>J. Exp. Med.</i> , <u>180</u> :795 (1994)   |
|    | 7 | Abraham, et al., <i>J. Clin. Invest.</i> , <u>93</u> :776 (1994)  |
| 25 | 8 | Mulligan, et al., <i>J. Immunology</i> , <u>150</u> :2407 (1993)  |

- 9 Cybulsky, et al., *Science*, 251:788 (1991)
- 10 Li, et al., *Arterioscler. Thromb.*, 13:197 (1993)
- 5 11 Sasseville, et al., *Am. J. Path.*, 144:27 (1994)
- 12 Yang, et al., *Proc. Nat. Acad. Science (USA)*, 90:10494 (1993)
- 13 Burkly, et al., *Diabetes*, 43:529 (1994)
- 10 14 Baron, et al., *J. Clin. Invest.*, 93:1700 (1994)
- 15 15 Hamann, et al., *J. Immunology*, 152:3238 (1994)
- 16 Yednock, et al., *Nature*, 356:63 (1992)
- 17 Baron, et al., *J. Exp. Med.*, 177:57 (1993)
- 18 van Dinther-Janssen, et al., *J. Immunology*, 147:4207 (1991)
- 20 19 van Dinther-Janssen, et al., *Annals. Rheumatic Dis.*, 52:672 (1993)
- 20 20 Elices, et al., *J. Clin. Invest.*, 93:405 (1994)
- 21 Postigo, et al., *J. Clin. Invest.*, 89:1445 (1991)
- 22 Paul, et al., *Transpl. Proceed.*, 25:813 (1993)
- 30 23 Okarhara, et al., *Can. Res.*, 54:3233 (1994)
- 24 Paavonen, et al., *Int. J. Can.*, 58:298 (1994)
- 25 25 Schadendorf, et al., *J. Path.*, 170:429 (1993)
- 35 26 Bao, et al., *Diff.*, 52:239 (1993)
- 27 Lauri, et al., *British J. Cancer*, 68:862 (1993)
- 40 28 Kawaguchi, et al., *Japanese J. Cancer Res.*, 83:1304 (1992)
- 29 29 Kogan, et al., *U.S. Patent No. 5,510,332*, issued April 23, 1996
- 45 30 International Patent Appl. Publication No. WO 96/01644

All of the above publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

5

#### State of the Art

VLA-4 (also referred to as  $\alpha 4\beta 1$  integrin and CD49d/CD29), first identified by Hemler and Takada<sup>1</sup>, is a member of the  $\beta 1$  integrin family of cell surface receptors, each of which comprises two subunits, an  $\alpha$  chain and a  $\beta$  chain. VLA-4 contains an  $\alpha 4$  chain and a  $\beta 1$  chain. There are at least nine  $\beta 1$  integrins, all sharing the same  $\beta 1$  chain and each having a distinct  $\alpha$  chain. These nine receptors all bind a different complement of the various cell matrix molecules, such as fibronectin, laminin, and collagen. VLA-4, for example, binds to fibronectin. VLA-4 also binds non-matrix molecules that are expressed by endothelial and other cells. These non-matrix molecules include VCAM-1, which is expressed on cytokine-activated human umbilical vein endothelial cells in culture. Distinct epitopes of VLA-4 are responsible for the fibronectin and VCAM-1 binding activities and each activity has been shown to be inhibited independently.<sup>2</sup>

10  
15  
20

Intercellular adhesion mediated by VLA-4 and other cell surface receptors is associated with a number of inflammatory responses. At the site of an injury or other inflammatory stimulus, activated vascular endothelial cells express molecules that are adhesive for leukocytes. The mechanics of leukocyte adhesion to endothelial cells involves, in part, the recognition and binding of cell surface receptors on leukocytes to the corresponding cell surface molecules on endothelial cells. Once bound, the leukocytes migrate across the blood vessel wall to enter the injured site and release chemical mediators to combat infection. For reviews of adhesion receptors of the immune system, *see*, for example, Springer<sup>3</sup> and Osborn<sup>4</sup>.

25  
30


Inflammatory brain disorders, such as experimental autoimmune encephalomyelitis (EAE), multiple sclerosis (MS) and meningitis, are examples of central nervous system disorders in which the endothelium/leukocyte adhesion mechanism results in destruction to otherwise healthy brain tissue. Large numbers of leukocytes migrate across the blood brain barrier (BBB) in subjects with these inflammatory diseases. The leukocytes release toxic mediators that cause extensive tissue damage resulting in impaired nerve conduction and paralysis.

In other organ systems, tissue damage also occurs via an adhesion mechanism resulting in migration or activation of leukocytes. For example, it has been shown that the initial insult following myocardial ischemia to heart tissue can be further complicated by leukocyte entry to the injured tissue causing still further insult (Vedder et al.<sup>5</sup>). Other inflammatory conditions mediated by an adhesion mechanism include, by way of example, asthma<sup>6-8</sup>, Alzheimer's disease, atherosclerosis<sup>9-10</sup>, AIDS dementia<sup>11</sup>, diabetes<sup>12-14</sup> (including acute juvenile onset diabetes), inflammatory bowel disease<sup>15</sup> (including ulcerative colitis and Crohn's disease), multiple sclerosis<sup>16-17</sup>, rheumatoid arthritis<sup>18-21</sup>, tissue transplantation<sup>22</sup>, tumor metastasis<sup>23-28</sup>, meningitis, encephalitis, stroke, and other cerebral traumas, nephritis, retinitis, atopic dermatitis, psoriasis, myocardial ischemia and acute leukocyte-mediated lung injury such as that which occurs in adult respiratory distress syndrome.

In view of the above, assays for determining the level VLA-4 in a biological sample containing VLA-4 would be useful, for example, to diagnosis VLA-4 mediated conditions. Additionally, despite these advances in the understanding of leukocyte adhesion, the art has only recently addressed the use of inhibitors of adhesion in the treatment of inflammatory brain diseases and other inflammatory conditions<sup>29,30</sup>. The present invention addresses these and other needs.

## SUMMARY OF THE INVENTION

This invention provides compounds which bind to VLA-4. Such compounds can be used, for example, to assay for the presence of VLA-4 in a sample and, in pharmaceutical compositions to inhibit cellular adhesion mediated by VLA-4, for example, binding of VCAM-1 to VLA-4. The compounds of this invention have a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about 15  $\mu$ M or less (as measured by Example 136 below) which compounds are defined by formula I below:



where

$R^1$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

$R^2$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and where  $R^1$  and  $R^2$  together with the nitrogen atom bound to  $R^2$  and the  $\text{SO}_2$  group bound to  $R^1$  form a heterocyclic or a substituted heterocyclic group;

$R^3$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and, when  $R^2$  does not form a heterocyclic group with  $R^1$ ,  $R^2$  and  $R^3$  together with the nitrogen atom bound to  $R^2$  and the carbon atom bound to  $R^3$  can form a heterocyclic or a substituted heterocyclic group;

$R^5$  is  $-(\text{CH}_2)_x\text{-Ar-R}^5$  where  $R^5$  is selected from the group consisting  
(a) substituted alkylcarbonylamino with the proviso that at least one of the substituents on the substituted alkyl moiety is selected from the group

consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl,

5 carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,

10 thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl,

15 -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl,

20 -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-

25 (substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl,

30 heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted

alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(b) alkoxyaryl substituted on the alkoxy moiety with a substituent selected from the group consisting of carboxyl and -COOR where R is alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

(c) aryl and heteroaryl;

(d) -NR'R' wherein each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic with the proviso that at least one of R' is substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic and with the further proviso that when R' is substituted alkyl at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl,



-OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted  
 heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl,  
 -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl,  
 -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-  
 5 heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl,  
 -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted  
 aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl,  
 -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is  
 10 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted  
 alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino,  
 mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino,  
 mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic  
 amino, and unsymmetric di-substituted amines having different substituents  
 15 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,  
 substituted heteroaryl, heterocyclic and substituted heterocyclic, and  
 substituted alkyl groups having amino groups blocked by conventional  
 blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted  
 alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl,  
 -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl,  
 20 -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl,  
 -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is  
 hydrogen or alkyl;

(e) -alkoxy-NR"R" wherein each R" is independently selected from  
 the group consisting of alkyl, substituted alkyl, aryl, substituted aryl,  
 25 cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl,  
 heterocyclic, and substituted heterocyclic with the proviso that when each R"  
 is substituted alkyl then at least one of the substituents on the substituted alkyl  
 moiety is selected from the group consisting of alkoxy, substituted alkoxy,  
 acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl  
 30 amidino, thioamidino, aminoacyl, aminocarbonylamino,  
 aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy,  
 cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted

alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,

5 oxy-carbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl,

10 -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is

15 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents

20 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-

25  
30

-- 10 --

heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

- (f) substituted alkenyl or substituted alkynyl with the proviso that at least one of the substituents on the substituted alkenyl/alkynyl moiety is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when substituted with substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl,

-- 11 --

-NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(g) substituted aryloxy and substituted heteroaryloxy with the proviso that at least one substituent on the substituted aryloxy/heteroaryloxy is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxy carbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(h) -alkoxy-saturated heterocyclic, -alkoxy-saturated substituted heterocyclic, -substituted alkoxy-heterocyclic and -substituted alkoxy-substituted saturated heterocyclic;

(i) -O-heterocyclic and -O-substituted heterocyclic;

(j) tetrazolyl;

(k) -NR-SO<sub>2</sub>-substituted alkyl where R is hydrogen, alkyl or aryl, with the proviso that at least one substituent on the alkyl moiety of the substituted alkylsulfonylamino is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino,

alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(l) alkenylsulfonylamino, alkynylsulfonylamino, substituted alkenylsulfonylamino and substituted alkynylsulfonylamino;

5 (m) substituted alkoxy with the proviso that the substitution on the alkyl moiety of said substituted alkoxy does not include alkoxy-NR<sup>7</sup>, unsaturated heterocyclyl, alkyloxy, aryloxy, heteroaryloxy, aryl, heteroaryl and aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, 10 alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(n) amidine and amidine substituted with from 1 to 3 substituents independently selected from alkyl, substituted alkyl, alkenyl, substituted 15 alkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

(o) -C(O)NR<sup>8</sup>R<sup>9</sup> where each R<sup>8</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted 20 heterocyclic with the proviso that when one R<sup>8</sup> is unsaturated heterocyclylalkyl, aryl, heteroaryl or aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, 25 alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea, then the other R<sup>8</sup> is alkyl, substituted alkyl (other than unsaturated heterocyclyl substituted-alkyl), cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl and heterocyclic and substituted heterocyclic;

30 (p) -NR<sup>12</sup>C(O)-R<sup>8</sup> where R<sup>8</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted

heterocyclic, and  $R^{12}$  is alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5 (q)  $-\text{SO}_2\text{-aryl}$ ,  $-\text{SO}_2\text{-substituted aryl}$ ,  $-\text{SO}_2\text{-heteroaryl}$ ,  $-\text{SO}_2\text{-substituted heteroaryl}$  or  $-\text{SO}_2\text{-alkyl}$ ;

(r)  $-\text{NR}'\text{C(O)NR}^9\text{R}^9$  wherein  $R'$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and each  $R^9$  is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic:

15 (s)  $-\text{NR}'\text{C(O)OR}^9$  wherein  $R'$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and,  $R^9$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(t)  $-\text{aminocarbonyl-(N-formylheterocycl)}$ ; and

20 (u)  $-\text{alkyl-C(O)NH-heterocycl}$  and  $-\text{alkyl-C(O)NH-substituted heterocycl}$ ;

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl,


x is an integer of from 1 to 4;

25 Q is  $-\text{C(X)NR}^7$  wherein  $R^7$  is selected from the group consisting of hydrogen and alkyl; and X is selected from the group consisting of oxygen and sulfur;

and pharmaceutically acceptable salts thereof

30 In another embodiment, the compounds of this invention can also be provided as prodrugs which convert (e.g., hydrolyze, metabolize, etc.) *in vivo* to a compound of formula I above. In a preferred example of such an embodiment, the carboxylic acid of the compound of formula I is modified

into a group which, *in vivo*, will convert to a carboxylic acid (including salts thereof). In a particularly preferred embodiment, such prodrugs are represented by compounds of formula IA:



where

R<sup>1</sup> is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and R<sup>1</sup> and R<sup>2</sup> together with the nitrogen atom bound to R<sup>2</sup> and the SO<sub>2</sub> group bound to R<sup>1</sup> can form a heterocyclic or a substituted heterocyclic group;

R<sup>3</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and, when R<sup>2</sup> does not form a heterocyclic group with R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> and the carbon atom bound to R<sup>3</sup> can form a heterocyclic or a substituted heterocyclic group;

R<sup>5</sup> is -(CH<sub>2</sub>)<sub>i</sub>-Ar-R<sup>5'</sup> where R<sup>5'</sup> is selected from the group consisting

(a) substituted alkylcarbonylamino with the proviso that at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted

cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,

5 thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl,

10 -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl,

15 -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-

20 (substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl,

25 heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

30



-- 16 --

(b) alkoxyaryl substituted on the alkoxy moiety with a substituent selected from the group consisting of carboxyl and -COOR where R is alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

(c) aryl and heteroaryl;

5 (d) -NR'R' wherein each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic with the proviso that at least one of R' is substituted  
10 alkyl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic and with the further proviso that when R' is substituted alkyl at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino,  
15 aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl,  
20 substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted  
25 heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl,  
30 -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl,

-NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(e) -alkoxy-NR"R" wherein each R" is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when each R" is substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted

thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,

5 oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-}$  substituted alkyl,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,

10  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is

15 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents

20 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with  $-\text{SO}_2\text{-alkyl}$ ,  $-\text{SO}_2\text{-substituted alkyl}$ ,  $-\text{SO}_2\text{-alkenyl}$ ,

25  $-\text{SO}_2\text{-substituted alkenyl}$ ,  $-\text{SO}_2\text{-cycloalkyl}$ ,  $-\text{SO}_2\text{-substituted cycloalkyl}$ ,  $-\text{SO}_2\text{-aryl}$ ,  $-\text{SO}_2\text{-substituted aryl}$ ,  $-\text{SO}_2\text{-heteroaryl}$ ,  $-\text{SO}_2\text{-substituted heteroaryl}$ ,  $-\text{SO}_2\text{-heterocyclic}$ ,  $-\text{SO}_2\text{-substituted heterocyclic}$  and  $-\text{SO}_2\text{NRR}$  where R is hydrogen or alkyl;

(f) substituted alkenyl or substituted alkynyl with the proviso that at

30 least one of the substituents on the substituted alkenyl/alkynyl moiety is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl,

heterocyclic, and substituted heterocyclic with the proviso that when substituted with substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,

-- 20 --

substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(g) substituted aryloxy and substituted heteroaryloxy with the proviso that at least one substituent on the substituted aryloxy/heteroaryloxy is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(h) -alkoxy-saturated heterocyclic, -alkoxy-saturated substituted heterocyclic, -substituted alkoxy-heterocyclic and -substituted alkoxy-substituted saturated heterocyclic;

(i) -O-heterocyclic and -O-substituted heterocyclic;

(j) tetrazolyl;

(k) -NR-SO<sub>2</sub>-substituted alkyl where R is hydrogen, alkyl or aryl, with the proviso that at least one substituent on the alkyl moiety of the substituted alkylsulfonylamino is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(l) alkenylsulfonylamino, alkynylsulfonylamino, substituted alkenylsulfonylamino and substituted alkynylsulfonylamino;

(m) substituted alkoxy with the proviso that the substitution on the alkyl moiety of said substituted alkoxy does not include alkoxy-NR"R", unsaturated heterocyclyl, alkyloxy, aryloxy, heteroaryloxy, aryl, heteroaryl

and aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(n) amidine and amidine substituted with from 1 to 3 substituents independently selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

(o)  $-C(O)NR''R''$  where each  $R''$  is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic with the proviso that when one  $R''$  is unsaturated heterocyclalkyl, aryl, heteroaryl or aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea, then the other  $R''$  is alkyl, substituted alkyl (other than unsaturated heterocyclalkyl substituted-alkyl), cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl and heterocyclic and substituted heterocyclic;

(p)  $-NR^{12}C(O)-R^8$  where  $R^8$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and  $R^{12}$  is alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(q)  $-SO_2$ -aryl,  $-SO_2$ -substituted aryl,  $-SO_2$ -heteroaryl,  $-SO_2$ -substituted heteroaryl or  $-SO_2$ -alkyl;

(r)  $-NR'C(O)NR^9R^9$  wherein  $R'$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted

cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and each  $R^9$  is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(s)  $-NR^1C(O)OR^9$  wherein  $R^1$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and,  $R^9$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(t) -aminocarbonyl-(N-formylheterocycl); and

(u) -alkyl-C(O)NH-heterocycl and -alkyl-C(O)NH-substituted heterocycl,

$Ar$  is aryl, heteroaryl, substituted aryl or substituted heteroaryl,  
 $x$  is an integer of from 1 to 4;

$R^6$  is selected from the group consisting of 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), amino, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, -O-(N-succinimidyl), -NH-adamantyl, -O-cholest-5-en-3- $\beta$ -yl, -NHOY where Y is hydrogen, alkyl, substituted alkyl, aryl, and substituted aryl,  $-NH(CH_2)_pCOOY$  where  $p$  is an integer of from 1 to 8 and Y is as defined above,  $-OCH_2NR^9R^{10}$  where  $R^9$  is selected from the group consisting of -C(O)-aryl and -C(O)-substituted aryl and  $R^{10}$  is selected from the group consisting of hydrogen and  $-CH_2COOR^{11}$  where  $R^{11}$  is alkyl, and  $-NHSO_2Z$  where Z is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

Q is  $-C(X)NR^7-$  wherein  $R^7$  is selected from the group consisting of hydrogen and alkyl; and X is selected from the group consisting of oxygen and sulfur;

and pharmaceutically acceptable salts thereof

with the proviso that when  $R^1$  is  $p\text{-CH}_3\text{-}\phi\text{-}$ ,  $R^6$  is methoxy, Q is

-C(O)NH-, and R<sup>2</sup> and R<sup>3</sup> are joined to form a pyrrolidinyl group, then R<sup>5</sup> is not *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(isopropyl)<sub>2</sub>]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-1-pyrrolidinyl]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-pyrimidinyl)piperazinyl]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-N-morpholinyl]-benzyl-, or *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-N-piperidinyl]-benzyl-.

Preferably, in the compounds of formula I and IA above, R<sup>1</sup> is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl.

Even more preferably R<sup>1</sup> is selected from the group consisting of 4-methylphenyl, methyl, benzyl, *n*-butyl, 4-chlorophenyl, 1-naphthyl, 2-naphthyl, 4-methoxyphenyl, phenyl, 2,4,6-trimethylphenyl, 2-(methoxycarbonyl)phenyl, 2-carboxyphenyl, 3,5-dichlorophenyl, 4-trifluoromethylphenyl, 3,4-dichlorophenyl, 3,4-dimethoxyphenyl, 4-(CH<sub>3</sub>C(O)NH-)phenyl, 4-trifluoromethoxyphenyl, 4-cyanophenyl, isopropyl, 3,5-di-(trifluoromethyl)phenyl, 4-*t*-butylphenyl, 4-*t*-butoxyphenyl, 4-nitrophenyl, 2-thienyl, 1-N-methyl-3-methyl-5-chloropyrazol-4-yl, phenethyl, 1-N-methylimidazol-4-yl, 4-bromophenyl, 4-amidinophenyl, 4-methylamidinophenyl, 4-[CH<sub>3</sub>SC(=NH)]phenyl, 5-chloro-2-thienyl, 2,5-dichloro-4-thienyl, 1-N-methyl-4-pyrazolyl, 2-thiazolyl, 5-methyl-1,3,4-thiadiazol-2-yl, 4-[H<sub>2</sub>NC(S)]phenyl, 4-aminophenyl, 4-fluorophenyl, 2-fluorophenyl, 3-fluorophenyl, 3,5-difluorophenyl, pyridin-3-yl, pyrimidin-2-yl, 4-(3'-dimethylamino-*n*-propoxy)-phenyl, and 1-methylpyrazol-4-yl.

Preferably, in the compounds of formula I and IA above, R<sup>2</sup> is hydrogen, methyl, phenyl, benzyl, -(CH<sub>2</sub>)<sub>2</sub>-2-thienyl, and -(CH<sub>2</sub>)<sub>2</sub>-φ.

In one embodiment, R<sup>1</sup> and R<sup>2</sup> together with the nitrogen atom bound to R<sup>2</sup> and the SO<sub>2</sub> group bound to R<sup>1</sup> are joined to form a heterocyclic group or substituted heterocyclic group. Preferred heterocyclic and substituted heterocyclic groups include those having from 5 to 7 ring atoms having 2 to 3 heteroatoms in the ring selected from nitrogen, oxygen and sulfur which ring



is optionally fused to another ring such as a phenyl or cyclohexyl ring to provide for a fused ring heterocycle of from 10 to 14 ring atoms having 2 to 4 heteroatoms in the ring selected from nitrogen, oxygen and sulfur.

Specifically preferred R<sup>1</sup>/R<sup>2</sup> joined groups include, by way of example,  
5 benzisothiazolonyl (saccharin-2-yl).

In one preferred embodiment, R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> substituent and the carbon bound to the R<sup>3</sup> substituent form a heterocyclic group or a substituted heterocyclic group of 4 to 6 ring atoms  
10 having 1 to 2 heteroatoms in the ring selected from nitrogen, oxygen and sulfur which ring is optionally substituted with 1 to 2 substituents selected from fluoro, methyl, hydroxy, amino, phenyl, thiophenyl, thiobenzyl or can be fused to another ring such as a phenyl or cycloalkyl ring to provide for a fused ring heterocycle of from 10 to 14 ring atoms having 1 to 2 heteroatoms in the  
15 ring selected from nitrogen, oxygen and sulfur. Such heterocyclic rings include azetidiny (e.g., L-azetidiny), thiazolidiny (e.g., L-thiazolidiny), piperidiny (e.g., L-piperidiny), piperiziny (e.g., L-piperiziny), dihydroindolyl (e.g., L-2,3-dihydroindol-2-yl), tetrahydroquinoliny (e.g., L-1,2,3,4-tetrahydroquinolin-2-yl), thiomorpholiny (e.g., L-thiomorpholin-3-yl), pyrrolidiny (e.g., L-pyrrolidiny), substituted pyrrolidiny such as 4-  
20 hydroxypyrrolidiny (e.g., 4- $\alpha$ -(or  $\beta$ )-hydroxy-L-pyrrolidiny), 4-fluoropyrrolidiny (e.g., 4- $\alpha$ -(or  $\beta$ )-fluoro-L-pyrrolidiny), 3-phenylpyrrolidiny (e.g., 3- $\alpha$ -(or  $\beta$ )-phenyl-L-pyrrolidiny), 3-thiophenylpyrrolidiny (e.g., 3- $\alpha$ -(or  $\beta$ )-thiophenyl-L-pyrrolidiny), 4-aminopyrrolidiny (e.g., 4- $\alpha$ -(or  $\beta$ )-amino-L-pyrrolidiny), 3-methoxypyrrolidiny (e.g., 3- $\alpha$ -(or  $\beta$ )-methoxy-L-pyrrolidiny), 4,4-di-methylpyrrolidiny, substituted piperiziny such as 4-N-Cbz-piperiziny, substituted thiazolidiny such as 5,5-dimethylthiazolidin-4-yl, 1,1-dioxo-thiazolidiny (e.g., L-1,1-dioxo-thiazolidin-2-yl), substituted 1,1-dioxo-thiazolidiny such as L-1,1-dioxo-5,5-dimethylthiazolidin-2-yl, 1,1-dioxothiophenoliny (e.g., L-1,1-dioxo-thiomorpholin-3-yl) and the like.  
30

Preferably, in the compounds of formula I and IA above, R<sup>3</sup> includes all of the isomers arising by substitution with methyl, phenyl, benzyl, diphenylmethyl, -CH<sub>2</sub>CH<sub>2</sub>-COOH, -CH<sub>2</sub>-COOH, 2-amidoethyl, *iso*-butyl, *t*-butyl, -CH<sub>2</sub>O-benzyl and hydroxymethyl. Additionally, in another preferred embodiment, R<sup>1</sup> and R<sup>2</sup> together with the nitrogen atom bound to R<sup>2</sup> can form a heterocyclic group or substituted heterocyclic group.

Q is preferably -C(O)NH- or -C(S)NH-.

10

R<sup>5</sup> is preferably selected from all possible isomers arising by substitution with the following groups:

4-[NH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-[HOOCCH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>NHBoc]benzyl, 4-[-NHC(O)CH(CH<sub>3</sub>)NHBoc]benzyl,  
 15 4-[-NHC(O)CH(CH<sub>2</sub>φ)NHBoc]benzyl, 4-[-NHC(O)CH<sub>2</sub>NHC(O)NH-3'-methylphenyl]benzyl, 4-[-NHC(O)CH(NHBoc)(CH<sub>2</sub>)<sub>4</sub>NHCbz]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>CH(C(O)OCH<sub>2</sub>φ)-NHCbz]benzyl, 4-φ-benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)NHBoc]benzyl,  
 4-[H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(BocHNCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-C(O)NH-)benzyl, 4-[φCH<sub>2</sub>OCH<sub>2</sub>(BocHN)CHC(O)NH-]benzyl,  
 20 4-[CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(N-methylpiperidin-4-oxy)-benzyl,  
 4-[CH<sub>2</sub>N(Boc)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[φCH<sub>2</sub>OCH<sub>2</sub>(H<sub>2</sub>N)CHC(O)NH-]benzyl, 4-[HO(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl, 4-[φCH<sub>2</sub>O(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 25 4-[HO(O)C(NH<sub>2</sub>)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 4-[CH<sub>3</sub>(*N*-Boc)NCH<sub>2</sub>C(O)NH-]benzyl, 4-[CH<sub>3</sub>NHCH<sub>2</sub>C(O)NH-]benzyl,  
 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>C(O)NH-]benzyl, 4-[-O-CH(COOH)φ]benzyl,  
 4-[2-carboxylphenyl]-benzyl, 4-[2-carboxylmethylphenyl]-benzyl  
 4-[φCH<sub>2</sub>OC(O)NHCH<sub>2</sub>CH<sub>2</sub>NH-]benzyl, 4-N[-(SO<sub>2</sub>)CH<sub>3</sub>]-  
 30 CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]benzyl, 4-*t*-butyl-O(O)CCH<sub>2</sub>-O-benzylNH]benzyl,  
 4-[*N,N*-di(4-*N,N*-dimethylamino)benzyl]amino]benzyl,  
 4-(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl-CH<sub>2</sub>NH)benzyl,

- 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1'-(4'-pyrimidinyl)-piperazinyl]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-pyrrolidinyl)]-  
 benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)]-benzyl,  
 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-O]-benzyl, 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>O]-benzyl,  
 5 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(1'-(4'-methylpiperazinyl))]benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-4-(3'-chlorophenyl)-piperazin-1-yl]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(φ)CH<sub>2</sub>CH<sub>3</sub>]-benzyl, 4-[-OCH<sub>2</sub>-3'-(N-Boc)-piperidinyl]-  
 benzyl, 4-[-O-(3-(N-Boc)-piperidinyl)]benzyl,  
 3-[-O-(N-methylpiperidin-4-yl)]benzyl, 4-[-O-(N-methylpiperidin-4-yl)]benzyl,  
 10 4-[di-*iso*-propylamino-CH<sub>2</sub>CH<sub>2</sub>O]-benzyl, 4-[*N*-3-methylbutyl-*N*-trifluoro-  
 methanesulfonyl]amino]benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-morpholinyl)]-benzyl,  
 4-[-OCH<sub>2</sub>CH(NHBoc)CH<sub>2</sub>cyclohexyl]-benzyl, 4-[OCH<sub>2</sub>CH<sub>2</sub>-(N'-piperidinyl)]-  
 benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(4-*m*-chlorophenyl)-piperazin-1-yl]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-homopiperidinyl)-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(benzyl)<sub>2</sub>]-benzyl,  
 15 3-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)benzyl]-benzyl, 4-[2-(2-azabicyclo[3.2.2]octan-2-  
 yl)ethyl-O]-benzyl, [cyclopentylacetylenyl]-benzyl, 4-[-C≡C-φ-4'φ]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-S(O)<sub>2</sub>-4'-CH<sub>3</sub>-φ]-benzyl, 4-[-C≡C-CH<sub>2</sub>NHC(O)NH<sub>2</sub>]-benzyl,  
 20 4-[-C≡C-CH<sub>2</sub>-O-(4'-COOCH<sub>2</sub>CH<sub>3</sub>)φ]-benzyl, 4-[-C≡C-CH(NH<sub>2</sub>)-cyclohexyl]-  
 benzyl, 4-[-C≡C-CH<sub>2</sub>-O-phenyl]-benzyl, 4-[-C≡C-CH<sub>2</sub>OCH<sub>3</sub>]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-(4'-C(O)OC<sub>2</sub>H<sub>5</sub>)phenyl]-benzyl, 4-[-C≡C-  
 CH<sub>2</sub>CH(C(O)OCH<sub>3</sub>)<sub>2</sub>]-benzyl, 4-[-C≡C-CH<sub>2</sub>CH(NHC(O)CH<sub>3</sub>)C(O)OH]-  
 benzyl, 4-[-C≡C-CH<sub>2</sub>NH-(4,5-dihydro-4-oxo-5-phenyl-oxazol-2yl)]-benzyl,  
 25 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(N-morpholino)]-benzyl, 4-[-OCH<sub>2</sub>COOH]-benzyl,  
 4-[-OCH<sub>2</sub>COO-*t*-butyl]-benzyl, 4-[-N(SO<sub>2</sub>CH<sub>3</sub>)(CH<sub>2</sub>)<sub>3</sub>-N(CH<sub>3</sub>)<sub>2</sub>]-benzyl,  
 4-[-NHS(O)<sub>2</sub>CF<sub>3</sub>]-benzyl, 4-[-C(=NH)NH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>Cl]-  
 benzyl, 4-[-OCH<sub>2</sub>C(O)NH-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)O-benzyl]-benzyl,  
 4-[-OCH<sub>2</sub>C(O)OH]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-(3-methoxypyrrol-  
 30 2-yl)-piperazinyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-*t*-  
 butyl]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-phenyl)-piperidinyl]-benzyl,

4-[-NHSO<sub>2</sub>-CH=CH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>Cl]-benzyl, 4-benzyl-  
benzyl, 4-[-OCH<sub>2</sub>C(O)piperidin-1-yl]benzyl,  
4-[-OCH<sub>2</sub>C(O)N(CH(CH<sub>3</sub>)<sub>2</sub>)<sub>2</sub>]benzyl, 4-amidinobenzyl, 4-acetamidobenzyl,  
4-(N-methyl)acetamidobenzyl, 4-(NHC(O)CH<sub>2</sub>NHC(O)NH-fluorescein)benzyl,  
5 4-(NHC(O)CH<sub>2</sub>CH(NH<sub>2</sub>)COOH, (1-toluenesulfonylimidizol-4-yl)-methyl-,  
[(1-N,N-dimethylaminosulfonyl)-imidizol-4-yl]methyl-, 4-(N-toluenesulfonyl-  
amino)benzyl, and 4-[N-methyltrifluoroacetamido]phenyl.

In the compounds of formula IA, R<sup>6</sup> is preferably 2,4-dioxo-  
10 tetrahydrofuran-3-yl (3,4-enol), methoxy, ethoxy, *iso*-propoxy, *n*-butoxy,  
*t*-butoxy, cyclopentoxy, *neo*-pentoxy, 2- $\alpha$ -*iso*-propyl-4- $\beta$ -methylcyclohexoxy,  
2- $\beta$ -isopropyl-4- $\beta$ -methylcyclohexoxy, -NH<sub>2</sub>, benzyloxy, -NHCH<sub>2</sub>COOH,  
-NHCH<sub>2</sub>CH<sub>2</sub>COOH, -NH-adamantyl, -NHCH<sub>2</sub>CH<sub>2</sub>COOCH<sub>2</sub>CH<sub>3</sub>, -NHSO<sub>2</sub>-*p*-  
CH<sub>3</sub>- $\phi$ , -NHOR<sup>8</sup> where R<sup>8</sup> is hydrogen, methyl, *iso*-propyl or benzyl, O-(N-  
15 succinimidyl), -O-cholest-5-en-3- $\beta$ -yl, -OCH<sub>2</sub>-OC(O)C(CH<sub>3</sub>)<sub>3</sub>,  
-O(CH<sub>2</sub>)<sub>z</sub>NHC(O)W where z is 1 or 2 and W is selected from the group  
consisting of pyrid-3-yl, N-methylpyridyl, and N-methyl-1,4-dihydro-pyrid-3-  
yl, -NR''C(O)-R' where R' is aryl, heteroaryl or heterocyclic and R'' is  
hydrogen or -CH<sub>2</sub>C(O)OCH<sub>2</sub>CH<sub>3</sub>.

Preferred compounds within the scope of formula I and IA above  
include by way of example:

25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxyl-  
carbonyl)glycyl]amino]-L-phenylalanine  
*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(glycyl)amino]-L-phenylalanine  
*N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(carboxy)propionamido]-L-  
30 phenylalanine  
*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxylcarbonyl)-L-  
alanyl]amino]-L-phenylalanine  
35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxylcarbonyl)-D-  
alanyl]amino]-L-phenylalanine

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl-D-phenylalanyl)amino]-L-phenylalanine
- 5      *N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[3-(fluorescein)thiouriedo]acetamido}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)glycyl]amino]-L-phenylalanine methyl ester
- 10      *N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[3-(3-methylphenyl)uriedo]acetamido}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*α-*tert*-butoxycarbonyl-*N*ε-carbobenzyloxy-L-lysyl)amino]-L-phenylalanine
- 15      *N*-(toluene-4-sulfonyl)-L-prolyl-4-[γ-(α-benzyl-*N*α-carbobenzyloxy-L-aspartyl)amino]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*α-*tert*-butoxycarbonyl-L-lysyl)amino]-L-phenylalanine
- 20      *N*-(toluene-4-sulfonyl)-L-prolyl-4-[γ-(L-aspartyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(4-aminobutyramido)-L-phenylalanine
- 25      *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-*tert*-butoxycarbonylamino)butyramido]-L-phenylalanine
- 30      *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-methylamino)butyramido]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-*tert*-butoxycarbonyl-*N*-methylamino)butyramido]-L-phenylalanine
- 35      *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*O*-benzyl)-L-seryl]amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[δ-(D,L-glutamyl)amino]-L-phenylalanine
- 40      *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonylsarcosyl)amino]-L-phenylalanine
- 45      *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N*-*tert*-butoxycarbonylsarcosyl)amino]-L-phenylalanine

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine ethyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(sarcosyl)amino]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-( $\alpha$ -carboxybenzyloxy)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenyl]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(methoxycarbonyl)phenyl]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[3-(*N,N*-dimethylamino)propyl]-*N*-[trifluoromethanesulfonyl]amino}-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[4-[(*tert*-butoxycarbonyl)methoxy]benzyl]amino}-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N,N*-di[4-(*N,N*-dimethylamino)benzyl]amino}-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl)methyl]amino}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-*N*-methyl-L-serinyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester

- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N*-ethyl-*N*-phenylamino)ethoxy]-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-diisopropylamino)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-cyclohexyl-2-(*N*-tert-butoxycarbonylamino)propoxy]-L-phenylalanine methyl ester
- 15 *N*-(thiophene-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(5-chlorothiophene-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-diethylamino)ethoxy]-L-phenylalanine
- N*-(2,5-dichlorothiophene-3-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 25 *N*-(1-methylpyrazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-diethylamino)propoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-3-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 35 *N*-(thiazole-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-diethylamino)propoxy]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine methyl ester

- N*-(1-methylimidazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 5 *N*-(2-methylthiadiazole-5-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 10 *N*-(4-cyanobenzenesulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-(3,3-dimethyl)prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine ethyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-(thiamorpholin-3-carbonyl)-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(cyclopentyl)ethynyl]-D,L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-{2-[4-(phenyl)phenyl]ethynyl}-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(toluene-4-sulfonyloxy)prop-1-ynyl]-D,L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(ureido)prop-1-ynyl]-D,L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(1-aminocyclohex-1-yl)ethynyl]-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(phenoxy)prop-1-ynyl]-D,L-phenylalanine



- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(phenoxy)prop-1-ynyl]-D,L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(methoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(methoxy)prop-1-ynyl]-D,L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4,4-di(methoxycarbonyl)but-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[4,4-di(methoxycarbonyl)but-1-ynyl]-D,L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(4-acetamido-4-carboxybut-1-ynyl)-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-(4-acetamido-4-carboxybut-1-ynyl)-D,L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[(4,5-dihydro-4-oxo-5-phenyloxazol-2-yl)amino]prop-1-ynyl}-D,L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)sarcosyl-4-{3-[(4,5-dihydro-4-oxo-5-phenyloxazol-2-yl)amino]prop-1-ynyl}-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenoxy]-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[4-(pyrimidin-2-yl)piperazin-1-yl]ethoxy}-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine methyl ester

-- 33 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(piperidin-1-yl)ethoxy]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine methyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[4-(3-methoxythien-2-yl)-4-hydroxypiperidin-1-yl]ethoxy}-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-3-(1-methylpiperidin-4-oxy)-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(1-methylpiperidin-4-oxy)-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-(1-methylpiperidin-4-oxy)-L-phenylalanine ethyl ester

- N*-(toluene-4-sulfonyl)-L-(1,1-dioxothiomorpholin-3-carbonyl)-4-(1-methylpiperidin-4-oxy)-L-phenylalanine ethyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-(1,1-dioxothiomorpholin-3-carbonyl)-4-(1-methylpiperidin-4-oxy)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-(1-methylpiperidin-4-oxy)-L-phenylalanine
- 10 *N*-( $\alpha$ -toluenesulfonyl)-L-prolyl-4-(1-methylpiperidin-4-oxy)-L-phenylalanine ethyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(trifluoromethanesulfonyl)amino-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(trifluoromethanesulfonyl)amino-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(chloromethanesulfonyl)amino-L-phenylalanine methyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-trifluoromethanesulfonyl-*N*-isobutyl)amino-L-phenylalanine methyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-benzylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine methyl ester
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(aminocarbonyl)methoxy]-L-phenylalanine

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butylaminocarbonyl)methoxy]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(4-phenyl-4-hydroxypiperidin-1-yl)ethoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(piperidin-1-ylcarbonyl)methoxy]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(amidino)phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(aminocarbonyl)phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine isopropyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methyltrifluoroacetamido)-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine *t*-butyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-(*N*-methylpiperidinoxy)-phenylalanine *t*-butyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-L-(4-methylpiperidinoxy) phenylalanine *t*-butyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine *t*-butyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine
- 45 and pharmaceutically acceptable salts thereof as well as any of the ester compounds recited above wherein one ester is replaced with another

ester selected from the group consisting of methyl ester, ethyl ester, *n*-propyl ester, isopropyl ester, *n*-butyl ester, isobutyl ester, *sec*-butyl ester and *tert*-butyl ester.


5           This invention also provides methods for binding VLA-4 in a biological sample which method comprises contacting the biological sample with a compound of formula I or IA above under conditions wherein said compound binds to VLA-4.

10           Certain of the compounds of formula I and IA above are also useful in reducing VLA-4 mediated inflammation *in vivo*.

          This invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of  
15           one or more the compounds of formula I or IA above with the exception that  $R^3$  and  $R^4$  are derived from L-amino acids or other similarly configured starting materials. Alternatively, racemic mixtures can be used.

          The pharmaceutical compositions may be used to treat VLA-4  
20           mediated disease conditions. Such disease conditions include, by way of example, asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes (including acute juvenile onset diabetis), inflammatory bowel disease (including ulcerative colitis and Crohn's disease), multiple sclerosis, rheumatoid arthritis, tissue transplantation, tumor metastasis, meningitis,  
25           encephalitis, stroke, and other cerebral traumas, nephritis, retinitis, atopic dermatitis, psoriasis, myocardial ischemia and acute leukocyte-mediated lung injury such as that which occurs in adult respiratory distress syndrome. Accordingly, this invention also provides methods for the treatment of an  
          inflammatory disease in a patient mediated by VLA-4 which methods  
30           comprise administering to the patient the pharmaceutical compositions described above.

Preferred compounds of formula I and IA above include those set forth in Table I below:



In all cases Q is -C(O)NH-

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[NH(C(O)CH <sub>2</sub> NHBoc]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[NH(C(O)CH <sub>2</sub> NH <sub>2</sub> ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[NH(C(O)CH <sub>2</sub> CH <sub>2</sub> C(O)OH]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	4-[NH(C(O)CH <sup>*</sup> (CH <sub>2</sub> )NHBoc]- benzyl- ( <sup>*</sup> corresponds to L isomer)	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	4-[NH(C(O)CH <sup>*</sup> (CH <sub>2</sub> )NHBoc]- benzyl- ( <sup>*</sup> corresponds to D isomer)	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>5</sup>	R <sup>6</sup>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		4-[NHC(O)CH <sup>*</sup> (CH <sub>3</sub> , $\phi$ )NHBoc]- benzyl- (* corresponds to D isomer)	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		[NHC(O)CH <sub>2</sub> NHC(O)NH-fluorescin]benzyl	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		4-[BocNHCH <sub>2</sub> C(O)NH]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[NHC(O)CH <sub>2</sub> NHC(O)NH- <i>m</i> - methylphenyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[NHC(O)CH(NHBoc)(CH <sub>3</sub> ),NHCbz]- benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[NHC(O)CH <sub>2</sub> CH-NHCbz]-benzyl   C(O)OCH <sub>3</sub> , $\phi$	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[NHC(O)CH <sub>2</sub> NHBoc]-benzyl   CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[NHC(O)CH <sub>2</sub> CH(NH <sub>2</sub> )COOH	-OH



R <sup>1</sup>	R <sup>2</sup>	R <sup>1</sup>	R <sup>1</sup>	R <sup>6</sup>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(H,NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CO)NH(benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(Boc-1HCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CO)NH(benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(CH <sub>3</sub> NHCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CO)NH(benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(CH <sub>3</sub> (Boc)NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> - CO)NH(benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(φCH <sub>2</sub> OCH <sub>2</sub> (H,N)CHCHCO)NH(benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(HO)C(Cbz-NH)CHCH <sub>2</sub> CH <sub>2</sub> - CO)NH(benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(HO)C(H,N)CHCH <sub>2</sub> CH <sub>2</sub> - CO)NH(benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(CH <sub>3</sub> (N-Boc)NCH <sub>2</sub> CO)NH(benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>1</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -(CH <sub>2</sub> ( <i>N</i> -Boc)NCH <sub>2</sub> C(O)NH-) <i>i</i> -benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(CH <sub>2</sub> NHCH <sub>2</sub> C(O)NH-) <i>i</i> -benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(CH <sub>2</sub> NI(CH <sub>2</sub> C(O)NH-) <i>i</i> -benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -(CH <sub>2</sub> NHCH <sub>2</sub> C(O)NH-) <i>i</i> -benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> C(O)NH-) <i>i</i> -benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> C(O)NH-) <i>i</i> -benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -(O-CH(COOH)φ)-benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -Cl <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[ <i>o</i> -carboxyphenyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[ <i>o</i> -carboxymethylphenyl]-benzyl-	-O-Cl <sub>3</sub> -φ
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[φCH <sub>2</sub> OC(O)NHCH <sub>2</sub> CH <sub>2</sub> NH-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[φCH <sub>2</sub> OC(O)NHCH <sub>2</sub> CH <sub>2</sub> NH-]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[N(CH <sub>3</sub> ) <sub>2</sub> -N(CH <sub>3</sub> ) <sub>2</sub> ]-benzyl-   SO <sub>2</sub> CH <sub>3</sub>	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[( <i>tert</i> -butyl-O(O)CCH <sub>2</sub> -O-benzyl)- NH-]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -Cl <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[( <i>N,N</i> -di(4- <i>N,N</i> - dimethylamino)benzyl)amino]benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -(2-formyl-1,2,3,4-tetrahydroisoquinolin-3- yl-CH <sub>2</sub> NH-)-benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		4-[(CH <sub>3</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-] $\phi$ -CH <sub>2</sub> -	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	-CH <sub>2</sub> OH (L isomer)	<i>p</i> -[(CH <sub>3</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> O-]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = 5,5- dimethylthiazo- lidin-4-yl		<i>p</i> -[(CH <sub>3</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> O-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[(CH <sub>3</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> O-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> N $\phi$ (CH <sub>2</sub> CH <sub>2</sub> )-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[di- <i>iso</i> -propylamino-(CH <sub>2</sub> CH <sub>2</sub> O-)-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> (NHBOC)(CH <sub>2</sub> cyclohexyl)- benzyl	-OCH <sub>3</sub>
2-thienyl-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> )-]-benzyl-	-OH
5-chloro-2- thienyl-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> )-]-benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> )]-benzyl-	-OH
2,5-dichloro- 4-thienyl	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CI <sub>2</sub> )]-benzyl-	-OH
1-N-methyl-4- pyrazolyl	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> )]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> )]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>m</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> )]-benzyl-	-OH
2-thiazolyl	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> )]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CI <sub>2</sub> )]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> )]-benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>3</sup>	R <sup>6</sup>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> )benzyl]-benzyl-	-OCH <sub>3</sub>
1-N-methyl-4- imidazolyl	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub> ]-benzyl-	-OH
2-methyl-4- thiadiazolyl	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub> ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sub>2</sub> /R <sub>3</sub> = thiazol-4-yl		<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub> ]-benzyl-	-OH
<i>p</i> -NC-φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O]benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> -		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O]benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O]benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>

R <sup>1</sup>	R <sup>2</sup>	R <sup>1</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[2-(2-azabicyclo[3.2.2]octan- 2-yl)ethyl-O-]benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> CH <sub>2</sub> -S-CH <sub>2</sub> - (L-thiomorpholin- 3-yl)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -O-]benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -O-]benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -O-]benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[2-(2-azabicyclo[3.2.2]octan- 2-yl)ethyl-O-]benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	H	-CH <sub>2</sub> -φ (L isomer)	<i>p</i> -(cyclopentyl-C≡C-)-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	H	-CH <sub>2</sub> -φ (L isomer)	<i>p</i> -[C≡C-φ- <i>p</i> -φ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	H	-CH <sub>2</sub> -φ (L isomer)	<i>p</i> -[C≡C-CH <sub>2</sub> -O-S(O) <sub>2</sub> - <i>p</i> -Cl <sub>2</sub> -φ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	H	-CH <sub>2</sub> -φ (L isomer)	<i>p</i> -[C≡C-CH <sub>2</sub> NHC(O)NH <sub>2</sub> ]-benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	
<i>p</i> -CH <sub>3</sub> -φ	H	-CH <sub>2</sub> -φ (L isomer)	<i>p</i> -[-C≡C-CH <sub>2</sub> -O- <i>p</i> -COOCH <sub>2</sub> CH <sub>2</sub> -φ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	H	-CH <sub>2</sub> -φ (L isomer)	<i>p</i> -[-C≡C-CH(NH <sub>2</sub> )-cyclohexyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -O-phenyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -O-phenyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -OCH <sub>2</sub> ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -OCH <sub>2</sub> ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -O- <i>p</i> -(-C(O)OC <sub>2</sub> H <sub>5</sub> )-phenyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -O- <i>p</i> -(-C(O)OC <sub>2</sub> H <sub>5</sub> )-phenyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	-CH <sub>3</sub>	H	<i>p</i> -[-C≡C-CH <sub>2</sub> -CH(C(O)OCH <sub>3</sub> )-]-benzyl-	-



R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> CH(C(O)OH)-benzyl-   NIC(O)CH <sub>3</sub> ]	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> NH-(4,5-dihydro-4-oxo- 5-phenyl-oxazol-2-yl)]benzyl	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> NH-(4,5-dihydro-4-oxo- 5-phenyl-oxazol-2-yl)]benzyl	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[O-( <i>o</i> -carboxyphenyl)]-benzyl	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> -1-(4- pyrimidinyl)-piperazinyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> -(1-piperidinyl)]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> -(1-pyrrolidinyl)]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -(1-piperidinyl)]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -1-(4- <i>m</i> -chlorophenyl)-piperazinyl]-benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>p</i> -1-OCH <sub>2</sub> -3-(N-Boc-piperidinyl)-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>p</i> -1-OCH <sub>2</sub> CH <sub>2</sub> -(N-morpholinyl)-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>p</i> -1-OCH <sub>2</sub> CH <sub>2</sub> -(N-piperidinyl)-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>p</i> -1-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -1-(4- <i>m</i> -chlorophenyl)-piperazinyl)-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>p</i> -1-OCH <sub>2</sub> CH <sub>2</sub> -(N-humopiperidinyl)-benzyl	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>p</i> -1-OCH <sub>2</sub> CH <sub>2</sub> -(N-humopiperidinyl)-benzyl	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>p</i> -1-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -1-(4-methyl)-piperazinyl)-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)		<i>m</i> -1-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -(1-pyrrolidiny)))-benzyl-	-OH

R <sup>1</sup>	R <sup>2</sup>	R <sup>1</sup>	R <sup>3</sup>	R <sup>6</sup>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -1-(4-methyl)- piperazinyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> -(N-morpholino)]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> -1-(4-hydroxy-4- (3-methoxy-pyrrol-2-yl)-piperazinyl)- benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[O-(3-(N-Boc)-piperidinyl)-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>m</i> -[O-(N-methyl-piperidin-4-yl)-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[O-(N-methylpiperidin-4-yl)]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> -		<i>p</i> -[(1-methylpiperidin-4-yl)-O]-benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> -CH <sub>2</sub> -SO <sub>2</sub> -CH <sub>2</sub> - (L-1,1-dioxathiomorpholin- 3-yl)		<i>p</i> -[(1-methylpiperidin-4-yl)-O]-benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> CH <sub>2</sub> -SO <sub>2</sub> -CH <sub>2</sub> - (L-1,1- dioxathiomorpholin- 3-yl)		<i>p</i> -[(1-methylpiperidin-4-yl)-O-]benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> CH <sub>2</sub> -SO <sub>2</sub> -CH <sub>2</sub> - (L-1,1- dioxathiomorpholin- 3-yl)		<i>p</i> -[(1-methylpiperidin-4-yl)-O-]benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -[(1-methylpiperidin-4-yl)-O-]benzyl-	-OH
φ-CH <sub>2</sub> -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[(1-methylpiperidin-4-yl)-O-]benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[NHSO <sub>2</sub> CF <sub>3</sub> ]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[NHSO <sub>2</sub> CF <sub>3</sub> ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ-	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[NHSO <sub>2</sub> CH <sub>2</sub> Cl]-benzyl-	-OCH <sub>3</sub>

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[·NHSO <sub>2</sub> -CH=CH <sub>2</sub> ]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[ <i>N</i> -3-methylbutyl- <i>N</i> - trifluoromethanesulfonyl]amino]benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[ <i>N</i> -vinylsulfonyl]amino]benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[·OCH <sub>2</sub> C(O)NH-benzyl]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[·OCH <sub>2</sub> C(O)O-benzyl]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[·OCH <sub>2</sub> C(O)NH-benzyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[·OCH <sub>2</sub> C(O)OH]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -[·OCH <sub>2</sub> C(O)NH <sub>2</sub> ]-benzyl-	-OCH <sub>3</sub>

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-OCH <sub>2</sub> C(O)NH],-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-OCH <sub>2</sub> C(O)NH- <i>t</i> -butyl]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-OCH <sub>2</sub> CH <sub>2</sub> -1-(4-hydroxy-4-phenyl)-piperidiny]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[(piperidin-1-yl)C(O)CH <sub>2</sub> -O]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[(CH <sub>3</sub> ) <sub>2</sub> CH] <sub>2</sub> NC(O)CH <sub>2</sub> -O]-benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[(CH <sub>3</sub> ) <sub>2</sub> CH] <sub>2</sub> NC(O)CH <sub>2</sub> -O]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	H	<i>p</i> -[-C(=NH)NH <sub>2</sub> ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	H	<i>p</i> -[-C(O)NH <sub>2</sub> ]-benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -( <i>N</i> -methylacclamido)benzyl-	-OCH <sub>2</sub> (CH <sub>3</sub> ) <sub>2</sub>

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -( <i>N</i> -methylacetamido)benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -( <i>N</i> -methyltrifluoroacetamido)benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		(1-toluenesulfonylimidazol-4-yl)methyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		1-(( <i>N</i> , <i>N</i> -dimethylaminosulfonyl)imidazol-4-yl)methyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -( <i>N</i> -toluenesulfonylamino)benzyl-	-OCH <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -( <i>N</i> -toluenesulfonylamino)benzyl-	-OH
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5-dimethylthiazolidin-4-yl)		<i>p</i> -(3- <i>N</i> , <i>N</i> -dimethylpropoxy)benzyl-	-OC(CH <sub>3</sub> ) <sub>3</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -( <i>N</i> -methylpiperidinoxy)benzyl-	-OC(CH <sub>3</sub> ) <sub>3</sub>

-- 55 --

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
<i>p</i> -CH <sub>3</sub> , $\phi$	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (1,5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -( <i>N</i> -methylpiperidin-4-yl)-benzyl-	-OC(CH <sub>3</sub> ) <sub>2</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)		<i>p</i> -( $\phi$ )-benzyl-	-OC(CH <sub>3</sub> ) <sub>2</sub>
<i>p</i> -CH <sub>3</sub> , $\phi$	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)		<i>p</i> -( $\phi$ )-benzyl-	-OH



### DETAILED DESCRIPTION OF THE INVENTION

As above, this invention relates to compounds which inhibit leukocyte adhesion and, in particular, leukocyte adhesion mediated by VLA-4.

However, prior to describing this invention in further detail, the following terms will first be defined.

#### Definitions

As used herein, "alkyl" refers to alkyl groups preferably having from 1 to 10 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, t-butyl, n-heptyl, octyl and the like.

"Substituted alkyl" refers to an alkyl group, preferably of from 1 to 10 carbon atoms, having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, alkylthioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxylaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted aryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl,

-NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

"Alkoxy" refers to the group "alkyl-O-" which includes, by way of example, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

"Substituted alkoxy" refers to the group "substituted alkyl-O-".

"Acyl" refers to the groups H-C(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, alkenyl-C(O)-, substituted alkenyl-C(O)-, alkynyl-C(O)-, substituted alkynyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, aryl-C(O)-, substituted aryl-C(O)-, heteroaryl-C(O)-, substituted heteroaryl-C(O)-, heterocyclic-C(O)-, and substituted heterocyclic-C(O)- wherein alkyl,

substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

5 "Acylamino" refers to the group -C(O)NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and where each R is joined to  
10 form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

15 "Thiocarbonylamino" refers to the group -C(S)NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and where each R is joined to  
20 form, together with the nitrogen atom a heterocyclic or substituted heterocyclic ring wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Acyloxy" refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-, alkenyl-C(O)O-, substituted alkenyl-C(O)O-, alkynyl-C(O)O-, substituted alkynyl-C(O)O-, aryl-C(O)O-, substituted aryl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, heteroaryl-C(O)O-, substituted heteroaryl-C(O)O-, heterocyclic-C(O)O-, and substituted heterocyclic-C(O)O- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl,  
30

-- 59 --

substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

5 "Alkenyl" refers to alkenyl group preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkenyl unsaturation.

"Substituted alkenyl" refers to alkenyl groups having from 1 to 5  
 10 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl,  
 15 carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
 25 oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2$ -alkyl,  $-\text{OS}(\text{O})_2$ -substituted alkyl,  $-\text{OS}(\text{O})_2$ -aryl,  $-\text{OS}(\text{O})_2$ -substituted aryl,  $-\text{OS}(\text{O})_2$ -heteroaryl,  $-\text{OS}(\text{O})_2$ -substituted heteroaryl,  $-\text{OS}(\text{O})_2$ -heterocyclic,  $-\text{OS}(\text{O})_2$ -substituted heterocyclic,  $-\text{OSO}_2$ -NRR where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2$ -alkyl,  $-\text{NRS}(\text{O})_2$ -substituted alkyl,  $-\text{NRS}(\text{O})_2$ -aryl,  $-\text{NRS}(\text{O})_2$ -substituted aryl,  
 30  $-\text{NRS}(\text{O})_2$ -heteroaryl,  $-\text{NRS}(\text{O})_2$ -substituted heteroaryl,  $-\text{NRS}(\text{O})_2$ -heterocyclic,  $-\text{NRS}(\text{O})_2$ -substituted heterocyclic,  $-\text{NRS}(\text{O})_2$ -NR-alkyl,

-- 60 --

-NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkenyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkenyl/substituted alkenyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

"Alkynyl" refers to alkynyl group preferably having from 2 to 10 carbon atoms and more preferably 3 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkynyl unsaturation.

"Substituted alkynyl" refers to alkynyl groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl,

guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl,  
 substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl,  
 substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic,  
 heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic,  
 5 cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted  
 heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
 oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2$ -alkyl,  $-\text{OS}(\text{O})_2$ -  
 substituted alkyl,  $-\text{OS}(\text{O})_2$ -aryl,  $-\text{OS}(\text{O})_2$ -substituted aryl,  $-\text{OS}(\text{O})_2$ -heteroaryl,  
 $-\text{OS}(\text{O})_2$ -substituted heteroaryl,  $-\text{OS}(\text{O})_2$ -heterocyclic,  $-\text{OS}(\text{O})_2$ -substituted  
 10 heterocyclic,  $-\text{OSO}_2$ -NRR where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2$ -alkyl,  
 $-\text{NRS}(\text{O})_2$ -substituted alkyl,  $-\text{NRS}(\text{O})_2$ -aryl,  $-\text{NRS}(\text{O})_2$ -substituted aryl,  
 $-\text{NRS}(\text{O})_2$ -heteroaryl,  $-\text{NRS}(\text{O})_2$ -substituted heteroaryl,  $-\text{NRS}(\text{O})_2$ -  
 heterocyclic,  $-\text{NRS}(\text{O})_2$ -substituted heterocyclic,  $-\text{NRS}(\text{O})_2$ -NR-alkyl,  
 $-\text{NRS}(\text{O})_2$ -NR-substituted alkyl,  $-\text{NRS}(\text{O})_2$ -NR-aryl,  $-\text{NRS}(\text{O})_2$ -NR-substituted  
 15 aryl,  $-\text{NRS}(\text{O})_2$ -NR-heteroaryl,  $-\text{NRS}(\text{O})_2$ -NR-substituted heteroaryl,  
 $-\text{NRS}(\text{O})_2$ -NR-heterocyclic,  $-\text{NRS}(\text{O})_2$ -NR-substituted heterocyclic where R is  
 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted  
 alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino,  
 mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino,  
 20 mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic  
 amino, unsymmetric di-substituted amines having different substituents  
 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,  
 substituted heteroaryl, heterocyclic and substituted heterocyclic and  
 substituted alkynyl groups having amino groups blocked by conventional  
 25 blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted  
 alkynyl groups substituted with  $-\text{SO}_2$ -alkyl,  $-\text{SO}_2$ -substituted alkyl,  $-\text{SO}_2$ -  
 alkenyl,  $-\text{SO}_2$ -substituted alkenyl,  $-\text{SO}_2$ -cycloalkyl,  $-\text{SO}_2$ -substituted  
 cycloalkyl,  $-\text{SO}_2$ -aryl,  $-\text{SO}_2$ -substituted aryl,  $-\text{SO}_2$ -heteroaryl,  $-\text{SO}_2$ -substituted  
 heteroaryl,  $-\text{SO}_2$ -heterocyclic,  $-\text{SO}_2$ -substituted heterocyclic and  $-\text{SO}_2$ -NRR  
 30 where R is hydrogen or alkyl.

-- 62 --

"Amidino" refers to the group  $\text{H}_2\text{NC}-$  and the term "alkylamidino"



5 refers to compounds having 1 to 3 alkyl groups (e.g., alkylHNC-).



10

"Thioamidino" refers to the group  $\text{RSC}-$  where R is hydrogen or



alkyl.

15

"Aminoacyl" refers to the groups -NRC(O)alkyl,

-NRC(O)substituted alkyl, -NRC(O)cycloalkyl, -NRC(O)substituted cycloalkyl, -NRC(O)alkenyl, -NRC(O)substituted alkenyl, -NRC(O)alkynyl,

-NRC(O)substituted alkynyl, -NRC(O)aryl, -NRC(O)substituted aryl,

20 -NRC(O)heteroaryl, -NRC(O)substituted heteroaryl, -NRC(O)heterocyclic,

and -NRC(O)substituted heterocyclic where R is hydrogen or alkyl and

wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl,

substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl,

heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are

25 as defined herein.

"Aminocarbonyloxy" refers to the groups -NRC(O)O-alkyl,

-NRC(O)O-substituted alkyl, -NRC(O)O-alkenyl, -NRC(O)O-substituted

alkenyl, -NRC(O)O-alkynyl, -NRC(O)O-substituted alkynyl, -NRC(O)O-

30 cycloalkyl, -NRC(O)O-substituted cycloalkyl, -NRC(O)O-aryl, -NRC(O)O-

substituted aryl, -NRC(O)O-heteroaryl, -NRC(O)O-substituted heteroaryl,

-NRC(O)O-heterocyclic, and -NRC(O)O-substituted heterocyclic where R is

hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted

alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl,

35 substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted

heterocyclic are as defined herein.

"Oxycarbonylamino" refers to the groups -OC(O)NH<sub>2</sub>, -OC(O)NRR, -OC(O)NR-alkyl, -OC(O)NR-substituted alkyl, -OC(O)NR-alkenyl, -OC(O)NR-substituted alkenyl, -OC(O)NR-alkynyl, -OC(O)NR-substituted alkynyl, -OC(O)NR-cycloalkyl, -OC(O)NR-substituted cycloalkyl, -OC(O)NR-aryl, -OC(O)NR-substituted aryl, -OC(O)NR-heteroaryl, -OC(O)NR-substituted heteroaryl, -OC(O)NR-heterocyclic, and -OC(O)NR-substituted heterocyclic where R is hydrogen, alkyl or where each R is joined to form, together with the nitrogen atom a heterocyclic or substituted heterocyclic ring and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Oxythiocarbonylamino" refers to the groups -OC(S)NH<sub>2</sub>, -OC(S)NRR, -OC(S)NR-alkyl, -OC(S)NR-substituted alkyl, -OC(S)NR-alkenyl, -OC(S)NR-substituted alkenyl, -OC(S)NR-alkynyl, -OC(S)NR-substituted alkynyl, -OC(S)NR-cycloalkyl, -OC(S)NR-substituted cycloalkyl, -OC(S)NR-aryl, -OC(S)NR-substituted aryl, -OC(S)NR-heteroaryl, -OC(S)NR-substituted heteroaryl, -OC(S)NR-heterocyclic, and -OC(S)NR-substituted heterocyclic where R is hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminocarbonylamino" refers to the groups -NRC(O)NRR, -NRC(O)NR-alkyl, -NRC(O)NR-substituted alkyl, -NRC(O)NR-alkenyl, -NRC(O)NR-substituted alkenyl, -NRC(O)NR-alkynyl, -NRC(O)NR-substituted alkynyl, -NRC(O)NR-aryl, -NRC(O)NR-substituted aryl, -NRC(O)NR-cycloalkyl, -NRC(O)NR-substituted cycloalkyl, -NRC(O)NR-heteroaryl, and -NRC(O)NR-substituted heteroaryl, -NRC(O)NR-heterocyclic,



-- 64 --

and -NRC(O)NR-substituted heterocyclic where each R is independently hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminothiocabonylamino" refers to the groups -NRC(S)NRR, -NRC(S)NR-alkyl, -NRC(S)NR-substituted alkyl, -NRC(S)NR-alkenyl, -NRC(S)NR-substituted alkenyl, -NRC(S)NR-alkynyl, -NRC(S)NR-substituted alkynyl, -NRC(S)NR-aryl, -NRC(S)NR-substituted aryl, -NRC(S)NR-cycloalkyl, -NRC(S)NR-substituted cycloalkyl, -NRC(S)NR-heteroaryl, and -NRC(S)NR-substituted heteroaryl, -NRC(S)NR-heterocyclic, and -NRC(S)NR-substituted heterocyclic where each R is independently hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aryl" or "Ar" refers to an unsaturated aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl or anthryl) which condensed rings may or may not be aromatic (e.g., 2-benzoxazolinone, 2H-1,4-benzoxazin-3(4H)-one-7-yl, and the like). Preferred aryls include phenyl and naphthyl.

Substituted aryl refers to aryl groups which are substituted with from 1 to 3 substituents selected from the group consisting of hydroxy, acyl,

acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy,  
 substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl,  
 amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy,  
 aminocarbonylamino, aminothiocarbonylamino, aryl, substituted aryl, aryloxy,  
 5 substituted aryloxy, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy,  
 substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
 carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl,  
 carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,  
 carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,  
 10 carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl,  
 substituted thioalkyl, thioaryl, substituted thioaryl, thioheteroaryl, substituted  
 thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic,  
 substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino,  
 guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic,  
 15 substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy,  
 substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
 oxycarbonylamino, oxythiocarbonylamino, -S(O)<sub>2</sub>-alkyl, -S(O)<sub>2</sub>-substituted  
 alkyl, -S(O)<sub>2</sub>-cycloalkyl, -S(O)<sub>2</sub>-substituted cycloalkyl, -S(O)<sub>2</sub>-alkenyl, -S(O)<sub>2</sub>-  
 substituted alkenyl, -S(O)<sub>2</sub>-aryl, -S(O)<sub>2</sub>-substituted aryl, -S(O)<sub>2</sub>-heteroaryl,  
 20 -S(O)<sub>2</sub>-substituted heteroaryl, -S(O)<sub>2</sub>-heterocyclic, -S(O)<sub>2</sub>-substituted  
 heterocyclic, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-  
 substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-  
 heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is  
 hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-  
 25 aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted  
 heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic,  
 -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl,  
 -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-  
 substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted  
 30 heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono-  
 and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-  
 substituted arylamino, mono- and di-heteroarylamino, mono- and di-

-- 66 --

substituted heteroaryl-amino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and amino groups on the substituted aryl blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or substituted with -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

"Aryloxy" refers to the group aryl-O- which includes, by way of example, phenoxy, naphthoxy, and the like.

"Substituted aryloxy" refers to substituted aryl-O- groups.

"Aryloxyaryl" refers to the group -aryl-O-aryl.

"Substituted aryloxyaryl" refers to aryloxyaryl groups substituted with from 1 to 3 substituents on either or both aryl rings selected from the group consisting of hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy, aminocarbonylamino, aminothiocabonylamino, aryl, substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic,

substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -S(O)<sub>2</sub>-alkyl, -S(O)<sub>2</sub>-substituted alkyl, -S(O)<sub>2</sub>-cycloalkyl, -S(O)<sub>2</sub>-substituted cycloalkyl, -S(O)<sub>2</sub>-alkenyl, -S(O)<sub>2</sub>-substituted alkenyl, -S(O)<sub>2</sub>-aryl, -S(O)<sub>2</sub>-substituted aryl, -S(O)<sub>2</sub>-heteroaryl, -S(O)<sub>2</sub>-substituted heteroaryl, -S(O)<sub>2</sub>-heterocyclic, -S(O)<sub>2</sub>-substituted heterocyclic, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and amino groups on the substituted aryl blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or substituted with -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

"Cycloalkyl" refers to cyclic alkyl groups of from 3 to 8 carbon atoms having a single cyclic ring including, by way of example, cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl and the like. Excluded from this definition are multi-ring alkyl groups such as adamantanyl, etc.

"Cycloalkenyl" refers to cyclic alkenyl groups of from 3 to 8 carbon atoms having single or multiple unsaturation but which are not aromatic.

"Substituted cycloalkyl" and "substituted cycloalkenyl" refer to a  
 5 cycloalkyl and cycloalkenyl groups, preferably of from 3 to 8 carbon atoms, having from 1 to 5 substituents selected from the group consisting of oxo (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino,  
 10 aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,  
 15 carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic,  
 20 cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted

alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

"Cycloalkoxy" refers to -O-cycloalkyl groups.

"Substituted cycloalkoxy" refers to -O-substituted cycloalkyl groups.

"Guanidino" refers to the groups -NRC(=NR)NRR, -NRC(=NR)NR-alkyl, -NRC(=NR)NR-substituted alkyl, -NRC(=NR)NR-alkenyl, -NRC(=NR)NR-substituted alkenyl, -NRC(=NR)NR-alkynyl, -NRC(=NR)NR-substituted alkynyl, -NRC(=NR)NR-aryl, -NRC(=NR)NR-substituted aryl, -NRC(=NR)NR-cycloalkyl, -NRC(=NR)NR-heteroaryl, -NRC(=NR)NR-substituted heteroaryl, -NRC(=NR)NR-heterocyclic, and -NRC(=NR)NR-substituted heterocyclic where each R is independently hydrogen and alkyl as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Guanidinosulfone" refers to the groups -NRC(=NR)NRSO<sub>2</sub>-alkyl,

-- 70 --

-NRC(=NR)NRSO<sub>2</sub>-substituted alkyl, -NRC(=NR)NRSO<sub>2</sub>-alkenyl,  
-NRC(=NR)NRSO<sub>2</sub>-substituted alkenyl, -NRC(=NR)NRSO<sub>2</sub>-alkynyl,  
-NRC(=NR)NRSO<sub>2</sub>-substituted alkynyl, -NRC(=NR)NRSO<sub>2</sub>-aryl,  
-NRC(=NR)NRSO<sub>2</sub>-substituted aryl, -NRC(=NR)NRSO<sub>2</sub>-cycloalkyl,  
5 -NRC(=NR)NRSO<sub>2</sub>-substituted cycloalkyl, -NRC(=NR)NRSO<sub>2</sub>-heteroaryl,  
and -NRC(=NR)NRSO<sub>2</sub>-substituted heteroaryl, -NRC(=NR)NRSO<sub>2</sub>-  
heterocyclic, and -NRC(=NR)NRSO<sub>2</sub>-substituted heterocyclic where each R is  
independently hydrogen and alkyl and wherein alkyl, substituted alkyl,  
alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl,  
10 substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
heterocyclic and substituted heterocyclic are as defined herein.

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo and  
preferably is either chloro or bromo.

15

"Heteroaryl" refers to an aromatic carbocyclic group of from 2 to 10  
carbon atoms and 1 to 4 heteroatoms selected from oxygen, nitrogen and  
sulfur within the ring. Such heteroaryl groups can have a single ring (e.g.,  
pyridyl or furyl) or multiple condensed rings (e.g., indoliziny or  
20 benzothieryl). Preferred heteroaryls include pyridyl, pyrrolyl, indolyl and  
furyl.

"Substituted heteroaryl" refers to heteroaryl groups which are  
substituted with from 1 to 3 substituents selected from the group consisting of  
25 hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl, substituted  
alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl, alkynyl,  
substituted alkynyl, amidino, alkylamidino, thioamidino, amino, aminoacyl,  
aminocarbonyloxy, aminocarbonylamino, aminothiobonylamino, aryl,  
substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted  
30 cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocycloxy,  
substituted heterocycloxy, carboxyl, carboxylalkyl, carboxyl-substituted  
alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl,

carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl,  
 carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido,  
 cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,  
 thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted  
 5 thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl,  
 substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl,  
 substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy,  
 substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,  
 heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino,  
 10 oxythiocarbonylamino, -S(O)<sub>2</sub>-alkyl, -S(O)<sub>2</sub>-substituted alkyl, -S(O)<sub>2</sub>-  
 cycloalkyl, -S(O)<sub>2</sub>-substituted cycloalkyl, -S(O)<sub>2</sub>-alkenyl, -S(O)<sub>2</sub>-substituted  
 alkenyl, -S(O)<sub>2</sub>-aryl, -S(O)<sub>2</sub>-substituted aryl, -S(O)<sub>2</sub>-heteroaryl, -S(O)<sub>2</sub>-  
 substituted heteroaryl, -S(O)<sub>2</sub>-heterocyclic, -S(O)<sub>2</sub>-substituted heterocyclic,  
 -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted  
 15 aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-  
 heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is  
 hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-  
 aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted  
 heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic,  
 20 -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl,  
 -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-  
 substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted  
 heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono-  
 and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-  
 25 substituted arylamino, mono- and di-heteroarylamino, mono- and di-  
 substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-  
 substituted heterocyclic amino, unsymmetric di-substituted amines having  
 different substituents selected from alkyl, substituted alkyl, aryl, substituted  
 aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted  
 30 heterocyclic and amino groups on the substituted aryl blocked by conventional  
 blocking groups such as Boc, Cbz, formyl, and the like or substituted with  
 -SO<sub>2</sub>NRR where R is hydrogen or alkyl.



"Heteroaryloxy" refers to the group -O-heteroaryl and "substituted heteroaryloxy" refers to the group -O-substituted heteroaryl.

5 "Heterocycle" or "heterocyclic" refers to a saturated or unsaturated group having a single ring or multiple condensed rings, from 1 to 10 carbon atoms and from 1 to 4 hetero atoms selected from nitrogen, sulfur or oxygen within the ring wherein, in fused ring systems, one or more the rings can be aryl or heteroaryl.

10 "Saturated heterocyclic" refers to heterocycles of single or multiple condensed rings lacking unsaturation in any ring (e.g., carbon to carbon unsaturation, carbon to nitrogen unsaturation, nitrogen to nitrogen unsaturation, and the like).

15 "Unsaturated heterocyclic" refers to non-aromatic heterocycles of single or multiple condensed rings having unsaturation in any ring (e.g., carbon to carbon unsaturation, carbon to nitrogen unsaturation, nitrogen to nitrogen unsaturation, and the like).

20 "Substituted heterocyclic" refers to heterocycle groups which are substituted with from 1 to 3 substituents selected from the group consisting of oxo (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino,  
25 aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,  
30 carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl,

- substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,
- 5 oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,
- 10  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is
- 15 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents
- 20 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with  $-\text{SO}_2\text{-alkyl}$ ,  $-\text{SO}_2\text{-substituted alkyl}$ ,  $-\text{SO}_2\text{-alkenyl}$ ,  $-\text{SO}_2\text{-substituted alkenyl}$ ,  $-\text{SO}_2\text{-cycloalkyl}$ ,  $-\text{SO}_2\text{-substituted cycloalkyl}$ ,  $-\text{SO}_2\text{-aryl}$ ,  $-\text{SO}_2\text{-substituted aryl}$ ,  $-\text{SO}_2\text{-heteroaryl}$ ,  $-\text{SO}_2\text{-substituted heteroaryl}$ ,  $-\text{SO}_2\text{-heterocyclic}$ ,  $-\text{SO}_2\text{-substituted heterocyclic}$  and  $-\text{SO}_2\text{NRR}$  where R is hydrogen or alkyl.
- 25
- 30 Examples of heterocycles and heteroaryls include, but are not limited to, azetidine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, dihydroindole, indazole, purine,

quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, 5 piperazine, indoline, phthalimide, 1,2,3,4-tetrahydroisoquinoline, 4,5,6,7-tetrahydrobenzo[b]thiophene, thiazole, thiazolidine, thiophene, benzo[b]thiophene, morpholino, thiomorpholino, piperidiny, pyrrolidine, tetrahydrofuranyl, and the like.

10 "Saturated substituted heterocyclic" refers to substituted heterocycles of single or multiple condensed rings lacking unsaturation in any ring (e.g., carbon to carbon unsaturation, carbon to nitrogen unsaturation, nitrogen to nitrogen unsaturation, and the like).

15 "Unsaturated substituted heterocyclic" refers to non-aromatic substituted heterocycles of single or multiple condensed rings having unsaturation in any ring (e.g., carbon to carbon unsaturation, carbon to nitrogen unsaturation, nitrogen to nitrogen unsaturation, and the like).

20 "Heterocyclyloxy" refers to the group -O-heterocyclic and "substituted heterocyclyloxy" refers to the group -O-substituted heterocyclic.

"Substituted alkylcarbonylamino" refers to the group -NHC(O)-substituted alkyl.

25

"Thiol" refers to the group -SH.

"Thioalkyl" refers to the group -S-alkyl.

30

"Substituted thioalkyl" refers to the group -S-substituted alkyl.

"Thiocycloalkyl" refers to the groups -S-cycloalkyl.

"Substituted thiocycloalkyl" refers to the group -S-substituted cycloalkyl.

5 "Thioaryl" refers to the group -S-aryl and "substituted thioaryl" refers to the group -S-substituted aryl.

"Thioheteroaryl" refers to the group -S-heteroaryl and "substituted thioheteroaryl" refers to the group -S-substituted heteroaryl.

10 "Thioheterocyclic" refers to the group -S-heterocyclic and "substituted thioheterocyclic" refers to the group -S-substituted heterocyclic.

"Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts of a compound of Formula I which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like.

15  
20

#### Compound Preparation

The compounds of this invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

25  
30

Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. Suitable protecting groups for various functional groups as well as suitable conditions for protecting and deprotecting particular functional groups are well known in the art. For example, numerous protecting groups are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Second Edition, Wiley, New York, 1991, and references cited therein.

Furthermore, the compounds of this invention will typically contain one or more chiral centers. Accordingly, if desired, such compounds can be prepared or isolated as pure stereoisomers, i.e., as individual enantiomers or diastereomers, or as stereoisomer-enriched mixtures. All such stereoisomers (and enriched mixtures) are included within the scope of this invention, unless otherwise indicated. Pure stereoisomers (or enriched mixtures) may be prepared using, for example, optically active starting materials or stereoselective reagents well-known in the art. Alternatively, racemic mixtures of such compounds can be separated using, for example, chiral column chromatography, chiral resolving agents and the like.

In a preferred method of synthesis, the compounds of formula I and IA wherein Q is  $-\text{C}(\text{O})\text{NR}^1-$  are prepared by first coupling an amino acid of formula II:



wherein  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  are as defined above, with a sulfonyl chloride of formula III:

-- 77 --



wherein  $\text{R}^1$  is as defined above, to provide an *N*-sulfonyl amino acid of formula IV:



wherein  $\text{R}^1\text{-R}^3$  are as defined above.

This reaction is typically conducted by contacting the amino acid of formula II with at least one equivalent, preferably about 1.1 to about 2 equivalents, of sulfonyl chloride III in an inert diluent such as dichloromethane and the like. Generally, the reaction is conducted at a temperature ranging from about  $-70^\circ\text{C}$  to about  $40^\circ\text{C}$  for about 1 to about 24 hours. Preferably, this reaction is conducted in the presence of a suitable base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, tertiary amines, such as triethylamine, diisopropylethylamine, *N*-methylmorpholine and the like. Alternatively, the reaction can be conducted under Schotten-Baumann-type conditions using aqueous alkali, such as sodium hydroxide and the like, as the base. Upon completion of the reaction, the resulting *N*-sulfonyl amino acid IV is recovered by conventional methods including neutralization, extraction, precipitation, chromatography, filtration, and the like.


The amino acids of formula II employed in the above reaction are either known compounds or compounds that can be prepared from known compounds by conventional synthetic procedures. Examples of suitable amino acids for use in this reaction include, but are not limited to, L-proline, *trans*-4-hydroxyl-L-proline, *cis*-4-hydroxyl-L-proline, *trans*-3-phenyl-L-proline, *cis*-3-phenyl-L-proline, L-(2-methyl)proline, L-pipecolinic acid, L-

azetidine-2-carboxylic acid, L-indoline-2-carboxylic acid, L-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, L-thiazolidine-4-carboxylic acid, L-(5,5-dimethyl)thiazolidine-4-carboxylic acid, L-thiamorpholine-3-carboxylic acid, glycine, 2-*tert*-butylglycine, D,L-phenylglycine, L-alanine,  $\alpha$ -methylalanine, *N*-methyl-L-phenylalanine, L-diphenylalanine, sarcosine, D,L-phenylsarcosine, L-aspartic acid  $\beta$ -*tert*-butyl ester, L-glutamic acid  $\gamma$ -*tert*-butyl ester, L-(*O*-benzyl)serine, 1-aminocyclopropanecarboxylic acid, 1-aminocyclobutanecarboxylic acid, 1-aminocyclopentanecarboxylic acid (cycloleucine) 1-aminocyclohexanecarboxylic acid, L-serine and the like. If  
5 desired, the corresponding carboxylic acid esters of the amino acids of formula II, such as the methyl esters, ethyl esters and the like, can be employed in the above reaction with the sulfonyl chloride III. Subsequent hydrolysis of the ester group to the carboxylic acid using conventional reagents and conditions, i.e., treatment with an alkali metal hydroxide in an inert diluent such as  
10 methanol/water, then provides the *N*-sulfonyl amino acid IV.

Similarly, the sulfonyl chlorides of formula III employed in the above reaction are either known compounds or compounds that can be prepared from known compounds by conventional synthetic procedures. Such compounds  
20 are typically prepared from the corresponding sulfonic acid, i.e., from compounds of the formula  $R^1-SO_3H$  where  $R^1$  is as defined above, using phosphorous trichloride and phosphorous pentachloride. This reaction is generally conducted by contacting the sulfonic acid with about 2 to 5 molar equivalents of phosphorous trichloride and phosphorous pentachloride, either  
25 neat or in an inert solvent, such as dichloromethane, at temperature in the range of about 0°C to about 80°C for about 1 to about 48 hours to afford the sulfonyl chloride. Alternatively, the sulfonyl chlorides of formula III can be prepared from the corresponding thiol compound, i.e., from compounds of the formula  $R^1-SH$  where  $R^1$  is as defined above, by treating the thiol with  
30 chlorine ( $Cl_2$ ) and water under conventional reaction conditions.

Examples of sulfonyl chlorides suitable for use in this invention include, but are not limited to, methanesulfonyl chloride, 2-propanesulfonyl chloride, 1-butanesulfonyl chloride, benzenesulfonyl chloride, 1-naphthalenesulfonyl chloride, 2-naphthalenesulfonyl chloride, *p*-toluenesulfonyl chloride,  $\alpha$ -toluenesulfonyl chloride, 4-acetamidobenzenesulfonyl chloride, 4-amidinobenzenesulfonyl chloride, 4-*tert*-butylbenzenesulfonyl chloride, 4-bromobenzenesulfonyl chloride, 2-carboxybenzenesulfonyl chloride, 4-cyanobenzenesulfonyl chloride, 3,4-dichlorobenzenesulfonyl chloride, 3,5-dichlorobenzenesulfonyl chloride, 3,4-dimethoxybenzenesulfonyl chloride, 3,5-ditrifluoromethylbenzenesulfonyl chloride, 4-fluorobenzenesulfonyl chloride, 4-methoxybenzenesulfonyl chloride, 2-methoxycarbonylbenzenesulfonyl chloride, 4-methylamidobenzenesulfonyl chloride, 4-nitrobenzenesulfonyl chloride, 4-thioamidobenzenesulfonyl chloride, 4-trifluoromethylbenzenesulfonyl chloride, 4-trifluoromethoxybenzenesulfonyl chloride, 2,4,6-trimethylbenzenesulfonyl chloride, 2-phenylethanesulfonyl chloride, 2-thiophenesulfonyl chloride, 5-chloro-2-thiophenesulfonyl chloride, 2,5-dichloro-4-thiophenesulfonyl chloride, 2-thiazolesulfonyl chloride, 2-methyl-4-thiazolesulfonyl chloride, 1-methyl-4-imidazolesulfonyl chloride, 1-methyl-4-pyrazolesulfonyl chloride, 5-chloro-1,3-dimethyl-4-pyrazolesulfonyl chloride, 3-pyridinesulfonyl chloride, 2-pyrimidinesulfonyl chloride, and the like. If desired, a sulfonyl fluoride, sulfonyl bromide or sulfonic acid anhydride may be used in place of the sulfonyl chloride in the above reaction to form the *N*-sulfonyl amino acids of formula IV.

The intermediate *N*-sulfonyl amino acids of formula IV can also be prepared by reacting a sulfonamide of formula V:




V



wherein R<sup>1</sup> and R<sup>2</sup> are as defined above, with a carboxylic acid derivative of the formula L(R<sup>3</sup>)CHCOOR where L is a leaving group, such as chloro, bromo, iodo, mesylate, tosylate and the like, R<sup>3</sup> is as defined above and R is hydrogen or an alkyl group. This reaction is typically conducted by contacting the sulfonamide V with at least one equivalent, preferably 1.1 to 2 equivalents, of the carboxylic acid derivative in the presence of a suitable base, such as triethylamine, in an inert diluent, such as DMF, at a temperature ranging from about 24°C to about 37°C for about 0.5 to about 4 hours. This reaction is further described in Zuckermann et al., *J. Am. Chem. Soc.*, 1992, 114, 10646-10647. Preferred carboxylic acid derivatives for use in this reaction are α-chloro and α-bromocarboxylic acid esters such as *tert*-butyl bromoacetate, and the like. When an carboxylic acid ester is employed in this reaction, the ester group is subsequently hydrolyzed using conventional procedures to afford an *N*-sulfonyl amino acid of formula IV.

The compounds of formula I are then prepared by coupling the intermediate *N*-sulfonyl amino acid of formula IV with an amino acid derivative of formula VI:



wherein R<sup>5</sup>-R<sup>7</sup> are as defined above. This coupling reaction is typically conducted using well-known coupling reagents such as carbodiimides, BOP reagent (benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphonate) and the like. Suitable carbodiimides include, by way of example, dicyclohexylcarbodiimide (DCC), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) and the like. If desired, polymer supported forms of carbodiimide coupling reagents may also be used including, for example, those described in *Tetrahedron Letters*, 34(48), 7685 (1993). Additionally,

well-known coupling promoters, such as N-hydroxysuccinimide, 1-hydroxybenzotriazole and the like, may be used to facilitate the coupling reaction.

5           This coupling reaction is typically conducted by contacting the *N*-sulfonylamino acid IV with about 1 to about 2 equivalents of the coupling reagent and at least one equivalent, preferably about 1 to about 1.2 equivalents, of amino acid derivative VI in an inert diluent, such as dichloromethane, chloroform, acetonitrile, tetrahydrofuran, *N,N*-  
10 dimethylformamide and the like. Generally, this reaction is conducted at a temperature ranging from about 0°C to about 37°C for about 12 to about 24 hours. Upon completion of the reaction, the compound of formula I is recovered by conventional methods including neutralization, extraction, precipitation, chromatography, filtration, and the like.


15           Alternatively, the *N*-sulfonyl amino acid IV can be converted into an acid halide and the acid halide coupled with amino acid derivative VI to provide compounds of formula I. The acid halide of VI can be prepared by contacting VI with an inorganic acid halide, such as thionyl chloride,  
20 phosphorous trichloride, phosphorous tribromide or phosphorous pentachloride, or preferably, with oxalyl chloride under conventional conditions. Generally, this reaction is conducted using about 1 to 5 molar equivalents of the inorganic acid halide or oxalyl chloride, either neat or in an inert solvent, such as dichloromethane or carbon tetrachloride, at temperature  
25 in the range of about 0°C to about 80°C for about 1 to about 48 hours. A catalyst, such as *N,N*-dimethylformamide, may also be used in this reaction.

          The acid halide of *N*-sulfonyl amino acid IV is then contacted with at least one equivalent, preferably about 1.1 to about 1.5 equivalents, of amino  
30 acid derivative VI in an inert diluent, such as dichloromethane, at a temperature ranging from about -70°C to about 40°C for about 1 to about 24 hours. Preferably, this reaction is conducted in the presence of a suitable base

-- 82 --

to scavenge the acid generated during the reaction. Suitable bases include, by way of example, tertiary amines, such as triethylamine, diisopropylethylamine, *N*-methyldmorpholine and the like. Alternatively, the reaction can be conducted under Schotten-Baumann-type conditions using aqueous alkali, such as sodium hydroxide and the like. Upon completion of the reaction, the compound of formula I is recovered by conventional methods including neutralization, extraction, precipitation, chromatography, filtration, and the like.

Alternatively, the compounds of formula I can be prepared by first forming a diamino acid derivative of formula VII:



wherein  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^5$  -  $\text{R}^7$  are as defined above. The diamino acid derivatives of formula VII can be readily prepared by coupling an amino acid of formula II with an amino acid derivative of formula VI using conventional amino acid coupling techniques and reagents, such carbodiimides, BOP reagent and the like, as described above. Diamino acid VII can then be sulfonated using a sulfonyl chloride of formula III and using the synthetic procedures described above to provide a compound of formula I.

The amino acid derivatives of formula VI employed in the above reactions are either known compounds or compounds that can be prepared from known compounds by conventional synthetic procedures. For example, amino acid derivatives of formula VI can be prepared by *C*-alkylating commercially available diethyl 2-acetamidomalonate (Aldrich, Milwaukee; Wisconsin, USA) with an alkyl or substituted alkyl halide. This reaction is typically conducted by treating the diethyl 2-acetamidomalonate with at least

one equivalent of sodium ethoxide and at least one equivalent of an alkyl or substituted alkyl halide in refluxing ethanol for about 6 to about 12 hours. The resulting C-alkylated malonate is then deacetylated, hydrolyzed and decarboxylated by heating in aqueous hydrochloric acid at reflux for about 6  
5 to about 12 hours to provide the amino acid, typically as the hydrochloride salt.

Examples of amino acid derivatives of formula VI suitable for use in the above reactions include, but are not limited to, L-4-nitrophenylalanine  
10 methyl ester, L-tyrosine methyl ester, D,L-homo-4-nitrophenylalanine methyl ester, L-(*O*-benzyl)tyrosine methyl ester, L-3,5-diiodotyrosine methyl ester, L-3-iodotyrosine methyl ester and the like. If desired, of course, other esters or amides of the above-described compounds may also be employed.

15 For ease of synthesis, the compounds of formula I are typically prepared as an ester, i.e., where R<sup>6</sup> is an alkoxy or substituted alkoxy group and the like. If desired, the ester group can be hydrolysed using conventional conditions and reagents to provide the corresponding carboxylic acid. Typically, this reaction is conducted by treating the ester with at least one  
20 equivalent of an alkali metal hydroxide, such as lithium, sodium or potassium hydroxide, in an inert diluent, such as methanol or mixtures of methanol and water, at a temperature ranging about 0°C to about 24°C for about 1 to about 12 hours. Alternatively, benzyl esters may be removed by hydrogenolysis using a palladium catalyst, such as palladium on carbon. The resulting  
25 carboxylic acids may be coupled, if desired, to amines such as β-alanine ethyl ester, hydroxyamines such as hydroxylamine and *N*-hydroxysuccinimide, alkoxyamines and substituted alkoxyamines such as *O*-methylhydroxylamine and *O*-benzylhydroxylamine, and the like, using conventional coupling reagents and conditions as described above.

30

As will be apparent to those skilled in the art, other functional groups present on any of the substituents of the compounds of formula I can be

readily modified or derivatized either before or after the above-described coupling reactions using well-known synthetic procedures. For example, a nitro group present on a substituent of a compound of formula I or an intermediate thereof may be readily reduced by hydrogenation in the presence of a palladium catalyst, such as palladium on carbon, to provide the corresponding amino group. This reaction is typically conducted at a temperature of from about 20°C to about 50°C for about 6 to about 24 hours in an inert diluent, such as methanol. Compounds having a nitro group on the R<sup>5</sup> substituent can be prepared, for example, by using a 4-nitrophenylalanine derivative and the like in the above-described coupling reactions.

Similarly, a pyridyl group can be hydrogenated in the presence of a platinum catalyst, such as platinum oxide, in an acidic diluent to provide the corresponding piperidinyl analogue. Generally, this reaction is conducted by treating the pyridine compound with hydrogen at a pressure ranging from about 20 psi to about 60 psi, preferably about 40 psi, in the presence of the catalyst at a temperature of about 20°C to about 50°C for about 2 to about 24 hours in an acidic diluent, such as a mixture of methanol and aqueous hydrochloric acid. Compounds having a pyridyl group can be readily prepared by using, for example,  $\beta$ -(2-pyridyl)-,  $\beta$ -(3-pyridyl)- or  $\beta$ -(4-pyridyl)-L-alanine derivatives in the above-described coupling reactions.

Additionally, when the R<sup>5</sup> substituent of a compound of formula I or an intermediate thereof contains a primary or secondary amino group, such amino groups can be further derivatized either before or after the above coupling reactions to provide, by way of example, amides, sulfonamides, ureas, thioureas, carbamates, secondary or tertiary amines and the like. Compounds having a primary amino group on the R<sup>5</sup> substituent may be prepared, for example, by reduction of the corresponding nitro compound as described above. Alternatively, such compounds can be prepared by using an amino acid derivative of formula VI derived from lysine, 4-aminophenylalanine and the like in the above-described coupling reactions.

By way of illustration, a compound of formula I or an intermediate thereof having a substituent containing a primary or secondary amino group, such as where R<sup>3</sup> is a (4-aminophenyl)methyl group, can be readily *N*-acylated using conventional acylating reagents and conditions to provide the

5 corresponding amide. This acylation reaction is typically conducted by treating the amino compound with at least one equivalent, preferably about 1.1 to about 1.2 equivalents, of a carboxylic acid in the presence of a coupling reagent such as a carbodiimide, BOP reagent (benzotriazol-1-yloxy-

10 tris(dimethylamino)phosphonium hexafluorophosphonate) and the like, in an inert diluent, such as dichloromethane, chloroform, acetonitrile, tetrahydrofuran, *N,N*-dimethylformamide and the like, at a temperature ranging from about 0°C to about 37°C for about 4 to about 24 hours. Preferably, a promoter, such as *N*-hydroxysuccinimide, 1-hydroxy-

15 benzotriazole and the like, is used to facilitate the acylation reaction. Examples of carboxylic acids suitable for use in this reaction include, but are not limited to, *N*-*tert*-butyloxycarbonylglycine, *N*-*tert*-butyloxycarbonyl-L-

20 phenylalanine, *N*-*tert*-butyloxycarbonyl-L-aspartic acid benzyl ester, benzoic acid, *N*-*tert*-butyloxycarbonylisonipecotic acid, *N*-methylisonipecotic acid, *N*-*tert*-butyloxycarbonylnipecotic acid, *N*-*tert*-butyloxycarbonyl-L-

25 tetrahydroisoquinoline-3-carboxylic acid, *N*-(toluene-4-sulfonyl)-L-proline and the like.

Alternatively, a compound of formula I or an intermediate thereof containing a primary or secondary amino group can be *N*-acylated using an

25 acyl halide or a carboxylic acid anhydride to form the corresponding amide. This reaction is typically conducted by contacting the amino compound with at least one equivalent, preferably about 1.1 to about 1.2 equivalents, of the acyl halide or carboxylic acid anhydride in an inert diluent, such as

30 dichloromethane, at a temperature ranging from about -70°C to about 40°C for about 1 to about 24 hours. If desired, an acylation catalyst such as 4-(*N,N*-dimethylamino)pyridine may be used to promote the acylation reaction. The acylation reaction is preferably conducted in the presence of a suitable

base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, tertiary amines, such as triethylamine, diisopropylethylamine, *N*-methylmorpholine and the like. Alternatively, the reaction can be conducted under Schotten-Baumann-type conditions using aqueous alkali, such as sodium hydroxide and the like.

Examples of acyl halides and carboxylic acid anhydrides suitable for use in this reaction include, but are not limited to, 2-methylpropionyl chloride, trimethylacetyl chloride, phenylacetyl chloride, benzoyl chloride, 2-bromobenzoyl chloride, 2-methylbenzoyl chloride, 2-trifluoro-methylbenzoyl chloride, isonicotinoyl chloride, nicotinoyl chloride, picolinoyl chloride, acetic anhydride, succinic anhydride, and the like. Carbamyl chlorides, such as *N,N*-dimethylcarbamyl chloride, *N,N*-diethylcarbamyl chloride and the like, can also be used in this reaction to provide ureas. Similarly, dicarbonates, such as di-*tert*-butyl dicarbonate, may be employed to provide carbamates.

In a similar manner, a compound of formula I or an intermediate thereof containing a primary or secondary amino group may be *N*-sulfonated to form a sulfonamide using a sulfonyl halide or a sulfonic acid anhydride.

Sulfonyl halides and sulfonic acid anhydrides suitable for use in this reaction include, but are not limited to, methanesulfonyl chloride, chloromethanesulfonyl chloride, *p*-toluenesulfonyl chloride, trifluoromethanesulfonic anhydride, and the like. Similarly, sulfamoyl chlorides, such as dimethylsulfamoyl chloride, can be used to provide sulfamides (e.g., >N-SO<sub>2</sub>-N<).

Additionally, a primary and secondary amino group present on a substituent of a compound of formula I or an intermediate thereof can be reacted with an isocyanate or a thioisocyanate to give a urea or thiourea, respectively. This reaction is typically conducted by contacting the amino compound with at least one equivalent, preferably about 1.1 to about 1.2 equivalents, of the isocyanate or thioisocyanate in an inert diluent, such as

toluene and the like, at a temperature ranging from about 24°C to about 37°C for about 12 to about 24 hours. The isocyanates and thioisocyanates used in this reaction are commercially available or can be prepared from commercially available compounds using well-known synthetic procedures. For example, isocyanates and thioisocyanates are readily prepared by reacting the appropriate amine with phosgene or thiophosgene. Examples of isocyanates and thioisocyanates suitable for use in this reaction include, but are not limited to, ethyl isocyanate, *n*-propyl isocyanate, 4-cyanophenyl isocyanate, 3-methoxyphenyl isocyanate, 2-phenylethyl isocyanate, methyl thioisocyanate, ethyl thioisocyanate, 2-phenylethyl thioisocyanate, 3-phenylpropyl thioisocyanate, 3-(*N,N*-diethylamino)propyl thioisocyanate, phenyl thioisocyanate, benzyl thioisocyanate, 3-pyridyl thioisocyanate, fluorescein isothiocyanate (isomer I) and the like.

Furthermore, when a compound of formula I or an intermediate thereof contains a primary or secondary amino group, the amino group can be reductively alkylated using aldehydes or ketones to form a secondary or tertiary amino group. This reaction is typically conducted by contacting the amino compound with at least one equivalent, preferably about 1.1 to about 1.5 equivalents, of an aldehyde or ketone and at least one equivalent based on the amino compound of a metal hydride reducing agent, such as sodium cyanoborohydride, in an inert diluent, such as methanol, tetrahydrofuran, mixtures thereof and the like, at a temperature ranging from about 0°C to about 50°C for about 1 to about 72 hours. Aldehydes and ketones suitable for use in this reaction include, by way of example, benzaldehyde, 4-chlorobenzaldehyde, valeraldehyde and the like.

In a similar manner, when a compound of formula I or an intermediate thereof has a substituent containing a hydroxyl group, the hydroxyl group can be further modified or derivatized either before or after the above coupling reactions to provide, by way of example, ethers, carbamates and the like. Compounds having a hydroxyl group on the R<sup>5</sup> substituent, for example, can



be prepared using an amino acid derivative of formula VI derived from tyrosine and the like in the above-described reactions.

By way of example, a compound of formula I or an intermediate thereof having a substituent containing a hydroxyl group, such as where R<sup>5</sup> is a (4-hydroxyphenyl)methyl group, can be readily *O*-alkylated to form ethers. This *O*-alkylation reaction is typically conducted by contacting the hydroxy compound with a suitable alkali or alkaline earth metal base, such as potassium carbonate, in an inert diluent, such as acetone, 2-butanone and the like, to form the alkali or alkaline earth metal salt of the hydroxyl group. This salt is generally not isolated, but is reacted *in situ* with at least one equivalent of an alkyl or substituted alkyl halide or sulfonate, such as an alkyl chloride, bromide, iodide, mesylate or tosylate, to afford the ether. Generally, this reaction is conducted at a temperature ranging from about 60°C to about 150°C for about 24 to about 72 hours. Preferably, a catalytic amount of sodium or potassium iodide is added to the reaction mixture when an alkyl chloride or bromide is employed in the reaction.

Examples of alkyl or substituted alkyl halides and sulfonates suitable for use in this reaction include, but are not limited to, *tert*-butyl bromoacetate, *N*-*tert*-butyl chloroacetamide, 1-bromoethylbenzene, ethyl  $\alpha$ -bromophenylacetate, 2-(*N*-ethyl-*N*-phenylamino)ethyl chloride, 2-(*N,N*-ethylamino)ethyl chloride, 2-(*N,N*-diisopropylamino)ethyl chloride, 2-(*N,N*-dibenzylamino)ethyl chloride, 3-(*N,N*-ethylamino)propyl chloride, 3-(*N*-benzyl-*N*-methylamino)propyl chloride, *N*-(2-chloroethyl)morpholine, 2-(hexamethyleneimino)ethyl chloride, 3-(*N*-methylpiperazine)propyl chloride, 1-(3-chlorophenyl)-4-(3-chloropropyl)piperazine, 2-(4-hydroxy-4-phenylpiperidine)ethyl chloride, *N*-*tert*-butyloxycarbonyl-3-piperidinemethyl tosylate, and the like.

30

Alternatively, a hydroxyl group present on a substituent of a compound of formula I or an intermediate thereof can be *O*-alkylating using the

-- 89 --

Mitsunobu reaction. In this reaction, an alcohol, such as 3-(*N,N*-dimethylamino)-1-propanol and the like, is reacted with about 1.0 to about 1.3 equivalents of triphenylphosphine and about 1.0 to about 1.3 equivalents of diethyl azodicarboxylate in an inert diluent, such as tetrahydrofuran, at a temperature ranging from about -10°C to about 5°C for about 0.25 to about 1 hour. About 1.0 to about 1.3 equivalents of a hydroxy compound, such as *N*-*tert*-butyltyrosine methyl ester, is then added and the reaction mixture is stirred at a temperature of about 0°C to about 30°C for about 2 to about 48 hours to provide the *O*-alkylated product.

In a similar manner, a compound of formula I or an intermediate thereof containing a aryl hydroxy group can be reacted with an aryl iodide to provide a diaryl ether. Generally, this reaction is conducted by forming the alkali metal salt of the hydroxyl group using a suitable base, such as sodium hydride, in an inert diluent such as xylenes at a temperature of about -25°C to about 10°C. The salt is then treated with about 1.1 to about 1.5 equivalents of cuprous bromide dimethyl sulfide complex at a temperature ranging from about 10°C to about 30°C for about 0.5 to about 2.0 hours, followed by about 1.1 to about 1.5 equivalents of an aryl iodide, such as sodium 2-iodobenzoate and the like. The reaction is then heated to about 70°C to about 150°C for about 2 to about 24 hours to provide the diaryl ether.

Additionally, a hydroxy-containing compound can also be readily derivatized to form a carbamate. In one method for preparing such carbamates, a hydroxy compound of formula I or an intermediate thereof is contacted with about 1.0 to about 1.2 equivalents of 4-nitrophenyl chloroformate in an inert diluent, such as dichloromethane, at a temperature ranging from about -25°C to about 0°C for about 0.5 to about 2.0 hours. Treatment of the resulting carbonate with an excess, preferably about 2 to about 5 equivalents, of a trialkylamine, such as triethylamine, for about 0.5 to 2 hours, followed by about 1.0 to about 1.5 equivalents of a primary or secondary amine provides the carbamate. Examples of amines suitable for

-- 90 --

using in this reaction include, but are not limited to, piperazine, 1-methylpiperazine, 1-acetylpiperazine, morpholine, thiomorpholine, pyrrolidine, piperidine and the like.

5           Alternatively, in another method for preparing carbamates, a hydroxy-containing compound is contacted with about 1.0 to about 1.5 equivalents of a carbamyl chloride in an inert diluent, such as dichloromethane, at a temperature ranging from about 25°C to about 70°C for about 2 to about 72  
10           hours. Typically, this reaction is conducted in the presence of a suitable base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, tertiary amines, such as triethylamine, diisopropylethylamine, *N*-methyldmorpholine and the like. Additionally, at least one equivalent (based on the hydroxy compound) of 4-(*N,N*-dimethylamino)pyridine is preferably added to the reaction mixture to facilitate the reaction. Examples of carbamyl  
15           chlorides suitable for use in this reaction include, by way of example, dimethylcarbamyl chloride, diethylcarbamyl chloride and the like.

             Likewise, when a compound of formula I or an intermediate thereof contains a primary or secondary hydroxyl group, such hydroxyl groups can be  
20           readily converted into a leaving group and displaced to form, for example, amines, sulfides and fluorides. For example, derivatives of 4-hydroxy-L-proline can be converted into the corresponding 4-amino, 4-thio or 4-fluoro-L-proline derivatives via nucleophilic displacement of the derivatized hydroxyl group. Generally, when a chiral compound is employed in these reactions, the  
25           stereochemistry at the carbon atom attached to the derivatized hydroxyl group is typically inverted.

---

             These reactions are typically conducted by first converting the hydroxyl group into a leaving group, such as a tosylate, by treatment of the  
30           hydroxy compound with at least one equivalent of a sulfonyl halide, such as *p*-toluenesulfonyl chloride and the like, in pyridine. This reaction is generally conducted at a temperature of from about 0°C to about 70°C for about 1 to

-- 91 --

about 48 hours. The resulting tosylate can then be readily displaced with sodium azide, for example, by contacting the tosylate with at least one equivalent of sodium azide in an inert diluent, such as a mixture of *N,N*-dimethylformamide and water, at a temperature ranging from about 0°C to about 37°C for about 1 to about 12 hours to provide the corresponding azido compound. The azido group can then be reduced by, for example, hydrogenation using a palladium on carbon catalyst to provide the amino (-NH<sub>2</sub>) compound.

Similarly, a tosylate group can be readily displaced by a thiol to form a sulfide. This reaction is typically conducted by contacting the tosylate with at least one equivalent of a thiol, such as thiophenol, in the presence of a suitable base, such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), in an inert diluent, such as *N,N*-dimethylformamide, at a temperature of from about 0°C to about 37°C for about 1 to about 12 hours to provide the sulfide. Additionally, treatment of a tosylate with morpholinosulfur trifluoride in an inert diluent, such as dichloromethane, at a temperature ranging from about 0°C to about 37°C for about 12 to about 24 hours affords the corresponding fluoro compound.

Furthermore, a compound of formula I or an intermediate thereof having a substituent containing an iodoaryl group, for example, when R<sup>5</sup> is a (4-iodophenyl)methyl group, can be readily converted either before or after the above coupling reactions into a biaryl compound. Typically, this reaction is conducted by treating the iodoaryl compound with about 1.1 to about 2 equivalents of an arylzinc iodide, such as 2-(methoxycarbonyl)phenylzinc iodide, in the presence of a palladium catalyst, such as palladium tetra(triphenylphosphine), in an inert diluent, such as tetrahydrofuran, at a temperature ranging from about 24°C to about 30°C until reaction completion. This reaction is further described, for example, in Rieke, *J. Org. Chem.* 1991, 56, 1445.

-- 92 --

In some cases, the compounds of formula I or intermediates thereof may contain substituents having one or more sulfur atoms. Such sulfur atoms will be present, for example, when the amino acid of formula II employed in the above reactions is derived from L-thiazolidine-4-carboxylic acid, L-(5,5-dimethyl)thiazolidine-4-carboxylic acid, L-thiamorpholine-3-carboxylic acid and the like. When present, such sulfur atoms can be oxidized either before or after the above coupling reactions to provide a sulfoxide or sulfone compound using conventional reagents and reaction conditions. Suitable reagents for oxidizing a sulfide compound to a sulfoxide include, by way of example, hydrogen peroxide, 3-chloroperoxybenzoic acid (MCPBA), sodium periodate and the like. The oxidation reaction is typically conducted by contacting the sulfide compound with about 0.95 to about 1.1 equivalents of the oxidizing reagent in an inert diluent, such as dichloromethane, at a temperature ranging from about -50°C to about 75°C for about 1 to about 24 hours. The resulting sulfoxide can then be further oxidized to the corresponding sulfone by contacting the sulfoxide with at least one additional equivalent of an oxidizing reagent, such as hydrogen peroxide, MCPBA, potassium permanganate and the like. Alternatively, the sulfone can be prepared directly by contacting the sulfide with at least two equivalents, and preferably an excess, of the oxidizing reagent. Such reactions are described further in March, "Advanced Organic Chemistry", 4th Ed., pp. 1201-1202, Wiley Publisher, 1992.

As described above, the compounds of formula I having an R<sup>2</sup> substituent other than hydrogen can be prepared using an N-substituted amino acid of formula II, such as sarcosine, N-methyl-L-phenylalanine and the like, in the above-described coupling reactions. Alternatively, such compounds can be prepared by N-alkylation of a sulfonamide of formula I or IV (where R<sup>2</sup> is hydrogen) using conventional synthetic procedures. Typically, this N-alkylation reaction is conducted by contacting the sulfonamide with at least one equivalent, preferably 1.1 to 2 equivalents, of an alkyl or substituted alkyl halide in the presence of a suitable base, such as potassium carbonate, in an inert diluent, such as acetone, 2-butanone and the like, at a temperature

-- 93 --

ranging from about 25°C to about 70°C for about 2 to about 48 hours.

Examples of alkyl or substituted alkyl halides suitable for use in this reaction include, but are not limited to, methyl iodide, and the like.

5           Additionally, the sulfonamides of formula I or IV wherein R<sup>2</sup> is hydrogen and R<sup>1</sup> is a 2-alkoxycarbonylaryl group can be intramolecularly cyclized to form 1,2-benzisothiazol-3-one derivatives or analogues thereof. This reaction is typically conducted by treating a sulfonamide, such as *N*-(2-methoxycarbonylphenylsulfonyl)glycine-L-phenylalanine benzyl ester, with  
10           about 1.0 to 1.5 equivalents of a suitable base, such as an alkali metal hydride, in a inert diluent, such as tetrahydrofuran, at a temperature ranging from about 0°C to about 30°C for about 2 to about 48 hours to afford the cyclized 1,2-benzisothiazol-3-one derivative.

15           Lastly, the compounds of formula I where Q is -C(S)NR<sup>2</sup> are can prepared by using an amino thionoacid derivative in place of amino acid II in the above described synthetic procedures. Such amino thionoacid derivatives can be prepared by the procedures described in Shalaky, et al., *J. Org. Chem.*, 61:9045-9048 (1996) and Brain, et al., *J. Org. Chem.*, 62:3808-3809 (1997)  
20           and references cited therein.

#### Pharmaceutical Formulations

          When employed as pharmaceuticals, the compounds of formula I and IA are usually administered in the form of pharmaceutical compositions.  
25           These compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds are effective as both injectable and oral compositions. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.

30           This invention also includes pharmaceutical compositions which contain, as the active ingredient, one or more of the compounds of formula I

-- 94 --

and IA above associated with pharmaceutically acceptable carriers. In making the compositions of this invention, the active ingredient is usually mixed with an excipient, diluted by an excipient or enclosed within such a carrier which can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, it may be necessary to mill the active compound to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it ordinarily is milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size is normally adjusted by milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxybenzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

-- 95 --

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 to about 100 mg, more usually about 10 to about 30 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

The active compound is effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It, will be understood, however, that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, 0.1 to about 500 mg of the active ingredient of the present invention.

The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to



-- 96 --

resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable solvents may be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device may be attached to a face masks tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

The following formulation examples illustrate the pharmaceutical compositions of the present invention.

#### Formulation Example 1

Hard gelatin capsules containing the following ingredients are prepared:

-- 97 --

	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
5	Active Ingredient	30.0
	Starch	305.0
	Magnesium stearate	5.0

The above ingredients are mixed and filled into hard gelatin capsules in 340 mg quantities.

10

Formulation Example 2

A tablet formula is prepared using the ingredients below:

	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/tablet)</u>
15	Active Ingredient	25.0
	Cellulose, microcrystalline	200.0
	Colloidal silicon dioxide	10.0
	Stearic acid	5.0

20

The components are blended and compressed to form tablets, each weighing 240 mg.

Formulation Example 3

25 A dry powder inhaler formulation is prepared containing the following components:

	<u>Ingredient</u>	<u>Weight %</u>
30	Active Ingredient	5
	Lactose	95

35 The active mixture is mixed with the lactose and the mixture is added to a dry powder inhaling appliance.

Formulation Example 4

Tablets, each containing 30 mg of active ingredient, are prepared as follows:

-- 98 --

	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/tablet)</u>
5	Active Ingredient	30.0 mg
	Starch	45.0 mg
	Microcrystalline cellulose	35.0 mg
	Polyvinylpyrrolidone (as 10% solution in water)	4.0 mg
	Sodium carboxymethyl starch	4.5 mg
10	Magnesium stearate	0.5 mg
	Talc	<u>1.0 mg</u>
	Total	120 mg

15        The active ingredient, starch and cellulose are passed through a No. 20 mesh U.S. sieve and mixed thoroughly. The solution of polyvinyl-pyrrolidone is mixed with the resultant powders, which are then passed through a 16 mesh U.S. sieve. The granules so produced are dried at 50° to 60°C and passed through a 16 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium  
20        stearate, and talc, previously passed through a No. 30 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

#### Formulation Example 5

25        Capsules, each containing 40 mg of medicament are made as follows:

	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
30	Active Ingredient	40.0 mg
	Starch	109.0 mg
	Magnesium stearate	<u>1.0 mg</u>
	Total	150.0 mg

35        The active ingredient, cellulose, starch, an magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 150 mg quantities.

#### Formulation Example 6

-- 99 --

Suppositories, each containing 25 mg of active ingredient are made as follows:

5	<u>Ingredient</u>	<u>Amount</u>
	Active Ingredient	25 mg
	Saturated fatty acid glycerides to	2,000 mg

10 The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2.0 g capacity and allowed to cool.

15 Formulation Example 7

Suspensions, each containing 50 mg of medicament per 5.0 ml dose are made as follows:

20	<u>Ingredient</u>	<u>Amount</u>
	Active Ingredient	50.0 mg
	Xanthan gum	4.0 mg
	Sodium carboxymethyl cellulose (11%)	
	Microcrystalline cellulose (89%)	50.0 mg
25	Sucrose	1.75 g
	Sodium benzoate	10.0 mg
	Flavor and Color	q.v.
	Purified water to	5.0 ml

30 The medicament, sucrose and xanthan gum are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of the microcrystalline cellulose and sodium carboxymethyl cellulose in water. The sodium benzoate, flavor, and color are diluted with some of the water and  
35 added with stirring. Sufficient water is then added to produce the required volume.

-- 100 --

Formulation Example 8

	<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
5	Active Ingredient	15.0 mg
	Starch	407.0 mg
	Magnesium stearate	<u>3.0 mg</u>
10	Total	425.0 mg

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 560 mg quantities.

15 Formulation Example 9

An intravenous formulation may be prepared as follows:

	<u>Ingredient</u>	<u>Quantity</u>
20	Active Ingredient	250.0 mg
	Isotonic saline	1000 ml

Formulation Example 10

A topical formulation may be prepared as follows:

25	<u>Ingredient</u>	<u>Quantity</u>
	Active Ingredient	1-10 g
	Emulsifying Wax	30 g
	Liquid Paraffin	20 g
30	White Soft Paraffin	to 100 g

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active  
35 ~~ingredient is added and stirring is continued until dispersed.~~ The mixture is then cooled until solid.

Another preferred formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal

-- 101 --

patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991, herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

When it is desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system used for the transport of biological factors to specific anatomical regions of the body is described in U.S. Patent 5,011,472 which is herein incorporated by reference.

Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latentiation by the conversion of hydrophilic drugs into lipid-soluble drugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier. Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions which can transiently open the blood-brain barrier.

#### Utility

The compounds of this invention can be employed to bind VLA-4 ( $\alpha_4\beta_1$  integrin) in biological samples and, accordingly have utility in, for example, assaying such samples for VLA-4. In such assays, the compounds can be bound to a solid support and the VLA-4 sample added thereto. The amount of VLA-4 in the sample can be determined by conventional methods such as use of a sandwich ELISA assay. Alternatively, labeled VLA-4 can be used in a

-- 102 --

competitive assay to measure for the presence of VLA-4 in the sample. Other suitable assays are well known in the art.

In addition, certain of the compounds of this invention inhibit, *in vivo*,  
5 adhesion of leukocytes to endothelial cells mediated by VLA-4 and,  
accordingly, can be used in the treatment of diseases mediated by VLA-4.  
Such diseases include inflammatory diseases in mammalian patients  
such as asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes  
(including acute juvenile onset diabetes), inflammatory bowel disease  
10 (including ulcerative colitis and Crohn's disease), multiple sclerosis,  
rheumatoid arthritis, tissue transplantation, tumor metastasis, meningitis,  
encephalitis, stroke, and other cerebral traumas, nephritis, retinitis, atopic  
dermatitis, psoriasis, myocardial ischemia and acute leukocyte-mediated lung  
injury such as that which occurs in adult respiratory distress syndrome.

15 The biological activity of the compounds identified above may be assayed  
in a variety of systems. For example, a compound can be immobilized on a  
solid surface and adhesion of cells expressing VLA-4 can be measured. Using  
such formats, large numbers of compounds can be screened. Cells suitable for  
20 this assay include any leukocytes known to express VLA-4 such as T cells, B  
cells, monocytes, eosinophils, and basophils. A number of leukocyte cell lines  
can also be used, examples include Jurkat and U937.

The test compounds can also be tested for the ability to competitively  
25 inhibit binding between VLA-4 and VCAM-1, or between VLA-4 and a  
labeled compound known to bind VLA-4 such as a compound of this  
invention or antibodies to VLA-4. In these assays, the VCAM-1 can be  
immobilized on a solid surface. VCAM-1 may also be expressed as a  
recombinant fusion protein having an Ig tail (e.g., IgG) so that binding to  
30 VLA-4 may be detected in an immunoassay. Alternatively, VCAM-1  
expressing cells, such as activated endothelial cells or VCAM-1 transfected  
fibroblasts can be used. For assays to measure the ability to block adhesion to

-- 103 --

brain endothelial cells, the assays described in International Patent Application Publication No. WO 91/05038 are particularly preferred. This application is incorporated herein by reference in its entirety.

5           Many assay formats employ labelled assay components. The labelling systems can be in a variety of forms. The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. A wide variety of labels may be used. The component may be labelled by any one of several methods. The most common method of  
10       detection is the use of autoradiography with  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^{32}\text{P}$  labelled compounds or the like. Non-radioactive labels include ligands which bind to labelled antibodies, fluorophores, chemiluminescent agents, enzymes and antibodies which can serve as specific binding pair members for a labelled ligand. The choice of label depends on sensitivity required, ease of  
15       conjugation with the compound, stability requirements, and available instrumentation.

          Appropriate *in vivo* models for demonstrating efficacy in treating inflammatory responses include EAE (experimental autoimmune  
20       encephalomyelitis) in mice, rats, guinea pigs or primates, as well as other inflammatory models dependent upon  $\alpha 4$  integrins.

          Compounds having the desired biological activity may be modified as necessary to provide desired properties such as improved pharmacological  
25       properties (e.g., *in vivo* stability, bio-availability), or the ability to be detected in diagnostic applications. For instance, inclusion of one or more D-amino acids in the sulfonamides of this invention typically increases *in vivo* stability. Stability can be assayed in a variety of ways such as by measuring the half-life of the proteins during incubation with peptidases or human plasma or serum.  
30       A number of such protein stability assays have been described (see, e.g., Verhoef, et al., Eur. J. Drug Metab. Pharmacokinet., 1990, 15(2):83-93).



-- 104 --

For diagnostic purposes, a wide variety of labels may be linked to the compounds, which may provide, directly or indirectly, a detectable signal. Thus, the compounds of the subject invention may be modified in a variety of ways for a variety of end purposes while still retaining biological activity. In addition, various reactive sites may be introduced at the terminus for linking to particles, solid substrates, macromolecules, or the like.

Labeled compounds can be used in a variety of *in vivo* or *in vitro* applications. A wide variety of labels may be employed, such as radionuclides (e.g., gamma-emitting radioisotopes such as technetium-99 or indium-111), fluorescers (e.g., fluorescein), enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors, chemiluminescent compounds, bioluminescent compounds, and the like. Those of ordinary skill in the art will know of other suitable labels for binding to the complexes, or will be able to ascertain such using routine experimentation. The binding of these labels is achieved using standard techniques common to those of ordinary skill in the art.

*In vitro* uses include diagnostic applications such as monitoring inflammatory responses by detecting the presence of leukocytes expressing VLA-4. The compounds of this invention can also be used for isolating or labeling such cells. In addition, as mentioned above, the compounds of the invention can be used to assay for potential inhibitors of VLA-4/VCAM-1 interactions.

For *in vivo* diagnostic imaging to identify, e.g., sites of inflammation, radioisotopes are typically used in accordance with well known techniques. The radioisotopes may be bound to the peptide either directly or indirectly using intermediate functional groups. For instance, chelating agents such as diethylenetriaminepentaacetic acid (DTPA) and ethylenediaminetetraacetic acid (EDTA) and similar molecules have been used to bind proteins to metallic ion radioisotopes.

-- 105 --

The complexes can also be labeled with a paramagnetic isotope for purposes of *in vivo* diagnosis, as in magnetic resonance imaging (MRI) or electron spin resonance (ESR), both of which were well known. In general, any conventional method for visualizing diagnostic imaging can be used.

5 Usually gamma- and positron-emitting radioisotopes are used for camera imaging and paramagnetic isotopes are used for MRI. Thus, the compounds can be used to monitor the course of amelioration of an inflammatory response in an individual. By measuring the increase or decrease in lymphocytes expressing VLA-4 it is possible to determine whether a particular therapeutic  
10 regimen aimed at ameliorating the disease is effective.

The pharmaceutical compositions of the present invention can be used to block or inhibit cellular adhesion associated with a number of diseases and disorders. For instance, a number of inflammatory disorders are associated  
15 with integrins or leukocytes. Treatable disorders include, e.g., transplantation rejection (e.g., allograft rejection), Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes (including acute juvenile onset diabetes), retinitis, cancer metastases, rheumatoid arthritis, acute leukocyte-mediated lung injury (e.g., adult respiratory distress syndrome), asthma, nephritis, and acute and chronic  
20 inflammation, including atopic dermatitis, psoriasis, myocardial ischemia, and inflammatory bowel disease (including Crohn's disease and ulcerative colitis). In preferred embodiments the pharmaceutical compositions are used to treat inflammatory brain disorders, such as multiple sclerosis (MS), viral meningitis and encephalitis.

25 Inflammatory bowel disease is a collective term for two similar diseases referred to as Crohn's disease and ulcerative colitis. Crohn's disease is an idiopathic, chronic ulceroconstrictive inflammatory disease characterized by sharply delimited and typically transmural involvement of all layers of the  
30 bowel wall by a granulomatous inflammatory reaction. Any segment of the gastrointestinal tract, from the mouth to the anus, may be involved, although the disease most commonly affects the terminal ileum and/or colon.

-- 106 --

Ulcerative colitis is an inflammatory response limited largely to the colonic mucosa and submucosa. Lymphocytes and macrophages are numerous in lesions of inflammatory bowel disease and may contribute to inflammatory injury.

5

Asthma is a disease characterized by increased responsiveness of the tracheobronchial tree to various stimuli potentiating paroxysmal constriction of the bronchial airways. The stimuli cause release of various mediators of inflammation from IgE-coated mast cells including histamine, eosinophilic and neutrophilic chemotactic factors, leukotrienes, prostaglandin and platelet activating factor. Release of these factors recruits basophils, eosinophils and neutrophils, which cause inflammatory injury.

10

Atherosclerosis is a disease of arteries (e.g., coronary, carotid, aorta and iliac). The basic lesion, the atheroma, consists of a raised focal plaque within the intima, having a core of lipid and a covering fibrous cap. Atheromas compromise arterial blood flow and weaken affected arteries. Myocardial and cerebral infarcts are a major consequence of this disease. Macrophages and leukocytes are recruited to atheromas and contribute to inflammatory injury.

20

Rheumatoid arthritis is a chronic, relapsing inflammatory disease that primarily causes impairment and destruction of joints. Rheumatoid arthritis usually first affects the small joints of the hands and feet but then may involve the wrists, elbows, ankles and knees. The arthritis results from interaction of synovial cells with leukocytes that infiltrate from the circulation into the synovial lining of the joints. See e.g., Paul, *Immunology* (3d ed., Raven Press, 1993).

25

Another indication for the compounds of this invention is in treatment of organ or graft rejection mediated by VLA-4. Over recent years there has been a considerable improvement in the efficiency of surgical techniques for transplanting tissues and organs such as skin, kidney, liver, heart, lung,

30

-- 107 --

pancreas and bone marrow. Perhaps the principal outstanding problem is the lack of satisfactory agents for inducing immunotolerance in the recipient to the transplanted allograft or organ. When allogeneic cells or organs are transplanted into a host (i.e., the donor and donee are different individuals from the same species), the host immune system is likely to mount an immune response to foreign antigens in the transplant (host-versus-graft disease) leading to destruction of the transplanted tissue. CD8<sup>+</sup> cells, CD4 cells and monocytes are all involved in the rejection of transplant tissues. Compounds of this invention which bind to alpha-4 integrin are useful, *inter alia*, to block alloantigen-induced immune responses in the donee thereby preventing such cells from participating in the destruction of the transplanted tissue or organ. See, e.g., Paul et al., *Transplant International* 9, 420-425 (1996); Georczynski et al., *Immunology* 87, 573-580 (1996); Georczynski et al., *Transplant. Immunol.* 3, 55-61 (1995); Yang et al., *Transplantation* 60, 71-76 (1995); Anderson et al., *APMIS* 102, 23-27 (1994).

A related use for compounds of this invention which bind to VLA-4 is in modulating the immune response involved in "graft versus host" disease (GVHD). See e.g., Schlegel et al., *J. Immunol.* 155, 3856-3865 (1995). GVHD is a potentially fatal disease that occurs when immunologically competent cells are transferred to an allogeneic recipient. In this situation, the donor's immunocompetent cells may attack tissues in the recipient. Tissues of the skin, gut epithelia and liver are frequent targets and may be destroyed during the course of GVHD. The disease presents an especially severe problem when immune tissue is being transplanted, such as in bone marrow transplantation; but less severe GVHD has also been reported in other cases as well, including heart and liver transplants. The therapeutic agents of the present invention are used, *inter alia*, to block activation of the donor T-cells thereby interfering with their ability to lyse target cells in the host.

30

A further use of the compounds of this invention is inhibiting tumor metastasis. Several tumor cells have been reported to express VLA-4 and

-- 108 --

compounds which bind VLA-4 block adhesion of such cells to endothelial cells. Steinback et al., *Urol. Res.* 23, 175-83 (1995); Orosz et al., *Int. J. Cancer* 60, 867-71 (1995); Freedman et al., *Leuk. Lymphoma* 13, 47-52 (1994); Okahara et al., *Cancer Res.* 54, 3233-6 (1994).

5

A further use of the compounds of this invention is in treating multiple sclerosis. Multiple sclerosis is a progressive neurological autoimmune disease that affects an estimated 250,000 to 350,000 people in the United States. Multiple sclerosis is thought to be the result of a specific autoimmune reaction in which certain leukocytes attack and initiate the destruction of myelin, the insulating sheath covering nerve fibers. In an animal model for multiple sclerosis, murine monoclonal antibodies directed against VLA-4 have been shown to block the adhesion of leukocytes to the endothelium, and thus prevent inflammation of the central nervous system and subsequent paralysis in the animals<sup>16</sup>.

15

Pharmaceutical compositions of the invention are suitable for use in a variety of drug delivery systems. Suitable formulations for use in the present invention are found in *Remington's Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia, PA, 17th ed. (1985).

20

In order to enhance serum half-life, the compounds may be encapsulated, introduced into the lumen of liposomes, prepared as a colloid, or other conventional techniques may be employed which provide an extended serum half-life of the compounds. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka, et al., U.S. Patent Nos. 4,235,871, 4,501,728 and 4,837,028 each of which is incorporated herein by reference.

25

The amount administered to the patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions are administered to a patient already

30

-- 109 --

suffering from a disease in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the inflammation, the age, weight and general condition of the patient, and the like.

The compositions administered to a patient are in the form of pharmaceutical compositions described above. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

The therapeutic dosage of the compounds of the present invention will vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. For example, for intravenous administration, the dose will typically be in the range of about 20  $\mu$ g to about 500  $\mu$ g per kilogram body weight, preferably about 100  $\mu$ g to about 300  $\mu$ g per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.1  $\mu$ g to 1 mg per kilogram body weight. Effective doses can be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of

-- 110 --

this invention. Unless otherwise stated, all temperatures are in degrees Celsius.

5

## EXAMPLES

In the examples below, the following abbreviations have the following meanings. If an abbreviation is not defined, it has its generally accepted meaning.

	aq or aq.	=	aqueous
10	AcOH	=	acetic acid
	bd	=	broad doublet
	bm	=	broad multiplet
	bs	=	broad singlet
	Bn	=	benzyl
15	Boc	=	<i>N</i> - <i>tert</i> -butoxycarbonyl
	Boc <sub>2</sub> O	=	di- <i>tert</i> -butyl dicarbonate
	BOP	=	benzotriazol-1-yloxy- tris(dimethylamino)phosphonium hexafluorophosphate
20	Cbz	=	carbobenzyloxy
	CHCl <sub>3</sub>	=	chloroform
	CH <sub>2</sub> Cl <sub>2</sub>	=	dichloromethane
	(COCl) <sub>2</sub>	=	oxalyl chloride
	d	=	doublet
25	dd	=	doublet of doublets
	dt	=	doublet of triplets
	DBU	=	1,8-diazabicyclo[5.4.0]undec-7-ene
	DCC	=	1,3-dicyclohexylcarbodiimide
	DMAP	=	4- <i>N,N</i> -dimethylaminopyridine
30	DME	=	ethylene glycol dimethyl ether
	DMF	=	<i>N,N</i> -dimethylformamide
	DMSO	=	dimethylsulfoxide
	EDC	=	1-(3-dimethylaminopropyl)-3- ethylcarbodiimide hydrochloride
35	Et <sub>3</sub> N	=	triethylamine
	Et <sub>2</sub> O	=	diethyl ether
	EtOAc	=	ethyl acetate
	EtOH	=	ethanol
	eq or eq.	=	equivalent
40	Fmoc	=	<i>N</i> -(9-fluorenylmethoxycarbonyl)
	FmocONSu	=	<i>N</i> -(9-fluorenylmethoxycarbonyl)- succinimide
	g	=	grams

-- 111 --

	h	=	hour
	H <sub>2</sub> O	=	water
	HBr	=	hydrobromic acid
	HCl	=	hydrochloric acid
5	HOBT	=	1-hydroxybenzotriazole hydrate
	hr	=	hour
	K <sub>2</sub> CO <sub>3</sub>	=	potassium carbonate
	L	=	liter
	m	=	multiplet
10	MeOH	=	methanol
	mg	=	milligram
	MgSO <sub>4</sub>	=	magnesium sulfate
	mL	=	milliliter
	mm	=	millimeter
15	mM	=	millimolar
	mmol	=	millimol
	mp	=	melting point
	N	=	normal
	NaCl	=	sodium chloride
20	Na <sub>2</sub> CO <sub>3</sub>	=	sodium carbonate
	NaHCO <sub>3</sub>	=	sodium bicarbonate
	NaOEt	=	sodium ethoxide
	NaOH	=	sodium hydroxide
25	NH <sub>4</sub> Cl	=	ammonium chloride
	NMM	=	N-methylmorpholine
	Phe	=	L-phenylalanine
	Pro	=	L-proline
	psi	=	pounds per square inch
	PtO <sub>2</sub>	=	platinum oxide
30	q	=	quartet
	quint.	=	quintet
	rt	=	room temperature
	s	=	singlet
	sat	=	saturated
35	t	=	triplet
	t-BuOH	=	tert-butanol
	TFA	=	trifluoroacetic acid
	THF	=	tetrahydrofuran
	TLC or tlc	=	thin layer chromatography
40	Ts	=	tosyl
	TsCl	=	tosyl chloride
	TsOH	=	tosylate
	μL	=	microliter
45			



-- 112 --

In the examples below, all temperatures are in degrees Celcius (unless otherwise indicated). The following Methods were used to prepare the compounds set forth below as indicated.

5

## Method 1

N-Tosylation Procedure

N-Tosylation of the appropriate amino acid was conducted via the method of Cupps, Boutin and Rapoport *J. Org. Chem.* 1985, 50, 3972.

10

## Method 2

Methyl Ester Preparation Procedure

Amino acid methyl esters were prepared using the method of Brenner and Huber *Helv. Chim. Acta* 1953, 36, 1109.

15

## Method 3

BOP Coupling Procedure

The desired dipeptide ester was prepared by the reaction of a suitable N-protected amino acid (1 equivalent) with the appropriate amino acid ester or amino acid ester hydrochloride (1 equivalent), benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate [BOP] (2.0 equivalent), triethylamine (1.1 equivalent), and DMF. The reaction mixture was stirred at room temperature overnight. The crude product is purified flash chromatography to afford the dipeptide ester.

25

## Method 4

Hydrogenation Procedure I

Hydrogenation was performed using 10% palladium on carbon (10% by weight) in methanol at 30 psi overnight. The mixture was filtered through a pad of Celite and the filtrate concentrated to yield the desired amino compound.

30

-- 113 --

## Method 5

Hydrolysis Procedure I

To a chilled (0°C) THF/H<sub>2</sub>O solution (2:1, 5 - 10 mL) of the appropriate ester was added LiOH (or NaOH) (0.95 equivalents). The temperature was maintained at 0°C and the reaction was complete in 1-3 hours. The reaction mixture was extracted with ethyl acetate and the aqueous phase was lyophilized resulting in the desired carboxylate salt.

## Method 6

Ester Hydrolysis Procedure II

To a chilled (0°C) THF/H<sub>2</sub>O solution (2:1, 5 - 10 mL) of the appropriate ester was added LiOH (1.1 equivalents). The temperature was maintained at 0°C and the reaction was complete in 1-3 hours. The reaction mixture was concentrated and the residue was taken up into H<sub>2</sub>O and the pH adjusted to 2-3 with aqueous HCl. The product was extracted with ethyl acetate and the combined organic phase was washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated to yield the desired acid.

## Method 7

Ester Hydrolysis Procedure III

The appropriate ester was dissolved in dioxane/H<sub>2</sub>O (1:1) and 0.9 equivalents of 0.5 N NaOH was added. The reaction was stirred for 3-16 hours and then concentrated. The resulting residue was dissolved in H<sub>2</sub>O and extracted with ethyl acetate. The aqueous phase was lyophilized to yield the desired carboxylate sodium salt.

## Method 8

Sulfonylation Procedure I

To the appropriately protected aminophenylalanine analog (11.2 mmol), dissolved in methylene chloride (25ml) and cooled to -78°C was added the desired sulfonyl chloride (12 mmol) followed by dropwise addition of pyridine (2 mL). The solution was allowed to warm to room temperature and was

-- 114 --

stirred for 48 hr. The reaction solution was transferred to a 250 mL separatory funnel with methylene chloride (100 mL) and extracted with 1N HCl (50 mL x 3), brine (50 mL), and water (100 mL). The organic phase was dried (MgSO<sub>4</sub>) and the solvent concentrated to yield the desired product.

5

## Method 9

Reductive Amination Procedure

Reductive amination of Tos-Pro-p-NH<sub>2</sub>-Phe with the appropriate aldehyde was conducted using acetic acid, sodium triacetoxyborohydride, methylene chloride and the combined mixture was stirred at room temperature overnight. The crude product was purified by flash chromatography.

10

## Method 10

BOC Removal Procedure

Anhydrous hydrochloride (HCl) gas was bubbled through a methanolic solution of the appropriate Boc-amino acid ester at 0°C for 15 minutes and the reaction mixture was stirred for three hours. The solution was concentrated to a syrup and dissolved in Et<sub>2</sub>O and reconcentrated. This procedure was repeated and the resulting solid was placed under high vacuum overnight.

15

20

## Method 11

Tert-Butyl Ester Hydrolysis Procedure I

The *tert*-butyl ester was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and treated with TFA. The reaction was complete in 1-3 hr at which time the reaction mixture was concentrated and the residue dissolved in H<sub>2</sub>O and lyophilized to yield the desired acid.

25

## Method 12

EDC Coupling Procedure I

To a CH<sub>2</sub>Cl<sub>2</sub> solution (5-20 mL) of N-(toluene-4-sulfonyl)-L-proline (1 equivalent), the appropriate amino acid ester hydrochloride (1 equivalent), N-methylmorpholine (1.1-2.2 equivalents) and 1-hydroxybenzotriazole (2

30

-- 115 --

equivalents) were mixed, placed in an ice bath and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide (1.1 equivalents) added. The reaction was allowed to rise to room temperature and stirred overnight. The reaction mixture was poured into H<sub>2</sub>O and the organic phase was washed with sat. NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub> or Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude product was purified by column chromatography.

## Method 13

EDC Coupling Procedure II

To a DMF solution (5-20 mL) of the appropriate N-protected amino acid (1 equivalent), the appropriated amino acid ester hydrochloride (1 equivalent), Et<sub>3</sub>N (1.1 equivalents) and 1-hydroxybenzotriazole (2 equivalents) were mixed, placed in an ice batch and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide (1.1 equivalents) added. The reaction was allowed to rise to room temperature and stirred overnight. The reaction mixture was partitioned between EtOAc and H<sub>2</sub>O and the organic phase washed with 0.2 N citric acid, H<sub>2</sub>O, sat. NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub> or Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude product was purified by column chromatography or preparative TLC.

## Method 14

Sulfonylation Procedure II

The appropriate sulfonyl chloride was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and placed in an ice bath. L-Pro-L-Phe-OMe • HCl (1 equivalent) and Et<sub>3</sub>N (1.1 equivalent) was added and the reaction allowed to warm to room temperature and stirred overnight under an atmosphere of nitrogen. The reaction mixture was concentrated and the residue partitioned between EtOAc and H<sub>2</sub>O and the organic phase washed with sat. NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub> or Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude product was purified by column chromatography or preparative TLC.

## Method 15

-- 116 --

Sulfonylation Procedure III

To a solution of L-Pro-L-4-(3-dimethylaminopropoxy)-Phe-OMe [prepared using the procedure described in Method 10] (1 equivalent) in  $\text{CH}_2\text{Cl}_2$  was added  $\text{Et}_3\text{N}$  (5 equivalents) followed by the appropriate sulfonyl chloride (1.1 equivalent). The reaction was allowed to warm to room temperature and stirred overnight under an atmosphere of nitrogen. The mixture was concentrated, dissolved in EtOAc, washed with sat.  $\text{NaHCO}_3$  and 0.2 N citric acid. The aqueous phase was made basic with solid  $\text{NaHCO}_3$  and the product extracted with EtOAc. The organic phase was washed with brine, dried ( $\text{MgSO}_4$  or  $\text{Na}_2\text{SO}_4$ ), filtered and concentrated. The crude methyl ester was purified by preparative TLC. The corresponding acid was prepared using the procedure described in Method 7.

## Method 16

Hydrogenation Procedure II

To a methanol (10 -15 mL) solution of the azlactone, was added NaOAc (1 equivalent) and 10% Pd/C. This mixture was placed on the hydrogenator at 40 psi  $\text{H}_2$ . After 8 - 16 hours, the reaction mixture was filtered through a pad of Celite and the filtrate concentrated to yield the dehydrodipeptide methyl ester. The ester was dissolved in dioxane/ $\text{H}_2\text{O}$  (5- 10 mL), to which was added 0.5 N NaOH (1.05 equivalents). After stirring for 1- 3 hours, the reaction mix was concentrated and the residue was redissolved in  $\text{H}_2\text{O}$  and washed with EtOAc. The aqueous phase was made acidic with 0.2 N HCl and the product was extracted with EtOAc. The combined organic phase was washed with brine (1 x 5 mL), dried ( $\text{MgSO}_4$  or  $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to yield the acid as approximately a 1:1 mixture of diastereomers.

## Method 17

Tert-Butyl Ester Hydrolysis Procedure II

The *tert*-butyl ester was dissolved in  $\text{CH}_2\text{Cl}_2$  (5 mL) and treated with TFA (5 mL). The reaction was complete in 1-3 hours at which time the reaction mixture was concentrated and the residue dissolved in  $\text{H}_2\text{O}$  and concentrated.

-- 117 --

The residue was redissolved in H<sub>2</sub>O and lyophilized to yield the desired product.

## Example 1 (2)

5

**Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(*N*-*tert*-butoxycarbonylglycyl)amino]-L-phenylalanine**

*N*-(Toluene-4-sulfonyl)-L-prolyl-L-4-aminophenylalanine methyl ester  
10 (2.00 g, 4.67 mmol) was dissolved 50 mL of dry DMF, with Boc-glycine (1.1 eq, 0.9 g), Et<sub>3</sub>N (2.2 eq, 1.43 mL), and BOP reagent (1.1 eq, 2.27g). The reaction mixture was stirred at room temperature for 12 hours. EtOAc (100 mL) was added. The organic layer was washed with NaHCO<sub>3</sub> saturated (50 mL), 10% citric acid (20 mL), and brine (5x 50 mL). The organic layer was  
15 dried over MgSO<sub>4</sub>. Upon evaporation of the solvents under reduced pressure, the crude material was eluted on column chromatography (silica gel; CHCl<sub>3</sub>/MeOH 9:1). The desired ester (1.90 g, 3.1 mmol) was isolated in 66% yield. The ester was then taken up in MeOH:H<sub>2</sub>O [1:1] (40 mL), with NaOH (1.1 eq, 176 mg), for 4 hours at room temperature. EtOAc was added as well  
20 as water. The aqueous layer was collected and acidified with 1N HCl to pH 2.5, and reextracted with EtOAc. The organic layer was dried over MgSO<sub>4</sub>. Upon filtration and evaporation of the solvents under reduced pressure, the title compound was isolated in 73% yield, as an oil (1.33g, 2.26 mmol).

NMR data was as follows:

25 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 8.65 (bs, 1H), 7.70 (d, 2H, *J* = 7.98 Hz), 7.50 (d, 1H, *J* = 7.14 Hz), 7.45 (d, 2H, *J* = 8.10 Hz), 7.32 (d, 2H, *J* = 7.80 Hz), 7.11 (d, 2H, *J* = 8.00 Hz), 5.75 (bs, 1H), 4.82 (m, 1H), 4.11 (m, 1H), 3.90 (bs, 2H), 3.38 (m, 1H), 3.21 3.10 (m, 3H), 2.41 (s, 3H), 1.99 (m, 1H), 1.55 (m, 3H), 1.43 (s, 9H).

30 <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ = 174.03, 172.19, 169.02, 157.28, 145.03, 137.17, 133.34, 132.58, 130.59, 128.44, 120.65, 81.16, 62.85, 54.05, 50.31, 37.34, 30.69, 28.87, 24.89, 22.14, 14.77.

Mass Spectroscopy: (FAB) 589 (M+H).

-- 118 --

## Example 2 (7)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(glycyl)amino]-L-phenylalanine

5 The product from Example 1 (2) (1.33 g, 2.26 mmol) was taken up in dry dichloromethane (20 mL) with trifluoroacetic acid (1.00 mL) and the reaction mixture was stirred at room temperature for 12 hours. The solvents were removed under reduced pressure. The crude material was left on a vacuum  
10 pump overnight, then dissolved in methanol and cooled to 0°C. Ethyl ether was added and the product was collected, as a trifluoroacetate salt, in 50% yield (660 mg, 1.09 mmol).

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD): δ = 7.51 (d, 2H, *J* = 8.25 Hz), 7.31 (d, 2H, *J* = 8.25 Hz), 7.19 (d, 2H, *J* = 7.92 Hz), 7.03 (d, 2H, *J* = 8.25 Hz), 4.43 (m, 1H), 3.87 (m, 1H), 3.63 (s, 2H), 3.13-2.82 (m, 4H), 2.21 (s, 3H), 1.56-1.26 (m, 4H).  
15

<sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD): δ = 174.39, 165.97, 146.34, 139.50, 135.38, 135.21, 131.28, 129.57, 121.47, 67.47, 63.86, 51.16, 42.72, 38.39, 32.20, 25.87, 22.09, 16.02.  
20

Mass Spectroscopy: (FAB) 489 (M<sup>+</sup>).

## Example 3 (23)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[3-(carboxy)propionamido]-L-phenylalanine

25 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-aminophenylalanine methyl ester (455 mg, 1 mmol) was dissolved in DMF (10 mL) with Et<sub>3</sub>N (1.1 eq, 153 μL) and succinic anhydride (1.1 eq, 110 mg). The desired monoester was isolated  
30 in 53% yield (288 mg, 0.52 mmol) as a foam. The monoester (80 mg, 0.15 mmol) was then hydrolyzed in MeOH: H<sub>2</sub>O 1:1 (5 mL) with NaOH (2.2 eq, 15 mg) for 4 hours at room temperature. EtOAc was added as well as water. The aqueous layer was collected and acidified with 1N HCl to pH 2.5, and  
35 reextracted with EtOAc. The organic layer was dried over MgSO<sub>4</sub>. Upon

-- 119 --

filtration and evaporation of the solvents under reduced pressure, the diacid was isolated as a foam.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD): δ = 7.51 (d, 2H, *J* = 8.31 Hz), 7.28 (d, 2H, *J* = 8.34 Hz), 7.19 (d, 2H, *J* = 8.10 Hz), 6.99 (d, 2H, *J* = 8.43 Hz), 4.48 (m, 1H), 3.91 (m, 1H), 3.17-2.83 (m, 4H), 2.43 (s, 4H), 2.21 (s, 3H), 1.44-1.29 (m, 4H).

<sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD): δ = 176.87, 174.68, 174.53, 173.30, 149.74, 146.31, 139.33, 135.50, 134.24, 131.61, 131.44, 129.53, 121.66, 80.05, 63.81, 55.37, 51.17, 38.25, 32.87, 32.24, 30.55, 25.89, 22.11.

Mass Spectroscopy: (FAB) 532 (M+H).

#### Example 4 (27)

#### Synthesis of 15 *N*-(Toluene-4-sulfonyl)-L-prolyl-4- [(*N*-tert-butoxycarbonyl-L-alanyl)amino]-L-phenylalanine

Following the experimental procedure described for Example 1 (2), *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-aminophenylalanine methyl ester (0.15 g, 0.337 mmol) was taken into 5 mL of DMF, with Et<sub>3</sub>N (2.0 eq, 95 μL), Boc-L-alanine (1.1 eq, 70 mg), and BOP (1.1 eq, 163 mg). The desired methyl ester was isolated as an oil in 49% yield (102 mg, 0.16 mmol). The ester was then hydrolyzed with NaOH (1.1 eq, 8 mg) in a 1:1 MeOH:H<sub>2</sub>O solution (4 mL). The monoacid was isolated as an amorphous solid in quantitative yields.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 9.00 (broad s, 1H), 7.70 (d, 2H, *J* = 8.25 Hz), 7.45 (d, 2H, *J* = 8.37 Hz), 7.32 (d, 2H, *J* = 8.25 Hz), 7.09 (d, 2H, *J* = 8.25 Hz), 5.65 (bs, 1H), 4.82 (m, 1H), 4.35 (bs, 1H), 4.11 (m, 1H), 3.38-3.12 (m, 4H), 2.40 (s, 3H), 1.95 (m, 1H), 1.53 (m, 3H), 1.40 (s, 12H).

<sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ = 174.71, 172.40, 172.09, 156.72, 144.98, 137.62, 133.43, 132.21, 130.58, 128.48, 120.47, 80.98, 62.87, 54.05, 51.10, 50.31, 37.23, 30.73, 28.88, 24.92, 22.16, 18.89.

Mass Spectroscopy: (FAB) 603 (M+H).



-- 120 --

## Example 5 (28)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(*N*-*tert*-butoxycarbonyl-D-alanyl)amino]-L-phenylalanine

5

Following the experimental procedure described for Example 1 (2), *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-aminophenylalanine methyl ester (0.15 g, 0.337 mmol) was taken into 5 mL of DMF, with Et<sub>3</sub>N (2.0 eq, 95  $\mu$ L), Boc-D-alanine (1.1 eq, 70 mg), and BOP (1.1 eq, 163 mg). The desired methyl ester was isolated as an oil in 33% yield (68 mg, 0.11 mmol). The ester was then hydrolyzed with NaOH (1.1 eq, 5 mg) in a 1:1 MeOH:H<sub>2</sub>O solution (4 mL). The monoacid was isolated as a film in quantitative yields.

10

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 8.90 (broad s, 1H), 7.70 (d, 2H, *J* = 8.25 Hz), 7.45 (d, 2H, *J* = 8.37 Hz), 7.32 (d, 2H, *J* = 8.25 Hz), 7.09 (d, 2H, *J* = 8.25 Hz), 5.65 (bs, 1H), 4.82 (m, 1H), 4.35 (bs, 1H), 4.11 (m, 1H), 3.38-3.12 (m, 4H), 2.40 (s, 3H), 1.95 (m, 1H), 1.53 (m, 3H), 1.40 (s, 12H).

15

<sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  = 174.71, 172.40, 172.09, 156.72, 144.98, 137.62, 133.43, 132.21, 130.58, 128.48, 120.47, 80.98, 62.87, 54.05, 51.10, 50.31, 37.23, 30.73, 28.88, 24.92, 22.16, 18.89.

20

Mass Spectroscopy: (FAB) 603 (M+H).

## Example 6 (29)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(*N*-*tert*-butoxycarbonyl-D-phenylalanyl)amino]-L-phenylalanine

25

Following the experimental procedure described for Example 1 (2), *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-aminophenylalanine methyl ester (0.15 g, 0.337 mmol) was taken into 5 mL of DMF, with Et<sub>3</sub>N (2.0 eq, 95  $\mu$ L), Boc-L-phenylalanine (1.1 eq, 99 mg), and BOP (1.1 eq, 163 mg). The desired methyl ester was isolated as an oil in 16% yield (37 mg, 0.05 mmol). The ester was then hydrolyzed with NaOH (1.1 eq, 3 mg) in a 1:1 MeOH:H<sub>2</sub>O solution (2 mL). The monoacid was isolated as an amorphous solid in quantitative yields.

30

NMR data was as follows:

35

-- 121 --

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 8.56 (broad s, 1H), 7.73 (d, 2H, J = 8.13 Hz), 7.43-7.22 (m, 10 H), 7.09 (d, 2H, J = 7.95 Hz), 5.78 (m, 1H), 4.87 (m, 1H), 4.65 (m, 1H), 4.11 (m, 1H), 3.44 (m, 1H), 3.29-3.01 (m, 4H), 2.43 (s, 3H), 2.08 (m, 1H), 1.60 (m, 3H), 1.33 (s, 9H).

<sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ = 171.88, 171.44, 170.02, 156.56, 144.96, 137.23, 136.83, 133.46, 132.78, 130.53, 130.33, 129.92, 129.36, 128.44, 127.64, 120.63, 81.16, 62.84, 53.96, 50.26, 38.91, 37.90, 30.35, 28.81, 24.96, 22.11.

Mass Spectroscopy: (FAB) 679 (M+H).

#### Example 7 (67)

##### Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-4- {2-[3-(fluorescein)thiouriedo]acetamido}-L-phenylalanine

The product of Example 2 (7) was treated with NaHCO<sub>3</sub> and fluorescein isothiocyanate (isomer I -- obtained from Aldrich Chemical Company) in EtOH and H<sub>2</sub>O. Acidification of the mixture and isolation of the resulting precipitate gave the title compound as an amorphous orange solid.

Mass Spectroscopy: (+FAB, 3-nitrobenzyl alcohol) 878 (MH+).

#### Example 8 (175)

##### Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-4- [(*N*-*tert*-butoxycarbonyl)glycyl]amino]-L-phenylalanine Methyl Ester

*N*-(Toluene-4-sulfonyl)-L-prolyl-L-(4-nitro)phenylalanine methyl ester (2.00 g, 4.48 mmol) was dissolved in MeOH (10 mL) with a catalytic amount of 10% Pd on C. The hydrogenation reaction was run at room temperature for 12 hours at 40 psi. Upon filtration of the reaction mixture over celite, the solvent was evaporated under reduced pressure yielding a pink foam in quantitative yields. The title compound was prepared therefrom and N-Boc glycine using the procedure described in Method 12.

NMR data was as follows:

-- 122 --

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 7.72 (d, 2H, J = 8.13 Hz), 7.45 (d, 2H, J = 8.49 Hz), 7.35 (d, 2H, J = 8.30 Hz), 7.10 (d, 2H, J = 8.37 Hz), 5.40 (m, 1H), 4.82 (m, 1H), 4.07 (m, 1H), 3.91 (s, 2H), 3.76 (s, 3H), 3.40 (m, 1H), 3.23 (m, 1H), 3.05 (m, 2H), 2.43 (s, 3H), 2.43 (m, 1H), 1.46 (m, 12H).

5 <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ = 177.61, 171.97, 171.58, 168.52, 144.93, 137.25, 133.43, 132.55, 131.36, 130.54, 130.33, 128.39, 120.53, 80.50, 62.81, 54.00, 53.08, 50.25, 37.83, 30.42, 28.93, 28.87, 24.81, 22.14.

Mass Spectroscopy: (FAB) 603 (M+H).

10

## Example 9 (306)

## Synthesis of

*N*-(Toluene-4-sulfonyl)-L-prolyl-4-(2-[3-(3-methylphenyl)uriedo]acetamido)-L-phenylalanine

15 The methyl ester was prepared by the reaction of *N*-(toluene-4-sulfonyl)-L-prolyl-L-(4-amino)phenylalanine with 2-(3-*m*-tolyl-ureido)acetic acid (BOP, triethylamine, DMF, stir at room temperature overnight). The crude product was purified flash chromatography (silica, 9:1 EtOAc:hexane) to afford the methyl ester. The methyl ester was hydrolyzed using 1M LiOH in THF. The  
20 title compound was isolated following acid/base work-up as a white solid.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz): δ = 12.8 (br s, 1H); 9.96 (s, 1H); 8.70 (s, 1H); 8.04 (d, 1H, J = 7.9Hz); 7.68 (d, 2H, J = 8Hz); 7.48 (d, 2H, J = 8.3Hz); 7.39 (d, 2H, J = 8.3Hz); 7.21 (s, 1H); 7.16 (d, 3H, J = 8.56Hz); 7.08 (t, 1H, J =  
25 7.79Hz); 6.7 (d, 1H, J = 7.68Hz); 6.38 (t, 1H, J = 5.6Hz); 4.43 (m, 1H); 4.1 (dd, 1H, J = 3.18, 8.23Hz); 3.89 (d, 2H, J = 5.49Hz); 3.3 (m, 1H); 3.05 (m, 2H); 2.92 (dd, 1H, J = 8.34, 13.63Hz); 2.38 (s, 3H); 2.22 (s, 3H); 1.38-1.62 (m, 4H).

IR (KBr, cm<sup>-1</sup>): 3380, 2990, 1650, 1605, 1540, 1230, 1155, 660, 580, 545.

30

Mass Spectroscopy: ( (+) FAB, m/e (%) ) 644 (100 [M+Na]<sup>+</sup>); 622 (25 [M+H]<sup>+</sup>).

-- 123 --

## Example 10 (380)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[γ-(L-aspartyl)amino]-L-phenylalanine

5 Substitution of Boc-L-aspartic benzyl ester for Boc-glycine and following the methods for preparation of Examples 1 (2) and 2 (7), gave the title compound as a white solid, mp = 163-170°C.

NMR data was as follows:

10 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ = 8.05 (d, 1H), 7.70 (d, 1H), 7.45 (dd, 4H), 7.18 (d, 2H), 4.40 (m, 1H), 4.10 (m, 1H), 2.40 (s, 3H), 1.60 (m, 6H)

Mass Spectroscopy: (FAB) (M+H)<sup>+</sup> 547.

## Example 11 (14)

15 Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
(α-carboxybenzyloxy)-L-phenylalanine

*N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester (1.32 g, 2.95 mmol) was dissolved in dry DMF (50 mL) at room temperature. To this was added K<sub>2</sub>CO<sub>3</sub> (1.1 eq, 440 mg) and ethyl α-bromophenylacetate (1.1 eq, 750 mg). The reaction was stirred for 12 hours at room temperature. Ethyl acetate (100 mL) was added, and the organic layer washed several times with brine. The organic layer was dried over MgSO<sub>4</sub>. Upon filtration and evaporation of the solvents under reduced pressure, a residue was isolated which was then taken up in MeOH: H<sub>2</sub>O 1:1 (30 mL) with NaOH (1.1 eq, 427 mg). The desired diacid was isolated in 88% yield (754 mg, 1.33 mmol).

NMR data was as follows:

30 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) : δ = 7.50 (d, 2H, *J* = 8.25 Hz), 7.35 (d, 2H, *J* = 7.14 Hz), 7.15 (m, 5H), 6.95 (d, 2H, *J* = 8.13 Hz), 6.71 (d, 2H, *J* = 8.37 Hz), 5.48 (s, 1H), 4.51 (m, 1H), 3.87 (m, 1H), 3.10-2.79 (m, 4H), 2.15 (s, 3H), 1.45-1.01 (m, 4H).

<sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD): δ = 174.71, 174.49, 173.98, 158.25, 146.39, 138.08, 135.37, 132.28, 131.74, 130.53, 130.36, 129.60, 129.00, 117.18, 79.95, 63.94, 55.31, 38.00, 32.23, 25.88, 22.32, 15.22.

**Mass Spectroscopy:** (FAB) 567 (M+H).

## Synthesis of

## 5

10

15

20

## Synthesis of

## 25

30

35

-- 125 --

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ = 7.81 (d, J=7.7, 1H), 7.72 (d, J=8.2, 2H), 7.54-7.49 (m, 1H), 7.42-7.16 (m, 9H), 4.93-4.86 (m, 1H), 4.11-4.07 (m, 1H), 3.80 (s, 3H), 3.64 (s, 3H), 3.42-3.28 (m, 2H), 3.15-3.08 (m, 2H), 2.43 (s, 3H), 2.09-2.04 (m, 1H), 1.58-1.45 (m, 3H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz): δ = 171.3, 170.8, 169.1, 144.3, 141.9, 140.1, 135.0, 132.9, 131.2, 130.8, 130.6, 129.9, 129.7, 129.0, 128.4, 127.8, 127.1, 62.2, 53.3, 52.5, 52.0, 49.7, 37.6, 29.7, 24.2, 21.5.

Mass Spectroscopy: (+FAB, 3-nitrobenzyl alcohol) 565 (MH<sup>+</sup>).

10

## Example 14 (35)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine

15

Following the experimental procedure described for the synthesis of Example 3 (23), the product from Example 15 (172) (32 mg, 0.051 mmol) was hydrolyzed in MeOH:H<sub>2</sub>O 1:1 (1 mL), with NaOH (1.1 eq, 3 mg). The title material was isolated in quantitative materials, as a foam.

NMR data was as follows:

20

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 7.68 (m, 2H), 7.30 (m, 5H), 7.20 (m, 2H), 4.77 (m, 1H), 4.46 (m, 1H), 4.18 (m, 2H), 3.65 (m, 1H), 3.47 (m, 1H), 3.12 (m, 2H), 2.78 (s, 1.5H), 2.54 (s, 1.5H), 2.42 (s, 3H), 1.26 (m, 3H).

<sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ = 171.47, 170.31, 144.78, 136.42, 135.77, 130.67, 129.88, 129.33, 127.80, 62.44, 60.99, 60.81, 54.00, 53.77, 38.32, 32.19, 31.96, 22.24, 14.76.

25

Mass Spectroscopy: (FAB) 449 (M+H).

## Example 15 (172)

30

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine  
Methyl Ester

*N*-(Toluene-4-sulfonyl)-L-prolyl-L-p-amino-phenylalanine methyl ester

35

(316 mg, 0.7 mmol) was dissolved in a KOAc-HOAc solution in dry methanol

-- 126 --

(10 mL) [pH 6.5], with NaCNBH<sub>3</sub> (10.0 eq, 446 mg), and carbobenzyloxy-glycinal (1.1 eq). The reaction mixture was stirred at room temperature overnight, and the solvents were evaporated under reduced pressure. EtOAc was added and the organic layer washed with brine, dried over MgSO<sub>4</sub>. Upon filtration and evaporation of the solvents, the crude material was eluted on a preparative plate (Silica gel, EtOAc/hexanes 1:1). The desired title compound was isolated, as a film in 5% yield (40 mg, 0.06 mmol).

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 7.71 (d, 2H, J = 6.90 Hz), 7.38 (m, 5H), 7.26 (m, 2H), 6.93 (d, 2H, J = 7.20 Hz), 6.54 (d, 2H, J = 7.50 Hz), 5.12 (m, 1H), 5.10 (s, 2H), 4.73 (m, 1H), 4.06 (m, 1H), 3.75 (s, 3H), 3.41 (m, 4H), 3.26 (m, 1H), 3.14 (m, 2H), 2.99 (m, 1H), 2.43 (s, 3H), 2.04 (m, 1H), 1.53 (m, 3H).

Mass Spectroscopy: (FAB) 623 (M+H).

#### Example 16 (368)

##### Synthesis of

*N*-(Toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[3-(*N,N*-dimethylamino)propyl]-*N*-(trifluoromethanesulfonyl)amino]-L-phenylalanine Methyl Ester

*N*-(Toluene-4-sulfonyl)-L-prolyl-L-(4-amino)phenylalanine methyl ester was reacted with trifluoromethane sulfonyl anhydride in pyridine to produce the corresponding trifluoromethane sulfonamide. This compound was alkylated on the nitrogen of the trifluoromethane sulfonamide group under Mitsunobu conditions using 3-dimethylamino-1-propanol in THF. After solvent evaporation and aqueous wash the mixture was purified by filtration to yield the product as a solid, mp = 45-55°C.

NMR data was as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400Mhz): δ = 7.70 (d, 2H); 7.34 (d, 3H); 7.25 (m, 3H); 4.88 (m, 1H); 4.10 (m, 1H); 3.85 (brd s, 2H); 3.79 (s, 3H); 3.32 (m, 2H); 3.08 (m, 2H); 2.43 (s, 3H); 2.18 (brd s, 5H); 1.98 (brd s, 1H); 1.75-1.30 (brd m, 9H).

-- 127 --

IR (KBr, cm<sup>-1</sup>): 3400; 2970; 2800; 1750; 1675; 1525; 1450; 1400; 1350; 1225; 1200; 1165; 1150; 1100; 1075; 1000; 950; 825; 675; 600; 550.

Mass Spectroscopy: (+FAB) 663 ( [M+H]<sup>+</sup> ); 603; 507; 438; 306; 191; 155; 91.

5

## Example 17 (510)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-*L*-prolyl-  
L-4-{*N,N*-di[4-(*N,N*-dimethylamino)benzyl]amino}-  
L-phenylalanine

10

The methyl ester was prepared by reductive amination of *N*-(toluene-4-sulfonyl)-*L*-prolyl-L-4-aminobenzyl-L-phenylalanine with 4-*N,N*-dimethylaminobenzaldehyde (acetic acid, sodium triacetoxyborohydride, methylene chloride), stirred at room temperature overnight. The crude product was purified by flash chromatography to afford the methyl ester which was hydrolyzed in the manner of Method 6 to provide the title compound.

15

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 7.7 (d, 2H, J = 8.34Hz); 7.54 (d, 1H, J = 5.71 Hz); 7.40 (d, 1H, J = 8.34Hz); 7.02 (d, 4H, J = 8.78Hz); 6.75 (d, 2H, J = 8.78Hz); 6.62 (d, 4H, J = 8.78Hz); 6.48 (d, 2H, J = 8.78Hz); 4.39 (s, 4H); 3.92 (dd, 1H, J = 2.85, 9.0Hz); 3.78 (m, 1H); 3.33 (s, 12H); 2.86 (m, 4H); 2.39 (s, 3H); 1.62 (m, 1H); 1.30 (m, 1H); 1.09 (m, 2H).

20

IR (KBr, cm<sup>-1</sup>) 3380, 1610, 1520, 1400, 1350, 1160, 800, 670.

25

MS ((+) FAB, m/e (%)) 698 (20[M+H]<sup>+</sup>).

Anal. Calc'd for C<sub>39</sub>H<sub>46</sub>N<sub>5</sub>O<sub>5</sub>Li 3H<sub>2</sub>O: C, 61.76; H, 6.91; N, 9.23. Found: C, 61.73; H, 6.91; N, 9.15.

## Example 18 (138)

30

Synthesis of  
*N*-(Toluene-4-sulfonyl)-*L*-prolyl-4-  
[3-(*N,N*-dimethylamino)propoxy]-*L*-phenylalanine

Triphenylphosphine (24.4 g, 92.9 mmol) and 3-dimethylamino-1-propanol (8.72 g, 84.5 mmol) were dissolved in THF (200 mL) and cooled in

35



-- 128 --

an ice bath. Diethyl azodicarboxylate (16.2 g, 14.6 mL, 92.9 mmol) was added dropwise via syringe over 5 minutes and stirred an additional 10 minutes before addition of *N*-Boc-tyrosine methyl ester (24.95 g, 94.5 mmol) as a solution in 100 mL THF via cannula. The mixture was stirred for 30 minutes at 0°C and 19 hr at room temperature. The mixture was concentrated on a rotary evaporator and taken up into Et<sub>2</sub>O (500 mL). The mixture was extracted with 0.2N HCl (3 x 350 mL) and the combined acidic extracts were made basic with solid NaHCO<sub>3</sub>. This mixture was extracted with EtOAc (3 x 300 mL) and the combined extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated in vacuo to give *N*-Boc-L-4-(3-dimethylaminopropoxy)phenylalanine methyl ester (26.5 g, 82 %).

*N*-Boc-L-4-(3-*N,N*-dimethylaminopropoxy)phenylalanine methyl ester (26.5 g, 69.5 mmol) was dissolved in MeOH (300 mL) and saturated with HCl gas. The mixture was stirred for 3 hr before removing the volatiles *in vacuo* to give L-4-(3-*N,N*-dimethylaminopropoxy)phenylalanine methyl ester dihydrochloride (23.6 g, 90%).

*N*-(Toluene-4-sulfonyl)-L-proline hydrate was coupled to L-4-(3-*N,N*-dimethylaminopropoxy)-phenylalanine methyl ester dihydrochloride using the procedure described in Method 13 to give *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-(3-*N,N*-dimethylaminopropoxy)phenylalanine methyl ester (16.3g, 89%). The title compound was prepared via hydrolysis of the methyl ester using 0.5 N NaOH in THF/water (14.71 g, 99%).

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ = 7.75 (d, 2H, *J* = 8.2 Hz), 7.67 (d, 1H, *J* = 5.6 Hz), 7.42 (d, 2H, *J* = 8.2 Hz), 7.00 (d, 2H, *J* = 8.5 Hz), 6.71 (d, 2H, *J* = 8.5 Hz), 3.98 (m, 2H), 3.90 (t, 2H, *J* = 6.5 Hz), 3.14-2.97 (4H), 2.40 (s, 3H), 2.32 (t, 2H, *J* = 7.0 Hz), 2.13 (s, 6H), 1.83-1.75 (3H), 1.45-1.36 (3H).

<sup>13</sup>C NMR (DMSO-d<sub>6</sub>): δ = 173.5, 169.7, 157.2, 144.1, 133.7, 130.9, 130.3, 128.1, 113.9, 65.9, 62.4, 56.0, 55.4, 49.4, 45.5, 36.1, 30.5, 27.3, 23.9, 21.4.

-- 129 --

**Mass Spectroscopy: FAB m/e 562 (M+2Na-H).**

### Example 19 (282)

## Synthesis of

***N*-(Toluene-4-sulfonyl)-*N*-methyl-L-serinyl-4-[3-(*N,N*-dimethylaminopropoxy)]-L-phenylalanine Methyl Ester**

*N*-Methyl-*N*-(toluene-*p*-sulfonyl)-L-serine (655 mg, 2.4 mmol) was taken up in DMF (100 mL) with L-4-(3-dimethylamino-propyloxy)phenylalanine methyl ester hydrochloride salt [see Example 18 (138) for preparation] (1.0 g, 2.4 mmol), HOBT (1.1 eq, 356 mg), Et<sub>3</sub>N (3.2 eq, 1.1 mL), and EDC (1.1 eq, 500 mg) following the procedure described in Method 13. The title compound was isolated in 70% yield (900 mg, 1.7 mmol) as an oil.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 7.68 (d, 2H, J = 8.40 Hz), 7.29 (d, 2H, J = 8.10 Hz), 7.08 (d, 1H, J = 7.80 Hz), 7.04 (d, 2H, J = 8.70 Hz), 6.82 (d, 2H, J = 8.70 Hz), 4.71 (m, 1H), 4.43 (m, 1H), 3.94 (t, 2H, J = 6.60 Hz), 3.75 (s, 3H), 3.69 (m, 1H), 3.51 (m, 1H), 3.14 (m, 1H), 2.91 (m, 1H), 2.59 (s, 3H), 2.45 (m, 2H), 2.42 (s, 3H), 2.20 (s, 6H), 1.87 (m, 2H).

<sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ = 172.04, 170.15, 158.66, 144.56, 136.02, 130.72, 130.46, 128.19, 127.87, 115.31, 66.70, 60.86, 60.65, 56.87, 54.01, 53.08, 45.95, 37.39, 31.85, 27.91, 22.13.

Mass Spectroscopy: (FAB) 536 (M+H).

### Example 20 (284)

## Synthesis of

***N*-(Toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine**

Triphenylphosphine (1.1 eq) and 3-dimethylamino-1-propanol (1 eq) were dissolved in THF and cooled in an ice bath. Diethyl azodicarboxylate (1 eq) was added dropwise via syringe over 5 min and stirred an additional 10 min before addition of *N*-Boc-tyrosine methyl ester (1.02 eq) as a solution in 100 mL THF via cannula. The mixture was stirred for 30 min at 0 °C and 19 hr at room temperature. The mixture was concentrated on a rotary evaporator and

-- 130 --

taken up into Et<sub>2</sub>O (500 mL). The mixture was extracted with 0.2N HCl (3 X 350 mL) and the combined acidic extracts were made basic with solid NaHCO<sub>3</sub>. This mixture was extracted with EtOAc (3 X 300 mL) and the combined extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated in vacuo to give *N*-Boc-L-4-(3-dimethylaminopropoxy)phenylalanine methyl ester.

*N*-Boc-L-4-(3-dimethylaminopropoxy)phenylalanine methyl ester (26.5 g, 69.5 mmol) was dissolved in MeOH (300 mL) and saturated with HCl gas. The mixture was stirred for 3 hr before removing the volatiles in vacuo to give L-4-(3-dimethylaminopropoxy)phenylalanine methyl ester dihydrochloride (23.6 g, 90%).

*N*-(Toluene-4-sulfonyl)-L-(5,5-dimethyl)-thiaproline was coupled to L-4-(3-dimethylaminopropoxy)-phenylalanine methyl ester dihydrochloride using the procedure described in Method 13 to give *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)-L-4-(3-dimethylaminopropoxy)phenylalanine methyl ester (16.3g, 89%). The title compound was prepared via hydrolysis of the methyl ester using 0.5 N NaOH in THF/water to provide a solid, mp = >200°C (dec.).

NMR data was as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300 MHz): δ = .85 (s, 3H), .94 (s, 3H), 1.78 (m, 2H), 2.23 (s, 3H), 2.28 (s, 6H), 2.57 (m, 2H), 2.83 (m, 2H), 3.71-3.74 (m, 2H), 4.22-4.27 (m, 2H), 4.41 (d, 1H, J = 9.1 Hz), 6.58 (d, 2H, J = 8.6 Hz), 6.98 (d, 2H, J = 8.6 Hz), 7.20 (d, 2H, J = 8.3 Hz), 7.54 (d, 2H, J = 8.3 Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD, 75 MHz): δ = 22.2, 25.3, 27.8, 30.5, 39.4, 45.2, 51.9, 56.0, 57.6, 58.2, 67.2, 75.1, 115.8, 129.8, 131.6, 132.1, 132.5, 135.4, 146.7, 159.5, 170.9, 178.2.

Mass Spectroscopy: (FAB+) 586 (M+H).

Anal. HPLC: (Microsorb-MV C18 Rev. Phase, 4.6 x 150mm; Gradient 1:1 CH<sub>3</sub>CN/H<sub>2</sub>O with 0.05% TFA; flow rate = 1.0 ml/min; l = 254 nm; sample vol. = 20 mL ) Run #1: 2.988 min retention time (100.0% purity) Run #2: 3.098 min retention time (100.0% purity).

-- 131 --

## Example 21 (287)

**Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine**

5

The title compound was prepared as in Example 20 (138), except that 2-dimethylaminoethanol was used in place of 3-dimethylamino-1-propanol.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ = 7.74 (d, 2H, J = 7.4 Hz), 7.67 (d, 1H, J = 7.9 Hz), 7.42 (d, 2H, J = 8.0 Hz), 7.01 (d, 2H, J = 7.9 Hz), 6.71 (d, 2H, J = 7.8 Hz), 3.95 (m, 4H), 3.14-2.98 (4H), 2.57 (t, 2H, J = 5.6 Hz), 2.40 (s, 3H), 2.18 (s, 6H), 1.74 (m, 1H), 1.40 (m, 3H).

10

<sup>13</sup>C NMR (DMSO-d<sub>6</sub>): δ = 173.5, 169.7, 157.0, 144.1, 133.8, 131.0, 130.9, 130.3, 128.1, 113.9, 66.0, 62.4, 58.0, 55.4, 49.4, 45.9, 36.2, 30.5, 23.9, 21.4.

15

Mass Spectroscopy: FAB m/c 504 (M+H).

## Example 22 (317)

**Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[2-(*N*-ethyl-*N*-phenylamino)ethoxy]-L-phenylalanine  
Methyl Ester**

20

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 2-(*N*-ethyl-*N*-phenyl)aminoethyl chloride in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide. The title compound was prepared using the procedure described in Method 7.

25

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 1.08 (3H, t, J=8.8Hz); 1.39 (1H, m); 1.55 (3H, m); 2.19 (3H, s); 2.85-3.0 (2H, m); 3.12 (1H, dd, J=6,10.6Hz); 3.41 (2H, q, J=6,8.6Hz); 3.6 (3H, s); 3.62 (2H, t, J=5.5Hz); 4.01 (2H, t, J=5.5Hz); 4.04 (1H, t, J=8.8Hz); 4.43 (1H, dd, J=7.6,7Hz); 6.55 (1H, t, J=8.8Hz); 6.65 (2H, d, J=10Hz); 6.8 (2H, d, J=10Hz); 7.1 (4H, m); 7.38 (2H, d, J=10Hz); 7.64 (2H, d, J=10Hz); 8.1 (1H, d, J=10Hz).

35

MS: +FAB,  $m/z$  594 ( $[MH]^+$ , 45 %), 157 (100%).

## 5

**Synthesis of**  
***N*-(Toluene-4-sulfonyl)-L-prolyl-4-**  
**[2-(*N,N*-diisopropylamino)ethoxy]-L-phenylalanine**

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 2-diisopropylaminoethyl chloride in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide. The title compound was prepared using the procedure described in Method 7 as a solid, mp = 121-124°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 0.9 (12H, d, J=8Hz); 1.25-1.6 (3H, m); 1.78 (1H, m); 2.38 (3H, s); 2.7 (2H, t, J=8,8Hz); 2.9 (2H, m); 3.0 (2H, q, J=5,5,5Hz); 3.1 (1H, dd, J=3,4,3Hz); 3.35 (1H, s); 3.78 (1H, t, J=7,7Hz); 3.8 (2H, t, J=5,5Hz); 3.95 (1H, d, J=10Hz); 6.5 (1H, d, J=10Hz); 6.65 (1H, d, J=10Hz); 6.85 (1H, d, J=10Hz); 6.9 (1H, d, J=10Hz); 7.4 (2H, d, J=10Hz); 7.62 (1H, m); 7.75 (2H, d, J=10Hz).

20 MS: +FAB, m/z 560 ([MH]<sup>+</sup>, 70 %), 154 (100).

### Example 24 (333)

25 **Synthesis of**  
***N*-(Thiophene-2-sulfonyl)-L-prolyl-4-**  
**[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine**

The title compound was prepared using Method 15 and was isolated as a white, hygroscopic solid, mp = >200°C.

NMR data was as follows:

30 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ = 8.03 (br s, 1H); 7.74 (m, 2H); 7.26 (br s, 1H); 7.04 (d, 2H, J = 7.0 Hz); 6.75 (d, 2H, J = 6.0 Hz); 4.16 (m, 1H); 4.06 (m, 1H); 3.85 (m, 2H); 3.33 (m, 1 H); 3.13 - 3.00 (br m, 3H); 2.62 (m, 2H); 2.35 (s, 3H); 1.90 (m, 2H); 1.75 (m, 1H); 1.90 - 1.50 (br m, 3H).

-- 133 --

<sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 75 MHz): δ = 173.4, 169.9, 157.3, 136.1, 134.3, 133.5, 130.8, 130.3, 128.7, 128.7, 114.1, 65.7, 62.5, 55.1, 54.9, 49.7, 43.9, 36.1, 30.6, 25.9, 24.1, 21.6.

Mass Spectroscopy: (PI-FAB) 532, (M)<sup>+</sup>.

5

#### Example 25 (334)

##### Synthesis of

##### *N*-(5-Chlorothiophene-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

10

The title compound was prepared using Method 15 and was isolated as a white, hygroscopic solid, mp = >200°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ = 7.62 (m, 2H); 7.44 (m, 2H); 6.99 (m, 2H); 6.71 (m, 2H); 4.05 (m, 1H); 3.92 (m, 3H); 3.26 (m, 1H); 3.09 - 2.97 (br m, 3H); 2.30 (m, 2H); 2.13 (s, 6H); 1.82 (m, 2H); 1.79 - 1.51 (br m, 4H).

15

<sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 75 MHz): δ = 173.3, 169.3, 157.2, 137.3, 134.3, 132.5, 130.9, 130.9, 119.8, 113.9, 65.9, 62.8, 56.0, 55.5, 49.7, 45.5, 36.1, 30.6, 27.2, 24.0, 22.8.

20

Mass Spectroscopy: (PI-FAB) 588, (M + Na)<sup>+</sup>.

#### Example 26 (336)

##### Synthesis of

##### *N*-(Toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-diethylamino)ethoxy]-L-phenylalanine

25

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 2-diethylaminoethyl chloride hydrochloride in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide. The title compound was prepared using the procedure described in Method 7 as a solid, mp = 105-109°C.

30

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 1.0 (6H, t, J = 8.8Hz); 1.25-1.45 (4H, m); 1.65 (2H, m); 2.18 (3H, s); 2.26 (2H, t, J = 2.2Hz); 2.65 (4H, q, J = 5.4, 5.5Hz); 2.8-3.1 (4H, m); 3.84 (1H, dd, J = 5.4, 5.5Hz); 3.9 (2H, t, J = 3.3Hz);

35

-- 134 --

4.1 (1H, d, J = 8Hz); 6.68 (2H, d, J = 10Hz); 6.95 (2H, d, J = 10Hz); 7.05 (2H, d, J = 10Hz); 7.4 (2H, d, J = 10Hz); 7.63 (1H, d, J = 4Hz); 7.73 (2H, t, J = 4,4Hz).

MS: +ESI, m/z 532.4 ([MH]<sup>+</sup>, 100 %).

5

#### Example 27 (340)

##### Synthesis of *N*-(2,5-Dichlorothiophene-3-sulfonyl)-L-prolyl- 4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

10

The title compound was prepared using Method 15 and was isolated as a white, hygroscopic solid, mp = >200°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ = 7.77 (d, 1H, J = 6.6 Hz); 7.43 (s, 1H); 7.07 (d, 2H, J = 7.4 Hz); 6.76 (d, 2H, J = 7.4 Hz); 4.33 (m, 1H); 4.15 (m, 1H); 3.89 (t, 2H, J = 6.1 Hz); 3.29 (m, 2H); 2.97 (m, 2H); 2.57 (t, 2H, J = 7.1 Hz); 2.31 (s, 6H); 1.85 (m, 4H); 1.65 (m, 2H).

15

<sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 75 MHz): δ = 173.4, 172.5, 170.0, 157.3, 134.2, 130.8, 130.5, 130.3, 127.7, 126.9, 114.1, 65.8, 61.9, 55.3, 54.9, 49.2, 44.2, 36.2, 30.9, 26.1, 24.3, 21.6.

20

Mass Spectroscopy: (PI-FAB) 600, (M)<sup>+</sup>.

#### Example 28 (341)

##### Synthesis of *N*-(1-Methylpyrazole-4-sulfonyl)-L-prolyl-4- [3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

25

The *N*-methylpyrazole sulfonyl chloride was prepared by adding *N*-methylpyrazole to chilled (0°C) chlorosulfonic acid. The reaction mixture was allowed to warm to room temperature and then heated to 100°C overnight under a stream of N<sub>2</sub>. The reaction mixture was then cooled to room temperature and chilled to 0°C. To this solution was added thionyl chloride (2.5 eq.) and the reaction was stirred at room temperature for 30 min, then warmed to 70 °C for two hours. The reaction was cooled to room temperature and then chilled in an ice bath. Water and ice were slowly added to the

30

35

- 135 -

reaction mixture to precipitate a white solid which was collected by filtration. The desired sulfonyl chloride was washed with cold water and hexane. The title compound was prepared using Method 15 and was isolated as a white, hygroscopic solid, mp = >200°C.

5 NMR data was as follows:

<sup>1</sup>H NMR(DMSO-d<sub>6</sub>, 300 MHz): δ = 8.44 (s, 1H); 7.89 (s, 1H); 7.66 (d, 1H, J = 5.6 Hz); 7.00 (d, 2H, J = 8.5 Hz); 6.70 (d, 2H, J = 8.6 Hz); 3.89 (m, 4H), 3.87 (s, 3H); 3.14 - 2.98 (br m, 4H); 2.34 (t, 2H, J = 7.2 Hz); 2.30 (s, 6H); 1.86 (m, 3H); 1.81 - 1.75 (br m, 3H).

10 <sup>13</sup>C NMR(DMSO-d<sub>6</sub>, 75 MHz): δ = 172.3, 169.8, 157.2, 138.9, 133.7, 130.9, 130.8, 117.4, 113.9, 65.9, 62.5, 55.9, 55.3, 49.5, 36.0, 30.6, 27.2, 23.9.

Mass Spectroscopy: (PI-FAB) 530, (M)+.

#### Example 29 (346)

15

#### Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-4- [3-(*N,N*-diethylamino)propoxy]-L-phenylalanine Methyl Ester

20 The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 3-diethylaminopropyl chloride in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide.

NMR data was as follows:

25 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 1.0 (6H, bs); 1.4-1.6 (4H, m); 1.85 (2H, m); 2.18 (3H, s); 2.5 (2H, bs); 2.5-2.8 (4H, bs); 2.9 (2H, m); 3.1 (1H, dd, J = 8,10,8Hz); 3.35 (1H, dd, J = 8,4,8Hz); 3.6 (3H, s); 3.94 (2H, t, J = 6,6Hz); 4.1 (1H, m); 4.48 (1H, dd, J = 8,6,8Hz); 6.8 (2H, d, J = 10Hz); 7.1 (2H, d, J = 10Hz); 7.4 (2H, d, J = 10Hz); 7.7 (2H, d, J = 10Hz); 8.2 (1H, d, J = 10Hz).

30 MS: +ESI, m/z 560.5 ([MH]<sup>+</sup>, 100 %).



-- 136 --

## Example 30 (351)

Synthesis of  
*N*-(Thiazole-2-sulfonyl)-L-prolyl-4-  
[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

5

The sulfonyl chloride was prepared from the thiol as taught by Roblin and Clapp, *JACS*, 72, 4890, 1950. The title compound was prepared using Method 15 and was isolated as a white, hygroscopic solid, mp = >200°C.

NMR data was as follows:

10

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ = 8.24 (d, 1H, J = 3.1 Hz); 8.15 (d, 1H, 3.1 Hz); 7.63 (d, 1H, J = 5.0 Hz); 6.98 (d, 2H; J = 8.6 Hz); 6.70 (d, 2H, J = 8.6 Hz); 4.24 (m, 1H); 3.88 (m, 3H); 3.36 (m, 2H); 3.28 - 2.98 (m, 2H); 2.31 (t, 2H, J = 7.0 Hz); 2.12 (s, 6H); 1.83 - 1.79 (br m, 4H); 1.79 (m, 1H); 1.45 (m, 1H).

15

<sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 75 MHz): δ = 172.3, 168.9, 162.5, 157.2, 145.5, 131.0, 130.9, 127.5, 113.8, 65.9, 63.3, 56.1, 55.5, 50.1, 45.6, 38.8, 38.5, 35.9, 30.7, 27.3, 24.0.

Mass Spectroscopy: (PI-FAB) 555, (M-H+Na)<sup>+</sup>.

20

## Example 31 (353)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine

25

The title compound was prepared from the product of Example 33 (356) using the procedure described in Method 7 as a solid, mp = 87-90°C.

NMR data was as follows:

30

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 7.75 (d, 2H, J=10Hz); 7.65 (d, 1H, J = 4Hz); 7.4 (d, 2H, J = 10Hz); 7.18-7.3 (m, 5H); 6.96 (d, 2H, J = 10Hz); 6.65 (d, 2H, J = 10Hz); 3.97 (d, 1H, J = 4Hz); 3.91 (t, 2H, J = 4,4Hz); 3.8 (q, 1H, J = 2,2,2Hz); 3.42 (s, 2H); 2.95-3.1 (m, 4H); 2.42 (t, 2H, J = 8,8Hz); 2.38 (s, 3H); 2.08 (s, 3H); 1.85 (t, 2H, 6,6Hz); 1.7 (m, 1H); 1.3 (bs, 4H).

35

-- 137 --

## Example 32 (354)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[3-(*N,N*-diethylamino)propoxy]-L-phenylalanine

The title compound was prepared from the product of Example 29 (346) using the procedure described in Method 7 as a solid, mp = 76-82°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 1.0 (6H, t, J = 8,8Hz); 1.38 (2H, m); 1.4-1.8 (4H, m); 2.38 (3H, s); 2.45 (2H, t, J = 7,7Hz); 2.6 (4H, q, J = 6,5,6Hz); 2.95 (2H, m); 3.05 (1H, dd, J = 5,5,5Hz); 3.15 (1H, dd, J = 4,5,4Hz); 3.9 (2H, t, J = 5,5Hz); 4.0 (1H, t, J = 6,6Hz); 4.1 (1H, m); 6.7 (2H, d, J = 10Hz); 6.95 (2H, d, J = 10Hz); 7.4 (2H, d, J = 10Hz); 7.7 (1H, d, J = 4Hz); 7.75 (2H, d, J = 10Hz).

## Example 33 (356)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine  
Methyl Ester

The title compound was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 3-(*N*-benzyl-*N*-methyl)aminopropyl chloride in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide to provide a solid, mp = 60-70°C.

## Example 34 (372)

Synthesis of  
*N*-(1-Methylimidazole-4-sulfonyl)-L-prolyl-  
4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

The title compound was prepared using Method 15 and was isolated as a white, hygroscopic solid, mp = >200°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ = 7.86 (s, 1H); 7.82 (s, 1H); 7.63 (d, 1H, J = 5.6 Hz); 6.96 (d, 2H, J = 8.2Hz); 6.97 (d, 2H, J = 8.2 hz); 4.12 - 4.09

-- 138 --

(br m, 1H); 3.89 (t, 2H,  $J = 6.5$  Hz); 3.70 (s, 3H); 3.13 (m, 2H); 3.00 (m, 2H); 2.33 (t, 2H, 7.2Hz); 2.13 (s, 6H); 1.82-1.75 (m, 3H), 1.60-1.40 (br m, 3H).

$^{13}\text{C}$  NMR (DMSO- $d_6$ , 75 MHz):  $\delta = 172.8, 169.9, 157.2, 140.7, 136.1, 130.9, 126.6, 113.9, 65.9, 62.9, 56.0, 55.3, 49.7, 45.5, 35.9, 33.9, 30.6, 27.3, 24.0, 21.9.$

Mass Spectroscopy: (PI-FAB) 552, (M-H+Na) $^+$ .

#### Example 35 (373)

##### Synthesis of *N*-(2-Methylthiadiazole-5-sulfonyl)-L-prolyl- 4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

The sulfonyl chloride was prepared from the thiol as taught by Roblin and Clapp, *JACS*, 72, 4890, 1950. The title compound was prepared using Method 15 and was isolated the as a white, hygroscopic solid, mp =  $>200^\circ\text{C}$ .

NMR data was as follows:

$^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz):  $\delta = 7.66$  (d, 1H,  $J = 3.2$  Hz); 7.02 (d, 2H,  $J = 8.5$  Hz); 6.72 (d, 2H,  $J = 8.2$  Hz); 4.28 (t, 1H; 5.8 Hz); 3.96-3.89 (br m, 3H); 3.37-3.23 (br m, 2H); 3.02 (m, 1H); 2.95 (m, 1H); 2.82 (s, 3H); 2.39 (t, 2H,  $J = 7.0$  Hz); 2.13 (s, 6H); 1.83 (m, 3H); 1.80-1.40 (m, 3H).

$^{13}\text{C}$  NMR (DMSO- $d_6$ , 75 MHz):  $\delta = 173.2, 170.9, 169.0, 165.9, 157.2, 130.9, 130.8, 113.9, 65.9, 63.1, 56.0, 55.6, 50.1, 45.4, 32.2, 30.9, 27.2, 24.1, 22.3, 15.9.$

Mass Spectroscopy: (PI-FAB) 548, (M) $^+$ .

#### Example 36 (393)

##### Synthesis of *N*-(Toluene-4-sulfonyl)-L-thiaprolyl-4- [3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

The title compound was prepared as in Example 18 (138) except L-thiaproline is used in place of L-proline.

NMR data was as follows:

$^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta = 7.57$  (d, 2H,  $J = 8.40$  Hz), 7.20 (d, 2H,  $J = 8.10$  Hz), 6.92 (d, 2H,  $J = 8.40$  Hz), 6.58 (d, 2H,  $J = 8.40$  Hz), 4.50 (dd,

-- 139 --

1H, J = 4.20, 7.50 Hz), 4.45 (d, 1H, J = 10.50 Hz), 4.17 (m, 1H), 3.87 (d, 1H, J = 10.50 Hz), 3.76 (t, 2H, J = 6.00 Hz), 3.10 (m, 1H), 2.99 (m, 2H), 2.80 (m, 1H), 2.55 (m, 2H), 2.35 (m, 1H), 2.25 (s, 6H), 2.22 (s, 3H), 1.79 (m, 2H).

<sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD): δ = 177.75, 170.46, 159.43, 146.91, 135.93, 132.42, 131.73, 129.80, 115.75, 67.25, 57.90, 57.70, 52.92, 45.30, 38.33, 34.73, 27.94, 24.74, 22.16.

Mass Spectroscopy: (FAB) 546 (M+H).

#### Example 37 (472)

10

##### Synthesis of *N*-(4-Cyanobenzenesulfonyl)-L-(5,5-dimethyl)thiaprolyl- 4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine Methyl Ester

15

The title compound was prepared following the procedure outlined for the preparation of Example 20 (284).

NMR data was as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.98-7.95 (d, 2H), 7.83-7.80 (d, 2H), 7.06-7.03 (d, 2H), 6.80-6.77 (d, 2H), 4.80 (m, 1H), 4.48 (m, 1H), 3.95 (m, 3H), 3.73 (s, 3H), 3.02 (m, 2H), 2.45 (m, 2H), 2.26 (s, 6H), 1.94 (m, 2H), 1.21 (s, 3H), 1.17 (s, 3H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 179.9, 168.2, 158.8, 141.1, 133.7, 130.9, 128.1, 117.9, 115.2, 74.2, 66.7, 56.9, 55.3, 53.9, 53.0, 51.1, 46.0, 38.0, 29.9, 28.0, 24.4.

25

#### Example 38 (514)

30

##### Synthesis of *N*-(Toluene-4-sulfonyl)-L-(thiamorpholin-3-carbonyl)- 4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

35

L-Thiamorpholine-5-carboxylic acid was prepared by the method of Larsson and Carlson (*Acta Chemica Scan.* 1994, **48**, 517-525). *N*-(Toluene-4-sulfonyl)-L-thiamorpholine-5-carboxylic acid using the procedure described in Method 1. The title compound was prepared following the procedure outlined for the preparation of Example 20 (284).

-- 140 --

NMR data was as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.53-7.47 (m, 2H), 7.20-7.14 (m, 2H), 7.00-6.85 (m, 2H), 6.58-6.54 (m, 2H), 4.70-4.57 (m, 1H), 4.22-4.14 (m, 1H), 3.77-3.71 (m, 3H), 3.25-3.09 (m, 1H), 2.93-2.69 (m, 4H), 2.44 (m, 3H), 2.22 (s, 3H), 2.18 (s, 6H), 1.92 (m, 1H), 1.68 (m, 2H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ 177.9, 177.8, 169.6, 169.4, 159.6, 159.5, 146.3, 146.1, 138.9, 138.9, 132.8, 132.4, 131.8, 130.5, 129.8, 129.2, 126.9, 115.8, 115.7, 67.4, 58.1, 57.8, 57.1, 45.6, 44.9, 38.8, 37.9, 28.2, 28.2, 27.9, 26.6, 26.3, 24.7, 22.1.

Example 39 (169)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[2-(carboxy)phenoxy]-L-phenylalanine Methyl Ester

*N*-(Toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester (2.14 g, 5.16 mmol) was added to a suspension of sodium hydride (60% in oil, 1.1 eq, 228 mg) in xylenes (50 mL) at 0°C. The reaction mixture was stirred for 5 minutes and cuprous bromide-dimethyl sulfide complex (1.4 eq, 1.48 g) was added. The reaction mixture was stirred at 23°C for 0.5 hours. 2-Iodo sodium benzoate (1.5 eq, 8.06 mmol) was added and the reaction mixture was refluxed for 12 hours. EtOAc (100 mL) was added, and the organic layer washed with NH<sub>4</sub>Cl, 10% HCl, and brine, dried over MgSO<sub>4</sub>. The crude material was eluted on column chromatography (silica gel), with CHCl<sub>3</sub>:MeOH 9:1, and the title compound was isolated as an oil.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 8.16 (broad d, 1H), 7.73 (m, 2H), 7.47 (m, 2H), 7.35 (m, 2H), 7.21 (m, 2H), 7.03 (m, 2H), 6.76 (m, 1H), 4.85 (m, 1H), 4.07 (m, 1H), 3.77 (s, 3H), 3.41 (m, 1H), 3.28 (m, 1H), 3.09 (m, 2H), 2.44 (s, 3H), 2.05 (m, 1H), 1.55 (m, 3H).

Mass Spectroscopy: (FAB) 567 (M+H).

-- 141 --

## Example 40 (309)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[2-[4-(pyrimidin-2-yl)piperazin-1-yl]ethoxy]-L-phenylalanine

The methyl ester was prepared by Mitsunobu reaction of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester following the procedure described for the preparation of Example 20 (284). The title compound was prepared using the procedure described in Method 7 as a solid, mp = 102-105°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 1.35 (4H,s); 1.4 (1H,m); 1.76 (1H,m); 2.38 (3H,s); 2.5 (2H, m); 2.66 (2H, t, J = 8.8Hz); 2.9-3.1 (4H,m); 3.68 (4H, t, J = 5.5Hz); 3.8 (1H, q, J = 4.6,4Hz); 3.95 (1H, dd, J = 2,10,2Hz); 4.05 (2H, t, 6.6Hz); 6.6 (1H, t, J = 5.5Hz); 6.7 (2H, d, J = 10Hz); 6.96 (2H, d, J = 10Hz); 7.4 (2H, d, J = 10Hz); 7.65 (1H, d, J = 4Hz); 7.74 (2H, d, J = 10Hz); 8.35 (2H, d, J = 5Hz).

## Example 41 (310)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[3-(piperidin-1-yl)propoxy]-L-phenylalanine

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 1-(2-chloropropyl)piperidine in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide. The title compound was prepared using the procedure described in Method 7 as a solid, mp = 122-125°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 7.74 (d, 2H, J = 10Hz); 7.65 (d, 1H, J = 4Hz); 7.4 (d, 2H, J = 10Hz); 6.96 (d, 2H, J = 10Hz); 6.65 (d, 2H, J = 10Hz); 3.96 (dd, 1H, J = 2,6,2Hz); 3.9 (t, 3H, J = 7,7Hz); 2.95-3.1 (m, 4H); 2.46 (t, 1H, J = 2,2Hz); 2.38 (s, 3H); 2.33 (t, 2H, J = 8.8Hz); 2.24 (m, 2H); 1.78 (m, 3H); 1.45 (t, 4H, J=5.5Hz); 1.38 (m, 4H).

- 142 -

## Example 42 (311)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
 [2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine

5

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 1-(2-chloroethyl)pyrrolidine in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide. The title compound was prepared using the procedure described in Method 7 as a solid, mp = 127-130°C.

10

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 7.73 (d, 2H, J = 10Hz); 7.65 (d, 1H, J = 4Hz); 7.4 (d, 2H, J = 10Hz); 6.95 (d, 2H, J = 10Hz); 6.65 (d, 2H, J = 10Hz); 3.95 (t, 3H, J = 4,6Hz); 3.7 (q, 1H, J = 2,4,2Hz); 2.95-3.1 (m, 4H); 2.72 (t, 2H, J = 6,6Hz); 2.48 (m, 3H); 2.38 (s, 3H); 1.74 (m, 1H); 1.64 (t, 4H, J = 4,4Hz); 1.42 (m, 1H); 1.37 (s, 3H).

15

## Example 43 (316)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
 {3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine

20

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 1-(3-chlorophenyl)-4-(3-chloropropyl)piperazine in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide. The title compound was prepared using the procedure described in Method 7 as a solid, mp = 116-8°C.

25

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 7.74 (d, 2H, J = 10Hz); 7.65 (d, 1H, J = 4Hz); 7.4 (d, 2H, J = 10Hz); 7.2 (t, 1H, J = 8,8Hz); 6.96 (d, 2H, J = 10Hz); 6.9 (s, 1H); 6.85 (d, 1H, J = 10Hz); 6.75 (d, 1H, J = 10Hz); 6.7 (d, 2H, J = 10Hz); 3.95 (d, 2H, J = 8Hz); 3.92 (t, 1H, J = 8,8Hz); 3.83 (q, 1H, J = 5,4,5Hz); 3.6 (t, 1H, 5,5Hz); 3.15 (t, 4H, J = 4,4Hz); 2.95-3.1 (m, 4H); 2.5 (m, 3H); 2.45 (t, 2H, J = 8,8Hz); 2.38 (s, 3H); 1.84 (t, 2H, J = 7,7Hz); 1.75 (m, 1H); 1.4 (q, 1H, J = 8,10,8Hz); 1.35 (s, 2H).

30

35

-- 143 --

## Example 44 (318)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-  
L-phenylalanine Methyl Ester

*N*-(Toluene-4-sulfonyl)-L-prolyl-O-[(1-*tert*-butoxycarbonyl)-piperidin-3-yl)methyl]-L-tyrosine methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with *N*-Boc-3-piperidinemethyl tosylate in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide to provide a solid, mp = 60-62°C.

The title compound was prepared from this product using the procedure described in Method 7 to provide a solid, mp = 82-84°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 12.77 (br s, 1H), 8.00 (d, 1H, J = 7.9Hz); 7.68 (d, 2H, J = 8.3Hz); 7.39 (d, 2H, J = 7.9Hz); 7.13 (d, 2H, J = 8.6Hz); 6.81 (d, 2H, 8.6Hz); 4.42 (m, 1H), 4.10 (m, 1H), 3.78 (m, 3H); 3.07 (dd, 1H, J = 9.7, 4.1Hz); 2.95 (dd, 1H, J = 19Hz, 5.1Hz); 2.90 (dd, 1H, J = 19Hz); 2.8 (m, 2H); 1.80 (m, 2H); 1.58 (m, 4H); 1.34 (br s 14H).

IR (KBr, cm<sup>-1</sup>): 3400, 2900, 1745, 1700, 1525, 1450, 1350, 1250, 1160, 850, 800, 700, 650, 600.

Mass Spectroscopy: (FAB, m/e (%)) 652 (75, (M+Na<sup>+</sup>)); 530.2 (75); 224 (100).

25

## Example 45 (322)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[2-(morpholin-4-yl)ethoxy]-L-phenylalanine

30

The methyl ester was prepared by alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with *N*-(2-chloroethyl)morpholine (cesium carbonate, DMF at 60°C under Argon for 72 hrs). The product was purified by flash column chromatography (silica, EtOAc) to afford the methyl ester as an off-white foam. The title compound was prepared using the procedure described in Method 6 as a solid, mp = 99-101°C.

35



-- 144 --

NMR data was as follows:

NB contains traces Tos-Pro-Phe (NMR)

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz):  $\delta$  = 7.73 (d, 2H, J = 8.6Hz); 7.62 (d, 1H, J = 5.5Hz); 7.40 (d, 2H, J = 8.2Hz); 6.69 (d, 2H); 3.95 (m, 3H); 3.8 (m, 1H); 3.55 (m, 4H); 3.1 (m, 3H); 2.62 (t, 2H, J = 4Hz); 2.4 (m, 4H); 2.38 (s, 3H); 1.7 (m, 2H); 1.4 (m, 3H).

IR (KBr, cm<sup>-1</sup>): 3400, 2950, 2850, 1610, 1510, 1425, 1340, 1245, 1150, 1125, 825, 800, 675, 575, 525.

Mass Spectroscopy: (FAB, m/e (%)) 552 (100, (M+H<sup>+</sup>)).

#### Example 46 (325)

#### Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-4- [2-(piperidin-1-yl)ethoxy]-L-phenylalanine

The methyl ester was prepared by Mitsunobu reaction of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with piperidineethanol following the procedure described for the preparation of Example 20 (284). The title compound was prepared using the procedure described in Method 7 as a solid, mp = 102-106°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz):  $\delta$  = 7.73(d, 2H, J = 8Hz); 7.60 (d, 1H, J = 5.5Hz); 7.39 (d, 2H, J = 8Hz); 6.96 (d, 2H, J = 8.6Hz); 6.68 (d, 2H, J = 8.6Hz); 3.96 (m, 3H); 3.84 (dd, 1H J = 5.2, 4.8Hz); 2.97-3.10 (m, 4H); 2.56 (t, 2H, J = 5.9Hz); 2.38 (br s, 6H); 1.72 (m, 1H); 1.34-1.48 (m, 10H).

IR (KBr, cm<sup>-1</sup>): 3400, 2900, 1660, 1610, 1510, 1450, 1350, 1150, 875, 675, 600, 550.

Mass Spectroscopy: (FAB, m/e (%)) 542 (100, (M-H<sup>+</sup>)); 196 (10); 155 (75).

-- 145 --

## Example 47 (326)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-  
L-phenylalanine Methyl Ester

5

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-  
L-prolyl-L-tyrosine methyl ester with 1-(2-chlorophenyl)-4-(3-chloro-  
propyl)piperazine in refluxing 2-butanone in the presence of potassium  
10 carbonate and sodium iodide.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 8.2 (d, 1H, J = 10Hz); 7.68 (d, 2H, J = 10Hz); 7.4 (d, 2H, J = 10Hz); 7.19 (t, 1H, J = 8,8Hz); 7.13 (d, 2H, J = 10Hz); 6.9 (s, 1H); 6.85 (d, 1H, J = 10Hz); 6.82 (d, 2H, J = 10Hz); 6.75 (d, 1H, J = 10Hz); 4.45 (q, 1H, J = 8,5,8Hz); 4.08 (t, 1H, J = 4,4Hz); 3.94 (t, 2H, J = 5,5Hz); 3.3 (s, 3H); 3.12 (bs, 4H); 3.1 (t, 1H, J = 8,8Hz); 2.96 (m, 2H); 2.47 (m, 3H); 2.38 (s, 3H); 1.85 (t, 2H, J = 6,6Hz); 1.57 (m, 3H); 1.4 (m, 1H).

15

MS: EI, m/z 682/684 ([MH]<sup>+</sup>, 18 %), 209 (26%).

20

## Example 48 (327)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[2-(azepan-1-yl)ethoxy]-L-phenylalanine

25

The title compound was prepared from the product of Example 49 (328) using the procedure described in Method 7 as a solid, mp: 105-107°C.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 7.75 (t, 2H, J = 8,8Hz); 7.45 (d, 1H, J = 8Hz); 7.4 (t, 2H, J = 8,8Hz); 7.05 (d, 1H, J = 10Hz); 6.96 (d, 1H, J = 10Hz); 6.7 (d, 1H, J = 10Hz); 4.1 (t, 1H, J = 5,5Hz); 3.92 (t, 3H, J = 8,8Hz); 3.82 (m, 1H); 3.6 (t, 1H); 2.8 (t, 2H, J = 6,6Hz); 2.66 (d, 4H, J = 5Hz); 2.5 (d, 3H, J = 10Hz); 1.76 (t, 1H, 7,7Hz); 1.65 (m, 1H); 1.5 (bs, 6H); 1.43 (t, 2H, J = 8,8Hz); 1.36 (m, 2H); 1.28 (s, 1H); 1.15 (s, 1H).

30

35

-- 146 --

## Example 49 (328)

**Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[2-(azepan-1-yl)ethoxy]-L-phenylalanine Methyl Ester**

The title compound was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 2-(hexamethylene-imino)ethyl chloride in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide to provide a solid, mp = 60-65°C.

## Example 50 (347)

**Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[3-(4-methylpiperazin-1-yl)propoxy]-  
L-phenylalanine Methyl Ester**

The methyl ester was prepared via *O*-alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 3-(*N*-methylpiperazine)propyl chloride in refluxing 2-butanone in the presence of potassium carbonate and sodium iodide.

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 8.2 (t, 1H, J = 10,10Hz); 7.7 (t, 2H, J = 10,12Hz); 7.4 (d, 2H, J = 10Hz); 7.1 (t, 2H, J = 10,10Hz); 6.8 (d, 2H, J = 10Hz); 4.44 (q, 1H, J = 4,6,6Hz); 4.1 (dd, 1H, J = 14,8,8Hz); 3.93 (t, 2H, J = 6,6Hz); 3.6 (s, 3H); 3.08 (q, 1H, J = 6,4,4Hz); 3.0 (dd, 1H, J = 12,4,4Hz); 2.9 (m, 2H); 2.38 (s, 3H); 2.2-2.35 (m, 10H); 2.12 (s, 3H); 1.8 (t, 2H, J = 6,6Hz); 1.55 (m, 3H); 1.41 (m, 1H).

MS: +ESI, m/z 587 ([MH]<sup>+</sup>, 100 %).

## Example 51 (355)

**Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine**

The title compound was prepared from the product of Example 50 (347) using the procedure described in Method 7 as a solid, mp = 80-83°C.

-- 147 --

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 Mhz): δ = 7.75 (d, 2H, J = 10Hz); 7.63 (d, 1H, J = 4Hz); 7.4 (d, 2H, J = 10Hz); 7.05 (d, 2H, J = 10Hz); 6.95 (d, 2H, J = 10Hz); 6.67 (d, 2H, J = 10Hz); 4.1 (d, 1H, J = 8Hz); 3.94 (d, 1H, J = 8Hz); 3.9 (t, 2H, J = 5.5Hz); 3.8 (bs, 1H); 3.08 (m, 1H); 2.91 (d, 1H, J = 10Hz); 2.85 (dd, 2H, J = 6,18,6Hz); 2.38 (s, 3H); 2.15-2.35 (m, 8H); 2.14 (s, 3H); 1.78 (q, 2H, J = 6,8,6Hz); 1.7 (m, 1H); 1.4 (m, 1H); 1.37 (m, 2H).

Example 52 (345)

10

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-  
4-*N*-(trifluoromethanesulfonyl)amino-  
L-phenylalanine Methyl Ester

15

The title compound was produced by reaction of *N*-(toluene-4-sulfonyl)-L-prolyl-(4-amino)phenylalanine methyl ester with trifluoromethanesulfonic acid anhydride in pyridine to provide a solid, mp = 75-78°C.

Example 53 (370)

20

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-  
4-*N*-(trifluoromethanesulfonyl)amino-L-phenylalanine

25

The title compound was prepared from the product of Example 52 (345) using the procedure described in Method 6.

NMR data was as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400MHz): δ = 8.08 (s, 1H); 7.7 (d, 2H); 7.56 (d, 1H); 7.34 (d, 2H); 7.22 (s, 2H); 4.85 (m, 1H); 4.16 (m, 1H); 3.40 (m, 3H); 3.09 (m, 2H); 2.44 (s, 3H); 1.86 (m, 1H); 1.50 (m, 3H).

30

IR (KBr, cm<sup>-1</sup>): 3390; 2950; 1750; 1650; 1525; 1425; 1375; 1340; 1200; 1150; 950; 825; 625; 590; 550.

Mass Spectroscopy: (+ESI) 564 ( [M+H]<sup>+</sup> ); 530; 462; 406; 362; 342; 335; 157.

35

-- 148 --

## Example 54 (387)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(*N*-benzylaminocarbony)methoxy]-L-phenylalanine  
Methyl Ester

5

The title compound was prepared by alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with *N*-benzyl-2-chloroacetamide (potassium carbonate, sodium iodide, refluxing butanone under Argon overnight). The product was purified by flash column chromatography (silica, 1:1 hexane:EtOAc) to afford the methyl ester as a white solid, mp = 57-59°C.

10

## Example 55 (389)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(benzyloxycarbony)methoxy]-L-phenylalanine

15

The title compound was prepared from the product of Example 4 (387) using the procedure described in Method 6 as a solid, mp = 79-81°C.

20

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 12.2 (br s, 1H); 8.60 (t, 1H); 8.03 (d, 1H, J = 7.9Hz); 7.70 (d, 2H, J = 6.6Hz); 7.39 (d, 2H, J = 8.4Hz); 7.15 (d, 2H, J = 8.6Hz); 6.87 (d, 2H, J = 8.6Hz); 4.48 (s, 2H); 4.42 (m, 1H); 4.31 (d, 2H, J = 6.3Hz); 4.10 (m, 1H); 3.0-3.2 (m, 2H); 2.8-2.9 (m, 2H); 2.38 (s, 3H); 1.2-1.6 (m, 4H).

25

IR (KBr, cm<sup>-1</sup>): 3400, 2950, 1725, 1660, 1525, 1510, 1450, 1350, 1240, 1150, 1080, 670, 575, 550.

Mass Spectroscopy: (FAB, m/e (%)) 602 (10, (M+Na<sup>+</sup>)); 580 (10, (M+H<sup>+</sup>)); 131 (100).

30

## Example 56 (390)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-  
4-[(carboxy)methoxy]-L-phenylalanine

35

Alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with t-butyl bromoacetate (potassium carbonate, DMF, under Argon for 72 hrs)

-- 149 --

gave *N*-(toluene-4-sulfonyl)-L-prolyl-L-*O*-(*tert*-butoxycarbonylmethyl)-tyrosine methyl ester after purification by flash column chromatography (silica, 1:1 hexane:EtOAc) to afford a white solid, mp = 55°C.

5        *N*-(Toluene-4-sulfonyl)-L-prolyl-L-*O*-(*tert*-butoxycarbonylmethyl)-tyrosine was prepared from the *N*-(toluene-4-sulfonyl)-L-prolyl-L-*O*-(*tert*-butoxycarbonylmethyl)-tyrosine methyl ester using the procedure described in Method 6 as a solid, mp = 69-70°C.

10        The title compound was prepared from *N*-(Toluene-4-sulfonyl)-L-prolyl-L-*O*-(*tert*-butoxycarbonylmethyl)-tyrosine by reaction with formic acid. Removal of the formic acid and titration with ether afforded the desired compound as a white solid, mp = 70-73°C.

NMR data was as follows:

15        <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 12.85 (br s, 2H); 8.01 (d, 1H, J = 7.9Hz); 7.69 (d, 2H, J = 8.3Hz); 7.39 (d, 2H, J = 8.3Hz); 7.13 (d, 2H, J = 8.8Hz); 6.79 (d, 2H, J = 8.6Hz); 4.6 (s, 2H); 4.42 (m, 1H); 4.10 (m, 1H); 3.08 (m, 1H); 2.95 (m, 2H); 2.38 (s, 3H); 1.5 (m, 4H).

IR (KBr, cm<sup>-1</sup>): 3350, 2950, 1730, 1625, 1510, 1425, 1340, 1175, 1160, 1075, 825, 675, 575, 550.

20        Mass Spectroscopy: (FAB, m/e (%)) 513 (100, (M+Na<sup>+</sup>)), 491 (75, (M+H<sup>+</sup>)).

#### Example 57 (407)

#### 25        Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-4- [(aminocarbonyl)-L-phenylalanine Methyl Ester

30        The title compound was prepared by alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 2-chloroacetamide (potassium carbonate, sodium iodide, refluxing butanone under Argon for 48 hr). The product was purified by flash column chromatography (silica, EtOAc, then 5% MeOH in EtOAc) to afford the title compound as a white solid, mp = 60-64°C.

-- 150 --

## Example 58 (408)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(aminocarbonyl)methoxy]-L-phenylalanine

5 The title compound was prepared from the product of Example 57 (407) using the procedure described in Method 6 as a solid, mp = 195-196°C.

NMR data was as follows:

10 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 12.2 (br s, 1H); 8.02 (d, 1H, J = 8.1Hz); 7.69 (d, 2H, J = 8.3Hz); 7.47, (br s, 1H); 7.40 (d, 2H, J = 7.9Hz); 7.35 (br s, 1H); 7.14 (d, 2H, J = 8.6Hz); 6.84 (d, 2H, J = 8.6Hz); 4.40 (m, 1H); 4.35 (s, 2H); 4.11 (dd, 1H); 3.09 (m, 1H); 2.91 (dd, 1H); 2.39 (s, 3H); 1.45-1.55 (m, 3H); 1.40 (m, 1H).

15 IR (KBr, cm<sup>-1</sup>): 3500, 3350, 3250, 2950, 1725, 1675, 1660, 1560, 1510, 1450, 1400, 1350, 1225, 1200, 1150, 1050, 825, 660, 575, 550.

Mass Spectroscopy: (FAB, m/e (%)) 488 (100, (M-H<sup>+</sup>)).

## Example 59 (409)

Synthesis of  
*N*-(Toluene-4-sulfonyl)-L-prolyl-4-  
[(*N*-tert-butylaminocarbonyl)methoxy]-L-phenylalanine

20 The methyl ester was prepared by alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-tyrosine methyl ester with 2-chloro-*N*-tert-butylacetamide (potassium carbonate, sodium iodide, reflux in butanone under Argon overnight). The  
25 product was purified by flash column chromatography (silica, 1:1 hexane:EtOAc) to afford the methyl ester as a white solid. The title compound was prepared using the procedure described in Method 6 as a solid, mp = 88-89°C.

30 NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 12.2 (br s, 1H); 8.02 (d, 1H, J = 8Hz); 7.68 (d, 2H, J = 8.3Hz); 7.39 (d, 3H, J = 8Hz); 7.14 (d, 2H, J = 8.8Hz); 6.82 (dd, 2H, J = 8.4, 2Hz); 4.4 (m, 1H); 4.32 (s, 2H); 4.10 (dd, 1H, J = 2.9, 8Hz); 3.07 (m, 1H); 3.0 (dd, 1H, J = 18.7, 27.5Hz); 2.94 (dd, 1H, J = 17.8, 26Hz); 2.39 (s, 3H); 1.5 (m, 3H); 1.4 (m, 1H); 1.29 (s, 9H).

35

-- 151 --

IR (KBr,  $\text{cm}^{-1}$ ): 3400, 2950, 1745, 1675, 1525, 1450, 1350, 1225, 1160, 1075, 825, 675, 575, 540.

Mass Spectroscopy: (FAB,  $m/e$  (%)) 544 (100, (M-H<sup>+</sup>)).

5

#### Example 60 (410)

#### Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-4-[2-(4-phenyl-4-hydroxypiperidin-1-yl)ethoxy]-L-phenylalanine Methyl Ester

10

The methyl ester was prepared by alkylation of *N*-(toluene-4-sulfonyl)-L-prolyl-L-*O*-(2-chloroethyl)tyrosine methyl ester with 4-hydroxy-4-phenyl piperidine (potassium carbonate, sodium iodide, in refluxing butanone under Argon for 72 hr). The product was purified by flash column chromatography (silica, 5% methanol in chloroform) to afford the methyl ester as a white foam. The title compound was prepared using the procedure described in Method 6 as a solid, mp = 122-123°C.

15

NMR data was as follows:

<sup>1</sup>H NMR (DMSO- $d_6$ , 400 MHz):  $\delta$  = 7.72 (m, 2H); 7.61 (d, 1H, J = 5.5Hz); 7.39 (d, 2H, J = 7.2Hz); 7.28 (m, 2H); 7.17 (m, 1H); 7.04 (d, 1H, J = 8.8Hz); 6.96 (d, 1H, J = 8.6Hz); 6.71 (dd, 2H, J = 2.4, 8.8Hz); 4.75 (s, 1H); 4.1 (m, 1H); 4.0 (m, 2H); 3.9 (q, 1H); 3.8 (q, 1H); 2.8-3.1 (m, 4H, overlapping signals); 2.7 (m, 4H, overlapping signals); 2.38 (s, 3H); 1.65 (m, 2H); 1.55 (m, 2H); 1.4 (m, 2H).

20

IR (KBr,  $\text{cm}^{-1}$ ): 3375, 2890, 1660, 1610, 1510, 1390, 1325, 1250, 1160, 1075, 1040, 700, 675, 575, 530.

25

Mass Spectroscopy: (FAB,  $m/e$  (%)) 648 (100, (M+2Li-H)<sup>+</sup>), 642 (90 (M+Li)<sup>+</sup>).

30

#### Example 61 (375)

#### Synthesis of *N*-(Toluene-4-sulfonyl)sarcosyl-D,L-4-(amidino)phenylalanine



-- 152 --

*N*-(Toluene-4-sulfonyl)sarcosyl-D,L-4-cyanophenylalanine methyl ester (see Example 61 (381)) (167 mg, 0.388 mmol) was dissolved in pyridine (6 mL) and then H<sub>2</sub>S gas was bubbled in until saturated. The mixture was stirred for 19 hours then the volatiles were removed under a stream of N<sub>2</sub>. The residue was taken up into EtOAc (50 mL) and washed with 5% aq KHSO<sub>4</sub> (2 X 25 mL). The organic solution was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated in vacuo to give *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-thiocarboxamidophenylalanine methyl ester. The thioamide was dissolved in acetone (10 mL). Iodomethane (1 mL) was added and the mixture was heated to reflux for 1 hr. The volatiles were removed in vacuo to give *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-methylthioimidatephenylalanine methyl ester hydroiodide (256 mg, 100%). The thioimidate was dissolved in MeOH (5 mL). Ammonium acetate (52 mg, 0.67 mmol) was added and the mixture was heated to reflux for 1.5 hr. The solvent was removed in vacuo and the residue was purified by preparative TLC (90:10:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH) to give *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-amidinophenylalanine methyl ester (75 mg, 38%). The title compound was prepared via hydrolysis of the methyl ester using 0.5 N NaOH in THF/water (66 mg, 87%).

NMR data was as follows:

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ = 7.66 (m, 4H), 7.43 (d, 2H, J = 7.7 Hz), 7.29 (d, 2H, J = 8.0 Hz), 4.10 (m, 1H), 3.57 (s, 2H), 3.20-3.06 (m, 2H), 2.54 (s, 3H), 2.40 (s, 3H).

Mass Spectroscopy: FAB m/e 433 (M+H).

25

Example 62 (381)

Synthesis of  
*N*-(Toluene-4-sulfonyl)sarcosyl-  
D,L-4-(aminocarbonyl)phenylalanine

30

*N*-(Toluene-4-sulfonyl)sarcosine was coupled to 4-cyanophenylalanine methyl ester hydrochloride (prepared via the method of Wagner, Voight, and Vieweg *Pharmazie* 1984, 39, 226-230) to give *N*-(toluene-4-sulfonyl)sarcosyl-

-- 153 --

D,L-4-cyanophenylalanine methyl ester. The compound was prepared via hydrolysis of the methyl ester using 0.5 N NaOH in THF/water.

*N*-(Toluene-4-sulfonyl)sarcosyl-D,L-4-cyanophenylalanine methyl ester  
5 (300 mg, 0.699 mmol) was slurried in EtOH (3mL). NaOH (10N, 98  $\mu$ L) and  $H_2O_2$  (475  $\mu$ L, 5.51 mmol) were added. The mixture was heated to 50 °C for 16 hr whereupon a white precipitate deposited. The mixture was cooled to room temp and made acidic with HCl (6N). The mixture was diluted with water (20 mL) and extracted with chloroform (4 X 25 mL). The organic  
10 extracts were dried ( $Na_2SO_4$ ), filtered and recrystallized from methanol to give the compound as a white solid (135 mg, 45%).


NMR data was as follows:

$^1H$  NMR (DMSO- $d_6$ ):  $\delta$  = 8.31 (br d, 1H,  $J$  = 3.6 Hz), 7.92 (br s, 1H), 7.72  
(d, 2H,  $J$  = 7.8 Hz), 7.62 (d, 2H,  $J$  = 7.9 Hz), 7.40-7.21 (5H), 4.47 (m, 1H),  
15 3.59 (m, 2H), 3.15 (m, 1H), 2.94 (m, 1H), 2.53 (s, 3H), 2.39 (s, 3H).

$^{13}C$  NMR (DMSO- $d_6$ ):  $\delta$  = 172.9, 168.0, 167.3, 143.7, 141.3, 134.2,  
132.8, 130.1, 129.4, 127.8, 127.7, 53.4, 52.4, 36.7, 36.0, 21.3.

Mass Spectroscopy: FAB  $m/e$  434 ( $M+H$ ).

20 Other compounds prepared by the methods described above include those set forth in Table II below as Examples 63-135:



Q = -C(O)NH- in all cases

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Example No.
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[NH(C(O)CH <sub>2</sub> (NHAc)(CH <sub>3</sub> ),NHCH <sub>2</sub> )- benzyl]-	-OH	63
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[NH(C(O)CH <sub>2</sub> (CH <sub>2</sub> (NHCH <sub>2</sub> )-benzyl)- C(O)OCH <sub>3</sub> φ	-OCH <sub>3</sub>	64
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -[NH(C(O)CH <sub>2</sub> (NHAc)-benzyl)- CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	-OH	65
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -(1,4-NHCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> (O)NH)-benzyl-	-OH	66
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -(1,4-NHCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> (O)NH)-benzyl-	-OH	67

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Example No.
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(CH <sub>2</sub> NHCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -C(O)NH)-benzyl-	-OH	68
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(CH <sub>2</sub> (Iboc)NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> - C(O)NH)-benzyl-	-OH	69
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(φCH <sub>2</sub> OCCH <sub>2</sub> (H,N)CHC(O)NH)-benzyl-	-OH	70
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(HO(O)C(Cbz-NH)CHCH <sub>2</sub> CH <sub>2</sub> - C(O)NH)-benzyl-	-OH	71
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(HO(O)C(H,N)CHCH <sub>2</sub> CH <sub>2</sub> - C(O)NH)-benzyl-	-OH	72
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(CH <sub>2</sub> (N-Boc)NCH <sub>2</sub> -C(O)NH)-benzyl-	-OH	73
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic (H <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - 1,5,5-dimethylthiazolidin- 4-yl)	<i>p</i> -(CH <sub>2</sub> (N-Iboc)NCH <sub>2</sub> -C(O)NH)-benzyl-	-OH	74
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(CH <sub>2</sub> NHCH <sub>2</sub> -C(O)NH)-benzyl-	-OCCH <sub>2</sub> CH <sub>3</sub>	75

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Example No.
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(CH <sub>2</sub> ) <sub>2</sub> NHCH <sub>2</sub> C(O)NH- <i>l</i> -benzyl-	-OH	76
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)	<i>p</i> -(CH <sub>2</sub> ) <sub>2</sub> NHCH <sub>2</sub> C(O)NH- <i>l</i> -benzyl-	-OH	77
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> C(O)NH- <i>l</i> -benzyl-	-OH	78
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> - (L-5,5- dimethylthiazolidin- 4-yl)	<i>p</i> -(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> C(O)NH- <i>l</i> -benzyl-	-OH	79
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -( <i>tert</i> -butyl-O)(O)CCH <sub>2</sub> -O-benzyl-	-OCH <sub>3</sub>	80
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(2-formyl-1,2,3,4-tetrahydroisoquinolin-3- yl)-CH <sub>2</sub> -NH- <i>l</i> -benzyl-	-OH	81
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)	<i>p</i> -(OCH <sub>2</sub> CH <sub>2</sub> (NH <i>Boc</i> )CH <sub>2</sub> cyclohexyl)- benzyl	-OCH <sub>3</sub>	82

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	Example No.
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>m</i> -[O(CH <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> ]benzyl-	-OH	83
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> -		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O]benzyl-	-OH	84
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic -CH <sub>2</sub> -S-(CH <sub>2</sub> ) <sub>2</sub> - (1-5,5- dimethylthiazolidin- 4-yl)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O]benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>	85
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[2-(2-azabicyclo[3.2.2]octan- 2-yl)ethyl O]benzyl-	-OCH <sub>3</sub>	86
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O]benzyl-	-OH	87
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[(CH <sub>2</sub> ) <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O]benzyl-	-OCH <sub>3</sub>	88
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[2-(2-azabicyclo[3.2.2]octan- 2-yl)ethyl O]benzyl-	-OH	89
<i>p</i> -CH <sub>3</sub> -φ	H	-CH <sub>2</sub> -φ (L isomer)	<i>p</i> -(cyclopentyl-C≡C-)benzyl-	-OH	90

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	Example No.
<i>p</i> -CH <sub>3</sub> , $\phi$ -	II	-CH <sub>2</sub> , $\phi$ (I, isomer)	<i>p</i> -[-C $\equiv$ C( $\phi$ - <i>p</i> - $\phi$ )-benzyl]-	-OH		91
<i>p</i> -CH <sub>3</sub> , $\phi$ -	II	-CH <sub>2</sub> , $\phi$ (I, isomer)	<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O-S(O) <sub>2</sub> - <i>p</i> -CH <sub>3</sub> , $\phi$ )-benzyl]-	-OH		92
<i>p</i> -CH <sub>3</sub> , $\phi$ -	II	-CH <sub>2</sub> , $\phi$ (I, isomer)	<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -NH(C(O)NH <sub>2</sub> )-benzyl)-	-OH		93
<i>p</i> -CH <sub>3</sub> , $\phi$ -	II	-CH <sub>2</sub> , $\phi$ (I, isomer)	<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O- <i>p</i> -(X)(X)(CH <sub>3</sub> , $\phi$ )-benzyl)-	-OH		94
<i>p</i> -CH <sub>3</sub> , $\phi$ -	II	-CH <sub>2</sub> , $\phi$ (I, isomer)	<i>p</i> -[-C $\equiv$ C(-CH(NH <sub>2</sub> )-cyclohexyl)-benzyl]-	-OH		95
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O-phenyl)-benzyl]-	-OH		96
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	II	<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O-phenyl)-benzyl]-	-OH		97
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O(CH <sub>3</sub> )-benzyl)-	-OH		98
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	II	<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O(CH <sub>3</sub> )-benzyl)-	-OH		99
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)		<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O- <i>p</i> -(C(O)OC <sub>2</sub> H <sub>5</sub> )-phenyl)-	-OH		100
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	II	<i>p</i> -[-C $\equiv$ C(-CH <sub>2</sub> -O- <i>p</i> -(C(O)OC <sub>2</sub> H <sub>5</sub> )-phenyl)-	-OH		101

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	Example No.
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> CH(C(=O)OCH <sub>3</sub> )]-benzyl-	-OH	102
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	H	<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> CH(C(=O)OCH <sub>3</sub> )]-benzyl-	-OH	103
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> CH(C(=O)OH)-benzyl-   NH(C(=O)CH <sub>3</sub> )]	-OH	104
<i>p</i> -H, $\phi$ -	-CH <sub>3</sub>	H	<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> CH(C(=O)OH)-benzyl-   NH(C(=O)CH <sub>3</sub> )]	-OH	105
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> NH-(4,5-dihydro-4-oxo- 5-phenyl-oxazol-2-yl)]benzyl	-OH	106
<i>p</i> -CH <sub>3</sub> , $\phi$ -	-CH <sub>3</sub>	H	<i>p</i> -[C $\equiv$ C-CH <sub>2</sub> NH-(4,5-dihydro-4-oxo- 5-phenyl-oxazol-2-yl)]benzyl	-OH	107
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -(1-piperidinyl)]-benzyl-	-OCH <sub>3</sub>	108
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>m</i> -[OCH <sub>2</sub> CH <sub>2</sub> -(1-pyrrolidinyl)]-benzyl-	-OH	109
<i>p</i> -CH <sub>3</sub> , $\phi$ -	R <sup>2</sup> /R <sup>3</sup> = cyclic 3 carbon atoms (L-pyrrolidinyl)		<i>p</i> -[OCH <sub>2</sub> CH <sub>2</sub> -(N-morpholino)]-benzyl-	-OH	110



R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	Example No
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)		<i>p</i> -[1-OC(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -1-(4-hydroxy-4-(3-methoxypyrrrol-2-yl)-piperazinyl)-benzyl-	-OCH <sub>3</sub>	111
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)		<i>p</i> -[1-(3-(N-Boc)-piperidinyl)-benzyl-	-OH	112
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)		<i>m</i> -[1-(N-methyl-piperidin-4-yl)-benzyl-	-OH	113
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)		<i>p</i> -[1-O-(N-methyl-piperidin-4-yl)-benzyl-	-OH	114
<i>p</i> -C(CH <sub>3</sub> ) <sub>2</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic CH <sub>2</sub> -S-C(CH <sub>3</sub> ) <sub>2</sub> -		<i>p</i> -[1-(1-methylpiperidin-4-yl)-O-(benzyl-	-OC(CH <sub>3</sub> ) <sub>2</sub> CH <sub>3</sub>	115
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic CH <sub>2</sub> CH <sub>2</sub> SO <sub>2</sub> CH <sub>2</sub> - (1,1,1-dioxathiomorpholin-3-yl)		<i>p</i> -[1-(1-methylpiperidin-4-yl)-O-(benzyl-	-OC(CH <sub>3</sub> ) <sub>2</sub> CH <sub>3</sub>	116
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic CH <sub>2</sub> CH <sub>2</sub> SO <sub>2</sub> CH <sub>2</sub> - (1,1,1-dioxathiomorpholin-3-yl)		<i>p</i> -[1-(1-methylpiperidin-4-yl)-O-(benzyl-	-OC(CH <sub>3</sub> ) <sub>2</sub> CH <sub>3</sub>	117

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Example No.
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic C(CH <sub>3</sub> ) <sub>2</sub> SO <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> (1,1,1,1-tetrahydro- dioxathiomorpholin- 3-yl)	<i>p</i> -(1-methylpiperidin-4-yl)-O- <i>l</i> -benzyl-	-OH	118
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic C(CH <sub>3</sub> ) <sub>2</sub> S-C(CH <sub>3</sub> ) <sub>2</sub> - (1,5,5-dimethylthiazolidin- 4-yl)	<i>p</i> -(1-methylpiperidin-4-yl)-O- <i>l</i> -benzyl-	-OH	119
φ-CH <sub>2</sub> -	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)	<i>p</i> -(1-methylpiperidin-4-yl)-O- <i>l</i> -benzyl-	-OCH <sub>2</sub> CH <sub>3</sub>	120
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)	<i>p</i> -1-NHSO <sub>2</sub> -C(CH <sub>3</sub> ) <sub>2</sub> -benzyl-	-OCH <sub>3</sub>	121
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)	<i>p</i> -1-NHSO <sub>2</sub> -C(CH <sub>3</sub> ) <sub>2</sub> -benzyl-	-OCH <sub>3</sub>	122
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)	<i>p</i> -[ <i>N</i> ,3-methylbutyl- <i>N</i> , trifluoromethanesulfonyl]amino]benzyl-	-OCH <sub>3</sub>	123
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidinyl)	<i>p</i> -[ <i>N</i> ,vinylsulfonyl]amino]benzyl-	-OH	124

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Example No.
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -(OC(=O)-O-benzyl)-	-OCH <sub>3</sub>	125
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -(piperidin-1-yl)C(O)CH <sub>2</sub> -O-benzyl-	-OH	126
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -(C(=O),C(=O),N(C(=O)CH <sub>2</sub> -O-benzyl)-	-OCH <sub>3</sub>	127
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -(CH <sub>3</sub> ) <sub>2</sub> C(=O),NC(O)CH <sub>2</sub> -O-benzyl-	-OH	128
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -( <i>N</i> -methylacetamido)benzyl-	-OCH(CH <sub>3</sub> ) <sub>2</sub>	129
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -( <i>N</i> -methylacetamido)benzyl-	-OH	130
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	<i>p</i> -( <i>N</i> -methyltrifluoroacetamido)benzyl-	-OCH <sub>3</sub>	131
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (1-pyrrolidiny)	(1-toluenesulfonylimidazol-4-yl)methyl-	-OCH <sub>3</sub>	132

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Example No.
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	1-((N,N-dimethylaminosulfonyl)-imidazol-4-yl)methyl-	-OCH <sub>3</sub>	133
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -(N-toluenesulfonylamino)benzyl-	-OCH <sub>3</sub>	134
<i>p</i> -CH <sub>3</sub> -φ	R <sup>2</sup> /R <sup>1</sup> = cyclic 3 carbon atoms (L-pyrrolidiny)	<i>p</i> -(N-toluenesulfonylamino)benzyl-	-OH	135

-- 164 --

Further to the above, the following additional compounds were prepared as Examples 136-140:

5 Example 136

**Synthesis of *N*-(Toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine *t*-butyl ester**

10 The title compound was prepared as in Example 20 except *N*-Boc-tyrosine *t*-butyl ester was used instead of *N*-Boc-tyrosine methyl ester.

MS: [(+)ESI], [M+H]<sup>+</sup> 620.

15 Example 137

**Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-L-(4-(*N*-methylpiperidinoxy)phenylalanine) *tert* butyl ester**

20 The title compound was prepared by BOP coupling of Tos-Pro-OH with Tyr-O-1-methyl piperidine *t*-butyl ester (prepared by Mitsunobu reaction). The crude product was purified by flash chromatography (silica, 95:5 EtOAc:Et<sub>3</sub>N) to afford a white solid (0.615 g, 60%).

MS ((+) ESI, *m/z* (%)) 586 (100 [M+H]<sup>+</sup>).

25 Anal. Calc'd for C<sub>31</sub>H<sub>43</sub>N<sub>3</sub>O<sub>6</sub>S: C, 63.57; H, 7.40; N, 7.17. Found: C, 63.11; H, 7.37; N, 6.96.

Example 138

30 **Synthesis of *N*-(Toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-L-(4-(*N*-methyl(piperidinoxy) phenylalanine *t*-butyl ester**

The title compound was prepared following the procedure described in Example 137 with substitution for the appropriate starting materials.

35

-- 165 --

Anal. Calc'd for  $C_{37}H_{45}N_3O_6S_2 \cdot 0.25 CH_2Cl_2$ ; C, 58.85; H, 7.02; N, 6.43.  
Found: C, 58.75; H, 6.92; N, 6.48.

MS (+ESI): 632 [M+H]<sup>+</sup>

5

## Example 139

10 **Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine  
tert-butyl ester**

The title compound was prepared from the corresponding triflate (which was prepared from (*N*-(Toluene-4-sulfonyl)-L-prolyl-L-tyrosine tert-butyl ester as taught by Tilley and coworkers, *J. Org. Chem.*, 55, 906, 1990). The dipeptide (505 mg, 0.8 mmol.), a catalytic amount of tetrakis(triphenyl-  
15 hosphine)palladium(0), potassium carbonate (201 mg, 1.5 eq.), phenylboronic acid (199 mg, 2.0 eq.), and 15 mL of toluene were refluxed for 10 hours with stirring. Ethyl acetate was added and the organic layer was washed with water, 1N NaOH, brine and dried over magnesium sulfate. Upon filtration, the solvent was evaporated under reduced pressure. The crude material was  
20 purified on a preparative plate (1:1 ethyl acetate:hexanes). The silica gel was washed several times with acetonitrile and ethyl acetate. The combined fractions were evaporated and the residue was dried under reduced pressure.

NMR data was as follows:

25 <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ = 7.70 (m, 1H); 7.57 (m, 3.5H); 7.45 (m, 3.5H); 7.28 (m, 5H); 4.78 (m, 1H); 4.06 (m, 1H); 3.30 (m, 2H); 3.06 (m, 2H); 2.40 (s, 3H); 2.05 (m, 1H); 1.42 (s, 9H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 171.02, 169.98, 144.4, 140.78, 139.82, 135.53, 132.82, 129.97, 129.9, 129.41, 128.78, 127.86, 127.04, 126.98, 82.65, 62.15,  
30 53.74, 49.49, 37.43, 29.67, 27.78, 23.92, 21.37.

-- 166 --

## Example 140

**Synthesis of *N*-(Toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine**

5       The title compound was prepared from the product of Example 139 using the procedure described in Method 11.

NMR data was as follows:

10       <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300 MHz): δ = 8.05 (m, 1H); 7.71 (d, 2H, J = 8.24 Hz); 7.55 (m, 4H); 7.30 (m, 8H); 4.71 (m, 1H); 4.09 (m, 1H); 3.30 (m, 3.30); 3.15 (m, 3H); 2.37 (s, 3H); 1.78 (m, 1H); 1.62 (m, 4H).

15       <sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 174.27, 145.88, 142.25, 141.23, 137.44, 135.12, 131.19, 131.15, 129.98, 129.09, 128.4, 128.17, 128.0, 63.25, 54.69, 50.52, 37.85, 31.52, 25.21, 21.43.

## Example 141

20       ***In vitro* Assay For Determining Binding of  
Candidate Compounds to VLA-4**

25       An *in vitro* assay was used to assess binding of candidate compounds to α<sub>4</sub>β<sub>1</sub> integrin. Compounds which bind in this assay can be used to assess VCAM-1 levels in biological samples by conventional assays (e.g., competitive assays). This assay is sensitive to IC<sub>50</sub> values as low as about 1nM.

30       The activity of α<sub>4</sub>β<sub>1</sub> integrin was measured by the interaction of soluble VCAM-1 with Jurkat cells (e.g., American Type Culture Collection Nos. TIB 152, TIB 153, and CRL 8163), a human T-cell line which expresses high levels of α<sub>4</sub>β<sub>1</sub> integrin. VCAM-1 interacts with the cell surface in an α<sub>4</sub>β<sub>1</sub> integrin-dependent fashion (Yednock, et al. J. Bio. Chem., 1995, 270:28740).

35       Recombinant soluble VCAM-1 was expressed as a chimeric fusion protein containing the seven extracellular domains of VCAM-1 on the N-

-- 167 --

terminus and the human IgG<sub>1</sub> heavy chain constant region on the C-terminus. The VCAM-1 fusion protein was made and purified by the manner described by Yednock, *supra*.

5 Jurkat cells were grown in RPMI 1640 supplemented with 10% fetal bovine serum, penicillin, streptomycin and glutamine as described by Yednock, *supra*.

10 Jurkat cells were incubated with 1.5 mM MnCl<sub>2</sub> and 5 µg/mL 15/7 antibody for 30 minutes on ice. Mn<sup>+2</sup> activates the receptor to enhance ligand binding, and 15/7 is a monoclonal antibody that recognizes an activated/ligand occupied conformation of α<sub>4</sub>β<sub>1</sub> integrin and locks the molecule into this conformation thereby stabilizing the VCAM-1/α<sub>4</sub>β<sub>1</sub> integrin interaction. Yednock, et al., *supra*. Antibodies similar to the 15/7 antibody have been  
15 prepared by other investigators (Luque, et al, 1996, J. Bio. Chem. 271:11067) and may be used in this assay.

Cells were then incubated for 30 minutes at room temperature with candidate compounds, in various concentrations ranging from 66 µM to 0.01  
20 µM using a standard 5-point serial dilution. 15 µL soluble recombinant VCAM-1 fusion protein was then added to Jurkat cells and incubated for 30 minutes on ice. (Yednock et al., *supra*).

Cells were then washed two times and resuspended in PE-conjugated goat  
25 F(ab')<sub>2</sub> anti-mouse IgG Fc (Immunotech, Westbrook, ME) at 1:200 and incubated on ice, in the dark, for 30 minutes. Cells were washed twice and analyzed with a standard fluorescence activated cell sorter ("FACS") analysis as described in Yednock, et al., *supra*.

30 Compounds having an IC<sub>50</sub> of less than about 15µM possess binding affinity to α<sub>4</sub>β<sub>1</sub>.



-- 168 --

When tested in this assay, each of the compounds in Examples 1-135 has an  $IC_{50}$  of 15  $\mu$ M or less.

5

## Example 142

***In vitro* Saturation Assay For Determining Binding of  
Candidate Compounds to  $\alpha_4\beta_1$**

10 The following describes an *in vitro* assay to determine the plasma levels needed for a compound to be active in the Experimental Autoimmune Encephalomyelitis ("EAE") model, described in the next example, or in other *in vivo* models.

15 Log-growth Jurkat cells are washed and resuspended in normal animal plasma containing 20  $\mu$ g/ml of the 15/7 antibody (described in the above example).

20 The Jurkat cells are diluted two-fold into either normal plasma samples containing known candidate compound amounts in various concentrations ranging from 66  $\mu$ M to 0.01  $\mu$ M, using a standard 12 point serial dilution for a standard curve, or into plasma samples obtained from the peripheral blood of candidate compound-treated animals.

25 Cells are then incubated for 30 minutes at room temperature, washed twice with phosphate-buffered saline ("PBS") containing 2% fetal bovine serum and 1mM each of calcium chloride and magnesium chloride (assay medium) to remove unbound 15/7 antibody.

30 The cells are then exposed to phycoerythrin-conjugated goat F(ab')<sub>2</sub> anti-mouse IgG Fc (Immunotech, Westbrook, ME), which has been adsorbed for any non-specific cross-reactivity by co-incubation with 5% serum from the animal species being studied, at 1:200 and incubated in the dark at 4°C for 30 minutes.

-- 169 --

Cells are washed twice with assay medium and resuspended in the same. They are then analyzed with a standard fluorescence activated cell sorter ("FACS") analysis as described in Yednock et al. J. Bio. Chem., 1995, 270:28740.

5

The data is then graphed as fluorescence versus dose, e.g., in a normal dose-response fashion. The dose levels that result in the upper plateau of the curve represent the levels needed to obtain efficacy in an *in vivo* model.

10

This assay may also be used to determine the plasma levels needed to saturate the binding sites of other integrins, such as the  $\alpha_v\beta_1$  integrin, which is the integrin most closely related  $\alpha_v\beta_1$  (Palmer et al, 1993, J. Cell Bio., 123:1289). Such binding is predictive of *in vivo* utility for inflammatory conditions mediated by  $\alpha_v\beta_1$  integrin, including by way of example, airway hyper-responsiveness and occlusion that occurs with chronic asthma, smooth muscle cell proliferation in atherosclerosis, vascular occlusion following angioplasty, fibrosis and glomerular scarring as a result of renal disease, aortic stenosis, hypertrophy of synovial membranes in rheumatoid arthritis, and inflammation and scarring that occur with the progression of ulcerative colitis and Crohn's disease.

15  
20

Accordingly, the above-described assay may be performed with a human colon carcinoma cell line, SW 480 (ATTC #CCL228) transfected with cDNA encoding  $\alpha_v$  integrin (Yokosaki et al., 1994, J. Bio. Chem., 269:26691), in place of the Jurkat cells, to measure the binding of the  $\alpha_v\beta_1$  integrin. As a control, SW 480 cells which express other  $\alpha$  and  $\beta_1$  subunits may be used.

25

Accordingly, another aspect of this invention is directed to a method for treating a disease in a mammalian patient, which disease is mediated by  $\alpha_v\beta_1$ , and which method comprises administering to said patient a therapeutically effective amount of a compound of this invention. Such compounds are preferably administered in a pharmaceutical composition described herein

30

-- 170 --

above. Effective daily dosing will depend upon the age, weight, condition of the patient which factors can be readily ascertained by the attending clinician. However, in a preferred embodiment, the compounds are administered from about 20 to 500  $\mu\text{g/kg}$  per day.

5

## Example 143

*In vivo* Evaluation

10 The standard multiple sclerosis model, Experimental Autoimmune (or Allergic) Encephalomyelitis ("EAE"), was used to determine the effect of candidate compounds to reduce motor impairment in rats or guinea pigs. Reduction in motor impairment is based on blocking adhesion between leukocytes and the endothelium and correlates with anti-inflammatory activity  
15 in the candidate compound. This model has been previously described by Keszthelyi et al., Neurology, 1996, 47:1053-1059, and measures the delay of onset of disease.

20 Brains and spinal cords of adult Hartley guinea pigs were homogenized in an equal volume of phosphate-buffered saline. An equal volume of Freund's complete adjuvant (100 mg *mycobacterium tuberculosis* plus 10 ml Freund's incomplete adjuvant) was added to the homogenate. The mixture was emulsified by circulating it repeatedly through a 20 ml syringe with a peristaltic pump for about 20 minutes.

25

Female Lewis rats (2-3 months old, 170-220 g) or Hartley guinea pigs (20 day old, 180-200 g) were anesthetized with isoflurane and three injections of the emulsion, 0.1 ml each, were made in each flank. Motor impairment onset is seen in approximately 9 days.

30

Candidate compound treatment began on Day 8, just before onset of symptoms. Compounds were administered subcutaneously ("SC"), orally ("PO") or intraperitoneally ("IP"). Doses were given in a range of 10mg/kg to

-- 171 --

200 mg/kg, bid, for five days, with typical dosing of 10 to 100 mg/kg SC, 10 to 50 mg/kg PO, and 10 to 100 mg/kg IP.

Antibody GG5/3 against  $\alpha_4\beta_1$  integrin (Keszthelyi et al., Neurology, 1996, 47:1053-1059), which delays the onset of symptoms, was used as a positive control and was injected subcutaneously at 3 mg/kg on Day 8 and 11.

Body weight and motor impairment were measured daily. Motor impairment was rated with the following clinical score:

10	0	no change
	1	tail weakness or paralysis
	2	hindlimb weakness
	3	hindlimb paralysis
15	4	moribund or dead

A candidate compound was considered active if it delayed the onset of symptoms, e.g., produced clinical scores no greater than 2 or slowed body weight loss as compared to the control.

When tested in this *in vivo* assay, the compounds of Examples 5, 12, 18 and 20 were active.

25

#### Example 144

##### Asthma Model Example

Inflammatory conditions mediated by  $\alpha_4\beta_1$  integrin include, for example, airway hyper-responsiveness and occlusion that occurs with chronic asthma. The following describes an asthma model which can be used to study the *in vivo* effects of the compounds of this invention for use in treating asthma.

Following the procedures described by Abraham et al, J. Clin. Invest, 93:776-787 (1994) and Abraham et al, Am J. Respir Crit Care Med, 156:696-703 (1997), both of which are incorporated by reference in their entirety,

-- 172 --

compounds of this invention are formulated into an aerosol and administered to sheep which are hypersensitive to *Ascaris suum* antigen. Compounds which decrease the early antigen-induced bronchial response and/or block the late-phase airway response, e.g., have a protective effect against antigen-induced late responses and airway hyper-responsiveness ("AHR"), are considered to be active in this model.

Allergic sheep which are shown to develop both early and late bronchial responses to inhaled *Ascaris suum* antigen are used to study the airway effects of the candidate compounds. Following topical anesthesia of the nasal passages with 2% lidocaine, a balloon catheter is advanced through one nostril into the lower esophagus. The animals are then intubated with a cuffed endotracheal tube through the other nostril with a flexible fiberoptic bronchoscope as a guide.

Pleural pressure is estimated according to Abraham (1994). Aerosols (see formulation below) are generated using a disposable medical nebulizer that provides an aerosol with a mass median aerodynamic diameter of 3.2  $\mu\text{m}$  as determined with an Andersen cascade impactor. The nebulizer is connected to a dosimeter system consisting of a solenoid valve and a source of compressed air (20 psi). The output of the nebulizer is directed into a plastic T-piece, one end of which is connected to the inspiratory port of a piston respirator. The solenoid valve is activated for 1 second at the beginning of the inspiratory cycle of the respirator. Aerosols are delivered at  $V_T$  of 500 ml and a rate of 20 breaths/minute. A 0.5% sodium bicarbonate solution only is used as a control.

To assess bronchial responsiveness, cumulative concentration-response curves to carbachol can be generated according to Abraham (1994). Bronchial biopsies can be taken prior to and following the initiation of treatment and 24

-- 173 --

hours after antigen challenge. Bronchial biopsies can be preformed according to Abraham (1994).

5 An *in vitro* adhesion study of alveolar macrophages can be performed according to Abraham (1994), and a percentage of adherent cells is calculated.

#### Aerosol Formulation

10 A solution of the candidate compound in 0.5% sodium bicarbonate/saline (w/v) at a concentration of 30.0 mg/mL is prepared using the following procedure:

#### A. Preparation of 0.5% Sodium Bicarbonate / Saline Stock Solution: 100.0 mL

15

Ingredient	Gram / 100.0 mL	Final Concentration
Sodium Bicarbonate	0.5 g	0.5%
Saline	q.s. ad 100.0 mL	q.s. ad 100%

#### Procedure:

- 20
1. Add 0.5g sodium bicarbonate into a 100 mL volumetric flask.
  2. Add approximately 90.0 mL saline and sonicate until dissolved.
  3. Q.S. to 100.0 mL with saline and mix thoroughly.

#### B. Preparation of 30.0 mg/mL Candidate Compound: 10.0 mL

25

Ingredient	Gram / 10.0 mL	Final Concentration
Candidate Compound	0.300 g	30.0 mg/mL
0.5% Sodium Bicarbonate / Saline Stock Solution	q.s. ad 10.0 mL	q.s. ad 100%

30

-- 174 --

## Procedure:

1. Add 0.300 g of the candidate compound into a 10.0 mL volumetric  
5 flask.
2. Add approximately 9.7 mL of 0.5% sodium bicarbonate / saline  
stock solution.
3. Sonicate until the candidate compound is completely dissolved.
4. Q.S. to 10.0 mL with 0.5% sodium bicarbonate / saline stock  
10 solution and mix thoroughly.


Using a conventional oral formulation, compounds of this invention  
would be active in this model.

15

-- 175 --

## WHAT IS CLAIMED IS:

1. A compound of formula I:



where

$\text{R}^1$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

$\text{R}^2$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and where  $\text{R}^1$  and  $\text{R}^2$  together with the nitrogen atom bound to  $\text{R}^2$  and the  $\text{SO}_2$  group bound to  $\text{R}^1$  form a heterocyclic or a substituted heterocyclic group;

$\text{R}^3$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and, when  $\text{R}^2$  does not form a heterocyclic group with  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  together with the nitrogen atom bound to  $\text{R}^2$  and the carbon atom bound to  $\text{R}^3$  can form a heterocyclic or a substituted heterocyclic group;

$\text{R}^3$  is  $-(\text{CH}_2)_x\text{-Ar-R}^{3'}$  where  $\text{R}^{3'}$  is selected from the group consisting

(a) substituted alkylcarbonylamino with the proviso that at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted



-- 176 --

cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl,  
 carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted  
 heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone,  
 thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,  
 5 thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted  
 thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic,  
 substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy,  
 heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted  
 heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl,  
 10 -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-  
 heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-  
 substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl,  
 -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-  
 substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl,  
 15 -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-  
 alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-  
 substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted  
 heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic  
 where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-  
 20 (substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted  
 arylamino, mono- and di-heteroarylamino, mono- and di-substituted  
 heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted  
 heterocyclic amino, and unsymmetric di-substituted amines having different  
 substituents selected from alkyl, substituted alkyl, aryl, substituted aryl,  
 25 heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic,  
 and substituted alkyl groups having amino groups blocked by conventional  
 blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted  
 alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl,  
 -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-  
 30 aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-  
 heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen  
 or alkyl;

-- 177 --

(b) alkoxyaryl substituted on the alkoxy moiety with a substituent selected from the group consisting of carboxyl and -COOR where R is alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

(c) aryl and heteroaryl;

5 (d) -NR'R' wherein each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic with the proviso that at least one of R' is substituted  
10 alkyl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic and with the further proviso that when R' is substituted alkyl at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino,  
15 aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl,  
20 substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted  
25 heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl,  
30 -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl,

-- 178 --

-NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(e) -alkoxy-NR"R" wherein each R" is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when each R" is substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted

-- 179 --

- thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,
- 5 oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,
- 10  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is
- 15 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents
- 20 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with  $-\text{SO}_2\text{-alkyl}$ ,  $-\text{SO}_2\text{-substituted alkyl}$ ,  $-\text{SO}_2\text{-alkenyl}$ ,
- 25  $-\text{SO}_2\text{-substituted alkenyl}$ ,  $-\text{SO}_2\text{-cycloalkyl}$ ,  $-\text{SO}_2\text{-substituted cycloalkyl}$ ,  $-\text{SO}_2\text{-aryl}$ ,  $-\text{SO}_2\text{-substituted aryl}$ ,  $-\text{SO}_2\text{-heteroaryl}$ ,  $-\text{SO}_2\text{-substituted heteroaryl}$ ,  $-\text{SO}_2\text{-heterocyclic}$ ,  $-\text{SO}_2\text{-substituted heterocyclic}$  and  $-\text{SO}_2\text{NRR}$  where R is hydrogen or alkyl;
- (f) substituted alkenyl or substituted alkynyl with the proviso that at least
- 30 one of the substituents on the substituted alkenyl/alkynyl moiety is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl,

-- 180 --

heterocyclic, and substituted heterocyclic with the proviso that when substituted with substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,

-- 181 --

substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(g) substituted aryloxy and substituted heteroaryloxy with the proviso that at least one substituent on the substituted aryloxy/heteroaryloxy is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(h) -alkoxy-saturated heterocyclic, -alkoxy-saturated substituted heterocyclic, -substituted alkoxy-heterocyclic and -substituted alkoxy-substituted saturated heterocyclic;

(i) -O-heterocyclic and -O-substituted heterocyclic;

(j) tetrazolyl;

(k) -NR-SO<sub>2</sub>-substituted alkyl where R is hydrogen, alkyl or aryl, with the proviso that at least one substituent on the alkyl moiety of the substituted alkylsulfonylamino is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(l) alkenylsulfonylamino, alkynylsulfonylamino, substituted alkenylsulfonylamino and substituted alkynylsulfonylamino;

(m) substituted alkoxy with the proviso that the substitution on the alkyl moiety of said substituted alkoxy does not include alkoxy-NR"R", unsaturated heterocyclyl, alkyloxy, aryloxy, heteroaryloxy, aryl, heteroaryl and

-- 182 --

aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(n) amidine and amidine substituted with from 1 to 3 substituents independently selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

(o)  $-C(O)NR''R'''$  where each  $R'''$  is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic with the proviso that when one  $R'''$  is unsaturated heterocyclalkyl, aryl, heteroaryl or aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea, then the other  $R'''$  is alkyl, substituted alkyl (other than unsaturated heterocyclalkyl substituted-alkyl), cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl and heterocyclic and substituted heterocyclic;

(p)  $-NR^{12}C(O)-R^8$  where  $R^8$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and  $R^{12}$  is alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(q)  $-SO_2$ -aryl,  $-SO_2$ -substituted aryl,  $-SO_2$ -heteroaryl,  $-SO_2$ -substituted heteroaryl or  $-SO_2$ -alkyl;

(r)  $-NR'C(O)NR^9R^9$  wherein  $R'$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted

-- 183 --

cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and each  $R^9$  is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic:

(s)  $-NR'C(O)OR^9$  wherein  $R'$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and,  $R^9$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(t) -aminocarbonyl-(N-formylheterocycl); and

(u) -alkyl-C(O)NH-heterocycl and -alkyl-C(O)NH-substituted heterocycl,


Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl,

x is an integer of from 1 to 4;

Q is  $-C(X)NR^7$  wherein  $R^7$  is selected from the group consisting of hydrogen and alkyl; and X is selected from the group consisting of oxygen and sulfur;

and pharmaceutically acceptable salts thereof.

2. A compound of formula IA:



where

$R^1$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;



-- 184 --

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and R<sup>1</sup> and R<sup>2</sup> together with the nitrogen atom bound to R<sup>2</sup> and the SO<sub>2</sub> group bound to R<sup>1</sup> can form a heterocyclic or a substituted heterocyclic group;

R<sup>3</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and, when R<sup>2</sup> does not form a heterocyclic group with R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> and the carbon atom bound to R<sup>3</sup> can form a heterocyclic or a substituted heterocyclic group;

R<sup>5</sup> is -(CH<sub>2</sub>)<sub>x</sub>-Ar-R<sup>5'</sup> where R<sup>5'</sup> is selected from the group consisting

(a) substituted alkylcarbonylamino with the proviso that at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted

heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl,

-- 185 --

- NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic
- where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted

-- 186 --

thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-

5 cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted

10 thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-

15 substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-

20 heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted

25 alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents

30 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted

-- 187 --

alkyl groups substituted with  $-\text{SO}_2$ -alkyl,  $-\text{SO}_2$ -substituted alkyl,  $-\text{SO}_2$ -alkenyl,  $-\text{SO}_2$ -substituted alkenyl,  $-\text{SO}_2$ -cycloalkyl,  $-\text{SO}_2$ -substituted cycloalkyl,  $-\text{SO}_2$ -aryl,  $-\text{SO}_2$ -substituted aryl,  $-\text{SO}_2$ -heteroaryl,  $-\text{SO}_2$ -substituted heteroaryl,  $-\text{SO}_2$ -heterocyclic,  $-\text{SO}_2$ -substituted heterocyclic and  $-\text{SO}_2\text{NRR}$  where R is hydrogen or alkyl;

(e) -alkoxy-NR"R" wherein each R" is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when each R" is substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2$ -alkyl,  $-\text{OS}(\text{O})_2$ -substituted alkyl,  $-\text{OS}(\text{O})_2$ -aryl,  $-\text{OS}(\text{O})_2$ -substituted aryl,  $-\text{OS}(\text{O})_2$ -heteroaryl,  $-\text{OS}(\text{O})_2$ -substituted heteroaryl,  $-\text{OS}(\text{O})_2$ -heterocyclic,  $-\text{OS}(\text{O})_2$ -substituted heterocyclic,  $-\text{OSO}_2$ -NRR where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2$ -alkyl,  $-\text{NRS}(\text{O})_2$ -substituted alkyl,  $-\text{NRS}(\text{O})_2$ -aryl,  $-\text{NRS}(\text{O})_2$ -substituted aryl,  $-\text{NRS}(\text{O})_2$ -heteroaryl,  $-\text{NRS}(\text{O})_2$ -substituted heteroaryl,  $-\text{NRS}(\text{O})_2$ -heterocyclic,  $-\text{NRS}(\text{O})_2$ -substituted heterocyclic,  $-\text{NRS}(\text{O})_2$ -NR-alkyl,

-- 188 --

-NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(f) substituted alkenyl or substituted alkynyl with the proviso that at least one of the substituents on the substituted alkenyl/alkynyl moiety is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when substituted with substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl,

-- 189 --

- substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;
- (g) substituted aryloxy and substituted heteroaryloxy with the proviso that at least one substituent on the substituted aryloxy/heteroaryloxy is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl,

-- 190 --

alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

5 (h) -alkoxy-saturated heterocyclic, -alkoxy-saturated substituted heterocyclic, -substituted alkoxy-heterocyclic and -substituted alkoxy-substituted saturated heterocyclic;

(i) -O-heterocyclic and -O-substituted heterocyclic;

(j) tetrazolyl;

10 (k) -NR-SO<sub>2</sub>-substituted alkyl where R is hydrogen, alkyl or aryl, with the proviso that at least one substituent on the alkyl moiety of the substituted alkylsulfonylamino is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, 15 alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(l) alkenylsulfonylamino, alkynylsulfonylamino, substituted alkenylsulfonylamino and substituted alkynylsulfonylamino;

(m) substituted alkoxy with the proviso that the substitution on the alkyl 20 moiety of said substituted alkoxy does not include alkoxy-NR"R", unsaturated heterocyclyl, alkyloxy, aryloxy, heteroaryloxy, aryl, heteroaryl and aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, 25 alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(n) amidine and amidine substituted with from 1 to 3 substituents independently selected from alkyl, substituted alkyl, alkenyl, substituted

alkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

30 (o) -C(O)NR"R" where each R" is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl,

-- 191 --

substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic with the proviso that when one R<sup>'''</sup> is unsaturated heterocyclylalkyl, aryl, heteroaryl or aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea, then the other R<sup>'''</sup> is alkyl, substituted alkyl (other than unsaturated heterocyclyl substituted-alkyl), cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl and heterocyclic and substituted heterocyclic;

(p) -NR<sup>12</sup>C(O)-R<sup>8</sup> where R<sup>8</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and R<sup>12</sup> is alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(q) -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl or -SO<sub>2</sub>-alkyl;

(r) -NR'<sup>9</sup>C(O)NR<sup>9</sup>R<sup>9</sup> wherein R' is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and each R<sup>9</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(s) -NR'<sup>9</sup>C(O)OR<sup>9</sup> wherein R' is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and, R<sup>9</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;



-- 192 --

(t) -aminocarbonyl-(N-formylheterocyclyl); and

(u) -alkyl-C(O)NH-heterocyclyl and -alkyl-C(O)NH-substituted heterocyclyl,

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl,

x is an integer of from 1 to 4;

R<sup>6</sup> is selected from the group consisting of 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), amino, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, -O-(N-succinimidyl), -NH-adamantyl, -O-cholest-5-en-3-β-yl, -NHOY where Y is hydrogen, alkyl, substituted alkyl, aryl, and substituted aryl, -NH(CH<sub>2</sub>)<sub>p</sub>COOY where p is an integer of from 1 to 8 and Y is as defined above, -OCH<sub>2</sub>NR<sup>9</sup>R<sup>10</sup> where R<sup>9</sup> is selected from the group consisting of -C(O)-aryl and -C(O)-substituted aryl and R<sup>10</sup> is selected from the group consisting of hydrogen and -CH<sub>2</sub>COOR<sup>11</sup> where R<sup>11</sup> is alkyl, and -NHSO<sub>2</sub>Z where Z is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

Q is -C(X)NR<sup>7</sup>- wherein R<sup>7</sup> is selected from the group consisting of hydrogen and alkyl; and X is selected from the group consisting of oxygen and sulfur,

and pharmaceutically acceptable salts thereof

with the proviso that when R<sup>1</sup> is *p*-CH<sub>3</sub>-φ-, R<sup>6</sup> is methoxy, Q is -C(O)NH-, and R<sup>2</sup> and R<sup>3</sup> are joined to form a pyrrolidinyl group, then R<sup>5</sup> is not *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(isopropyl)]<sub>2</sub>-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-1-pyrrolidinyl]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-pyrimidinyl)piperazinyl]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-N-morpholinyl]-benzyl-, or *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-N-piperidinyl]-benzyl-.

3. A compound according to Claims 1 or 2 wherein R<sup>1</sup> is selected

from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl.

-- 193 --

4. A compound according to Claim 3 wherein R<sup>1</sup> is selected from the group consisting of 4-methylphenyl, methyl, benzyl, *n*-butyl, 4-chlorophenyl, 1-naphthyl, 2-naphthyl, 4-methoxyphenyl, phenyl, 2,4,6-trimethylphenyl, 2-(methoxycarbonyl)phenyl, 2-carboxyphenyl, 3,5-dichlorophenyl, 4-trifluoromethylphenyl, 3,4-dichlorophenyl, 3,4-dimethoxyphenyl, 4-(CH<sub>3</sub>C(O)NH-)phenyl, 4-trifluoromethoxyphenyl, 4-cyanophenyl, isopropyl, 3,5-di-(trifluoromethyl)phenyl, 4-*t*-butylphenyl, 4-*t*-butoxyphenyl, 4-nitrophenyl, 2-thienyl, 1-N-methyl-3-methyl-5-chloropyrazol-4-yl, phenethyl, 1-N-methylimidazol-4-yl, 4-bromophenyl, 4-amidinophenyl, 4-methylamidinophenyl, 4-[CH<sub>3</sub>SC(=NH)]phenyl, 5-chloro-2-thienyl, 2,5-dichloro-4-thienyl, 1-N-methyl-4-pyrazolyl, 2-thiazolyl, 5-methyl-1,3,4-thiadiazol-2-yl, 4-[H<sub>2</sub>NC(S)]phenyl, 4-aminophenyl, 4-fluorophenyl, 2-fluorophenyl, 3-fluorophenyl, 3,5-difluorophenyl, pyridin-3-yl, pyrimidin-2-yl, 4-(3'-dimethylamino-*n*-propoxy)-phenyl, and 1-methylpyrazol-4-yl.
5. A compound according to Claims 1 or 2 wherein R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, phenyl, benzyl, -(CH<sub>2</sub>)<sub>2</sub>-2-thienyl, and -(CH<sub>2</sub>)<sub>2</sub>-φ.
6. A compound according to Claims 1 or 2 wherein R<sup>1</sup> and R<sup>2</sup> together with the nitrogen atom bound to R<sup>2</sup> and the SO<sub>2</sub> group bound to R<sup>1</sup> are joined to form a heterocyclic group or substituted heterocyclic group.
7. A compound according to Claims 1 or 2 wherein R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> substituent and the carbon bound to the R<sup>3</sup> substituent form a heterocyclic group or a substituted heterocyclic group.
8. A compound according to Claims 1 or 2 wherein R<sup>3</sup> is selected from the group consisting of methyl, phenyl, benzyl, diphenylmethyl, -CH<sub>2</sub>CH<sub>2</sub>-COOH, -CH<sub>2</sub>-COOH, 2-amidoethyl, *iso*-butyl, *t*-butyl, -CH<sub>2</sub>O-benzyl and hydroxymethyl.

9. A compound according to Claims 1 or 2 wherein Q is -C(O)NH- or -C(S)NH-.

- 5 10. A compound according to Claims 1 or 2 wherein R<sup>5</sup> is selected from the group consisting of:
- 4-[NH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-[HOOCCH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>NHBoc]benzyl, 4-[-NHC(O)CH(CH<sub>3</sub>)NHBoc]benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>φ)NHBoc]benzyl, 4-[-NHC(O)CH<sub>2</sub>NHC(O)NH-3'-  
 10 methylphenyl]benzyl, 4-[-NHC(O)CH(NHBoc)(CH<sub>2</sub>)<sub>4</sub>NHCbz]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>CH(C(O)OCH<sub>2</sub>φ)-NHCbz]benzyl, 4-φ-benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)NHBoc]benzyl,  
 4-[H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(BocHNCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-)benzyl, 4-[φCH<sub>2</sub>OCH<sub>2</sub>(BocHN)CHC(O)NH-]benzyl,  
 15 4-[CH<sub>3</sub>NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(N-methylpiperidin-4-oxy)-benzyl,  
 4-[CH<sub>3</sub>N(Boc)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[φCH<sub>2</sub>OCH<sub>2</sub>(H<sub>2</sub>N)CHC(O)NH-]benzyl, 4-[HO(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-]benzyl, 4-[φCH<sub>2</sub>O(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 4-[HO(O)C(NH<sub>2</sub>)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 20 4-[CH<sub>3</sub>(N-Boc)NCH<sub>2</sub>C(O)NH-]benzyl, 4-[CH<sub>3</sub>NHCH<sub>2</sub>C(O)NH-]benzyl,  
 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>C(O)NH-]benzyl, 4-[-O-CH(COOH)φ]benzyl,  
 4-[2-carboxylphenyl]-benzyl, 4-[2-carboxylmethylphenyl]-benzyl  
 4-[φCH<sub>2</sub>OC(O)NHCH<sub>2</sub>CH<sub>2</sub>NH-]benzyl, 4-N[-(SO<sub>2</sub>)CH<sub>3</sub>]-  
 CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]benzyl, 4-*t*-butyl-O(O)CCH<sub>2</sub>-O-benzylNH]benzyl,  
 25 4-[N,N-di(4-N,N-dimethylamino)benzyl)amino]benzyl,  
 4-(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl-CH<sub>2</sub>NH)benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1'-(4'-pyrimidinyl)-piperazinyl]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-pyrrolidinyl)]-  
 benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)]-benzyl,  
 30 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-O-]benzyl, 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>O-]benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(1'-(4'-methylpiperazinyl))]benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-4-(3'-chlorophenyl)-piperazin-1-yl]-benzyl,

- 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(φ)CH<sub>2</sub>CH<sub>3</sub>]-benzyl, 4-[-OCH<sub>2</sub>-3'-(N-Boc)-piperidinyl]-benzyl, 4-[-O-(3-(N-Boc)-piperidinyl)]benzyl,  
 3-[-O-(N-methylpiperidin-4-yl)]benzyl, 4-[-O-(N-methylpiperidin-4-yl)]benzyl,  
 4-[di-*iso*-propylamino-CH<sub>2</sub>CH<sub>2</sub>O-]-benzyl, 4-[*N*-3-methylbutyl-*N*-trifluoro-  
 5 methanesulfonyl)amino]benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-morpholinyl)]-benzyl,  
 4-[-OCH<sub>2</sub>CH(NHBoc)CH<sub>2</sub>cyclohexyl]-benzyl, 4-[OCH<sub>2</sub>CH<sub>2</sub>-(N-piperidinyl)]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(4-*m*-chlorophenyl)-piperazin-1-yl]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-homopiperidinyl)]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(benzyl)<sub>2</sub>]-benzyl,  
 3-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl,  
 10 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)benzyl]-benzyl, 4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethyl-O]-benzyl, [cyclopentylacetylenyl]-benzyl, 4-[-C≡C-φ-4'φ]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-S(O)<sub>2</sub>-4'-CH<sub>3</sub>-φ]-benzyl, 4-[-C≡C-CH<sub>2</sub>NHC(O)NH<sub>2</sub>]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-(4'-COOCH<sub>2</sub>CH<sub>3</sub>)φ]-benzyl, 4-[-C≡C-CH(NH<sub>2</sub>)-cyclohexyl]-  
 15 benzyl, 4-[-C≡C-CH<sub>2</sub>-O-phenyl]-benzyl, 4-[-C≡C-CH<sub>2</sub>OCH<sub>3</sub>]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-(4'-C(O)OC<sub>2</sub>H<sub>5</sub>)phenyl]-benzyl, 4-[-C≡C-CH<sub>2</sub>CH(C(O)OCH<sub>3</sub>)<sub>2</sub>]-benzyl, 4-[-C≡C-CH<sub>2</sub>CH(NHC(O)CH<sub>3</sub>)C(O)OH]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>NH-(4,5-dihydro-4-oxo-5-phenyl-oxazol-2-yl)]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(N-morpholino)]-benzyl, 4-[-OCH<sub>2</sub>COOH]-benzyl,  
 20 4-[-OCH<sub>2</sub>COO-*t*-butyl]-benzyl, 4-[-N(SO<sub>2</sub>CH<sub>3</sub>)(CH<sub>2</sub>)<sub>3</sub>-N(CH<sub>3</sub>)<sub>2</sub>]-benzyl,  
 4-[-NHS(O)<sub>2</sub>CF<sub>3</sub>]-benzyl, 4-[-C(=NH)NH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>Cl]-benzyl,  
 4-[-OCH<sub>2</sub>C(O)NH-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)O-benzyl]-benzyl,  
 4-[-OCH<sub>2</sub>C(O)OH]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-(3-methoxypyrrol-2-yl)-piperazinyl)-benzyl, 4-[-OCH<sub>2</sub>C(O)NH<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-*t*-butyl]-benzyl,  
 25 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-phenyl)-piperidinyl]-benzyl,  
 4-[-NHSO<sub>2</sub>-CH=CH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>Cl]-benzyl, 4-benzyl-benzyl, 4-[-OCH<sub>2</sub>C(O)piperidin-1-yl]benzyl,  
 4-[-OCH<sub>2</sub>C(O)N(CH(CH<sub>3</sub>)<sub>2</sub>)<sub>2</sub>]-benzyl, 4-amidinobenzyl, 4-acetamidobenzyl,  
 4-(N-methyl)acetamidobenzyl, 4-[-NHC(O)CH<sub>2</sub>NHC(O)NH-fluorescin]benzyl,  
 30 4-[-NHC(O)CH<sub>2</sub>CH(NH<sub>2</sub>)COOH, (1-toluenesulfonylimidazol-4-yl)-methyl-, [(1-*N,N*-dimethylaminosulfonyl)-imidazol-4-yl]methyl-, 4-(*N*-toluenesulfonyl-amino)benzyl, and 4-[*N*-methyltrifluoroacetamido]phenyl.

-- 196 --

11. A compound according to Claim 2 wherein R<sup>6</sup> is selected from the group consisting of 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), methoxy, ethoxy, *iso*-propoxy, *n*-butoxy, *t*-butoxy, cyclopentoxy, *neo*-pentoxy, 2- $\alpha$ -*iso*-propyl-4- $\beta$ -methylcyclohexoxy, 2- $\beta$ -isopropyl-4- $\beta$ -methylcyclohexoxy,  
 5 -NH<sub>2</sub>, benzyloxy, -NHCH<sub>2</sub>COOH, -NHCH<sub>2</sub>CH<sub>2</sub>COOH, -NH-adamantyl, -NHCH<sub>2</sub>CH<sub>2</sub>COOCH<sub>2</sub>CH<sub>3</sub>, -NHSO<sub>2</sub>-*p*-CH<sub>3</sub>- $\phi$ , -NHOR<sup>8</sup> where R<sup>8</sup> is hydrogen, methyl, *iso*-propyl or benzyl, O-(N-succinimidyl),  
 -O-cholest-5-en-3- $\beta$ -yl, -OCH<sub>2</sub>-OC(O)C(CH<sub>3</sub>)<sub>2</sub>, -O(CH<sub>2</sub>)<sub>z</sub>NHC(O)W where z is 1 or 2 and W is selected from the group consisting of pyrid-3-yl, N-  
 10 methylpyridyl, and N-methyl-1,4-dihydro-pyrid-3-yl, -NR''C(O)-R' where R' is aryl, heteroaryl or heterocyclic and R'' is hydrogen or  
 -CH<sub>2</sub>C(O)OCH<sub>2</sub>CH<sub>3</sub>.

12. A compound selected from the group consisting of:

15

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)glycyl]amino]-L-phenylalanine

20

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(glycyl)amino]-L-phenylalanine

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(carboxy)propionamido]-L-phenylalanine

25

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)-L-alanyl]amino]-L-phenylalanine

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)-D-alanyl]amino]-L-phenylalanine

30

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)-D-phenylalanyl]amino]-L-phenylalanine

35

*N*-(toluene-4-sulfonyl)-L-prolyl-4-(2-[3-(fluorescein)thiourido]acetamido)-L-phenylalanine

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)glycyl]amino]-L-phenylalanine methyl ester

40

*N*-(toluene-4-sulfonyl)-L-prolyl-4-(2-[3-(3-methylphenyl)urido]acetamido)-L-phenylalanine

-- 197 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*α-*tert*-butoxycarbonyl)-*N*ε-carbobenzyloxy-L-lysyl]amino]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[γ-(α-benzyl-*N*α-carbobenzyloxy-L-aspartyl)amino]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*α-*tert*-butoxycarbonyl)-L-lysyl]amino]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[γ-(L-aspartyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(4-aminobutyramido)-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-*tert*-butoxycarbonylamino)butyramido]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-methylamino)butyramido]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-*tert*-butoxycarbonyl)-*N*-methylamino]butyramido]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*O*-benzyl)-L-seryl]amino]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[δ-(D,L-glutamyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonylsarcosyl)amino]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N*-*tert*-butoxycarbonylsarcosyl)amino]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine ethyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(sarcosyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine

-- 198 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-( $\alpha$ -carboxybenzyloxy)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenyl]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(methoxycarbonyl)phenyl]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[3-(*N,N*-dimethylamino)propyl]-*N*-[trifluoromethanesulfonyl]amino}-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[4-[(*tert*-butoxycarbonyl)methoxy]benzyl]amino}-L-phenylalanine methyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N,N*-di[4-(*N,N*-dimethylamino)benzyl]amino}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl)methyl]amino}-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-*N*-methyl-L-serinyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N*-ethyl-*N*-phenylamino)ethoxy]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-diisopropylamino)ethoxy]-L-phenylalanine
- 
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-cyclohexyl-2-(*N-tert*-butoxycarbonylamino)propoxy]-L-phenylalanine methyl ester
- 45 *N*-(thiophene-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

-- 199 --

- N*-(5-chlorothiophene-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)-propoxy]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-diethylamino)ethoxy]-L-phenylalanine
- N*-(2,5-dichlorothiophene-3-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 10 *N*-(1-methylpyrazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)-propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-diethylamino)propoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(thiazole-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-diethylamino)propoxy]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(1-methylimidazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)-propoxy]-L-phenylalanine
- N*-(2-methylthiadiazole-5-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)-propoxy]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(4-cyanobenzenesulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-(3,3-dimethyl)prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine ethyl ester



-- 200 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine methyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-(thiamorpholin-3-carbonyl)-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(cyclopentyl)ethynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(4-(phenyl)phenyl)ethynyl]-D,L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(toluene-4-sulfonyloxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(urcido)prop-1-ynyl]-D,L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(1-aminocyclohex-1-yl)ethynyl]-D,L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(phenoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(phenoxy)prop-1-ynyl]-D,L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(methoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(methoxy)prop-1-ynyl]-D,L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4,4-di(methoxycarbonyl)but-1-ynyl]-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4,4-di(methoxycarbonyl)but-1-ynyl]-D,L-phenylalanine

-- 201 --

- N*-(toluene-4-sulfonyl)sarcosyl-4-{4,4-di(methoxycarbonyl)but-1-ynyl}-D,L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(4-acetamido-4-carboxybut-1-ynyl)-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-(4-acetamido-4-carboxybut-1-ynyl)-D,L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[(4,5-dihydro-4-oxo-5-phenyloxazol-2-yl)amino]prop-1-ynyl}-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-{3-[(4,5-dihydro-4-oxo-5-phenyloxazol-2-yl)amino]prop-1-ynyl}-D,L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[4-(pyrimidin-2-yl)piperazin-1-yl]ethoxy}-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(piperidin-1-yl)ethoxy]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine methyl ester

-- 202 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[4-(3-methoxythien-2-yl)-4-hydroxypiperidin-1-yl]ethoxy}-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-3-(1-methylpiperidin-4-oxy)-D,L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(1-methylpiperidin-4-oxy)-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-(1-methylpiperidin-4-oxy)-L-phenylalanine ethyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-(1,1-dioxothiomorpholin-3-carbonyl)-4-(1-methylpiperidin-4-oxy)-L-phenylalanine ethyl ester
- N*-(toluene-4-sulfonyl)-L-(1,1-dioxothiomorpholin-3-carbonyl)-4-(1-methylpiperidin-4-oxy)-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-(1-methylpiperidin-4-oxy)-L-phenylalanine
- 40 *N*-( $\alpha$ -toluenesulfonyl)-L-prolyl-4-(1-methylpiperidin-4-oxy)-L-phenylalanine ethyl ester
- 
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(trifluoromethanesulfonyl)amino-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(trifluoromethanesulfonyl)amino-L-phenylalanine

-- 203 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(chloromethanesulfonyl)amino-L-phenylalanine methyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-trifluoromethanesulfonyl)-*N*-isobutylamino-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-benzylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine methyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(aminocarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butylaminocarbonyl)methoxy]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(4-phenyl-4-hydroxypiperidin-1-yl)ethoxy]-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(piperidin-1-ylcarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 45 *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(amidino)phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(aminocarbonyl)phenylalanine

-- 204 --

*N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine isopropyl ester

*N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine

*N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methyltrifluoroacetamido)-L-phenylalanine methyl ester

*N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine *t*-butyl ester

*N*-(toluene-4-sulfonyl)-L-prolyl-L-4-(*N*-methylpiperidinoxy)-phenylalanine *t*-butyl ester


*N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-L-(4-methylpiperidinoxy) phenylalanine *t*-butyl ester

*N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine *t*-butyl ester

*N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine

and pharmaceutically acceptable salts thereof as well as any of the ester compounds recited above wherein one ester is replaced with another ester selected from the group consisting of methyl ester, ethyl ester, *n*-propyl ester, isopropyl ester, *n*-butyl ester, isobutyl ester, *sec*-butyl ester and *tert*-butyl ester.

13. A method for binding VLA-4 in a biological sample which method comprises contacting the biological sample with a compound :



IA

where

$\text{R}^1$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

-- 205 --

$R^2$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $R^1$  and  $R^2$  together with the nitrogen atom bound to  $R^2$  and the  $SO_2$  group bound to  $R^1$  can form a heterocyclic or a substituted heterocyclic group;

$R^3$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and, when  $R^2$  does not form a heterocyclic group with  $R^1$ ,  $R^2$  and  $R^3$  together with the nitrogen atom bound to  $R^2$  and the carbon atom bound to  $R^3$  can form a heterocyclic or a substituted heterocyclic group;

$R^5$  is  $-(CH_2)_x-Ar-R^5$  where  $R^5$  is selected from the group consisting

(a) substituted alkylcarbonylamino with the proviso that at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-OS(O)_2$ -alkyl,  $-OS(O)_2$ -substituted alkyl,  $-OS(O)_2$ -aryl,  $-OS(O)_2$ -substituted aryl,  $-OS(O)_2$ -heteroaryl,  $-OS(O)_2$ -substituted heteroaryl,  $-OS(O)_2$ -heterocyclic,  $-OS(O)_2$ -substituted heterocyclic,  $-OSO_2$ -NRR where R is hydrogen or alkyl,

-- 206 --

- NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;
- (b) alkoxyaryl substituted on the alkoxy moiety with a substituent selected from the group consisting of carboxyl and -COOR where R is alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,
- (c) aryl and heteroaryl;
- (d) -NR'R' wherein each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic with the proviso that at least one of R' is substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic and with the further proviso that when R' is substituted alkyl at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino,

-- 207 --

thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino,  
 thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino,  
 aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen,  
 hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-  
 5 cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted  
 aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl,  
 carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl,  
 substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl,  
 substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted  
 10 thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic,  
 substituted thioheterocyclic, heterocyclic, substituted heterocyclic,  
 cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted  
 heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
 oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-  
 15 substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl,  
 -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted  
 heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl,  
 -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl,  
 -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-  
 20 heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl,  
 -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted  
 aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl,  
 -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is  
 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted  
 25 alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino,  
 mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino,  
 mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic  
 amino, and unsymmetric di-substituted amines having different substituents  
 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,  
 30 substituted heteroaryl, heterocyclic and substituted heterocyclic, and  
 substituted alkyl groups having amino groups blocked by conventional  
 blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted



-- 208 --

alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(e) -alkoxy-NR"R" wherein each R" is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when each R" is substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl,

-- 209 --

-NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(f) substituted alkenyl or substituted alkynyl with the proviso that at least one of the substituents on the substituted alkenyl/alkynyl moiety is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when substituted with substituted alkyl then at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl,

-- 210 --

- substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with  $-\text{SO}_2\text{-alkyl}$ ,  $-\text{SO}_2\text{-substituted alkyl}$ ,  $-\text{SO}_2\text{-alkenyl}$ ,  $-\text{SO}_2\text{-substituted alkenyl}$ ,  $-\text{SO}_2\text{-cycloalkyl}$ ,  $-\text{SO}_2\text{-substituted cycloalkyl}$ ,  $-\text{SO}_2\text{-aryl}$ ,  $-\text{SO}_2\text{-substituted aryl}$ ,  $-\text{SO}_2\text{-heteroaryl}$ ,  $-\text{SO}_2\text{-substituted heteroaryl}$ ,  $-\text{SO}_2\text{-heterocyclic}$ ,  $-\text{SO}_2\text{-substituted heterocyclic}$  and  $-\text{SO}_2\text{-NRR}$  where R is hydrogen or alkyl;
- (g) substituted aryloxy and substituted heteroaryloxy with the proviso that at least one substituent on the substituted aryloxy/heteroaryloxy is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl,

-- 211 --

alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

5 (h) -alkoxy-saturated heterocyclic, -alkoxy-saturated substituted heterocyclic, -substituted alkoxy-heterocyclic and -substituted alkoxy-substituted saturated heterocyclic;

(i) -O-heterocyclic and -O-substituted heterocyclic;

(j) tetrazolyl;

10 (k) -NR-SO<sub>2</sub>-substituted alkyl where R is hydrogen, alkyl or aryl, with the proviso that at least one substituent on the alkyl moiety of the substituted alkylsulfonylamino is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

15 (l) alkenylsulfonylamino, alkynylsulfonylamino, substituted alkenylsulfonylamino and substituted alkynylsulfonylamino;

(m) substituted alkoxy with the proviso that the substitution on the alkyl moiety of said substituted alkoxy does not include alkoxy-NR"R", unsaturated heterocyclyl, alkylloxy, aryloxy, heteroaryloxy, aryl, heteroaryl and aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

25 (n) amidine and amidine substituted with from 1 to 3 substituents independently selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

30 (o) -C(O)NR""R"" where each R"" is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl,

-- 212 --

substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic with the proviso that when one R<sup>'''</sup> is unsaturated heterocyclalkyl, aryl, heteroaryl or aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea, then the other R<sup>'''</sup> is alkyl, substituted alkyl (other than unsaturated heterocyclalkyl substituted-alkyl), cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl and heterocyclic and substituted heterocyclic;

(p) -NR<sup>12</sup>C(O)-R<sup>8</sup> where R<sup>8</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and R<sup>12</sup> is alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(q) -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl or -SO<sub>2</sub>-alkyl;

(r) -NR'C(O)NR<sup>9</sup>R<sup>9</sup> wherein R' is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and each R<sup>9</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

(s) -NR'C(O)OR<sup>9</sup> wherein R' is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and, R<sup>9</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

-- 213 --

(t) -aminocarbonyl-(N-formylheterocycl); and

(u) -alkyl-C(O)NH-heterocycl and -alkyl-C(O)NH-substituted heterocycl,

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl,

x is an integer of from 1 to 4;

R<sup>6'</sup> is selected from the group consisting of, hydroxy, 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), amino, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, -O-(N-succinimidyl), -NH-adamantyl, -O-cholest-5-en-3-β-yl, -NHOY where Y is hydrogen, alkyl, substituted alkyl, aryl, and substituted aryl, -NH(CH<sub>2</sub>)<sub>p</sub>COOY where p is an integer of from 1 to 8 and Y is as defined above, -OCH<sub>2</sub>NR<sup>9</sup>R<sup>10</sup> where R<sup>9</sup> is selected from the group consisting of -C(O)-aryl and -C(O)-substituted aryl and R<sup>10</sup> is selected from the group consisting of hydrogen and -CH<sub>2</sub>COOR<sup>11</sup> where R<sup>11</sup> is alkyl, and -NHSO<sub>2</sub>Z where Z is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

Q is -C(X)NR<sup>7</sup>- wherein R<sup>7</sup> is selected from the group consisting of hydrogen and alkyl; and X is selected from the group consisting of oxygen and sulfur;

and pharmaceutically acceptable salts thereof

with the proviso that when R<sup>1</sup> is *p*-CH<sub>3</sub>-φ-, R<sup>6</sup> is methoxy, Q is -C(O)NH-, and R<sup>2</sup> and R<sup>3</sup> are joined to form a pyrrolidinyl group, then R<sup>3</sup> is not *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(isopropyl)]<sub>2</sub>-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-1-pyrrolidinyl]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-pyrimidinyl)piperazinyl]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-N-morpholinyl]-benzyl-, or *p*-[-OCH<sub>2</sub>CH<sub>2</sub>-N-piperidinyl]-benzyl-

under conditions wherein said compound binds to VLA-4.

14. The method according to Claim 13 wherein R<sup>1</sup> is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl.

-- 214 --

15. The method according to Claim 14 wherein R<sup>1</sup> is selected from the group consisting of 4-methylphenyl, methyl, benzyl, *n*-butyl, 4-chlorophenyl, 1-naphthyl, 2-naphthyl, 4-methoxyphenyl, phenyl, 2,4,6-trimethylphenyl, 2-(methoxycarbonyl)phenyl, 2-carboxyphenyl, 3,5-dichlorophenyl, 4-trifluoromethylphenyl, 3,4-dichlorophenyl, 3,4-dimethoxyphenyl, 4-(CH<sub>3</sub>C(O)NH-)phenyl, 4-trifluoromethoxyphenyl, 4-cyanophenyl, isopropyl, 3,5-di-(trifluoromethyl)phenyl, 4-*t*-butylphenyl, 4-*t*-butoxyphenyl, 4-nitrophenyl, 2-thienyl, 1-N-methyl-3-methyl-5-chloropyrazol-4-yl, phenethyl, 1-N-methylimidazol-4-yl, 4-bromophenyl, 4-amidinophenyl, 4-methylamidinophenyl, 4-[CH<sub>3</sub>SC(=NH)]phenyl, 5-chloro-2-thienyl, 2,5-dichloro-4-thienyl, 1-N-methyl-4-pyrazolyl, 2-thiazolyl, 5-methyl-1,3,4-thiadiazol-2-yl, 4-[H<sub>2</sub>NC(S)]phenyl, 4-aminophenyl, 4-fluorophenyl, 2-fluorophenyl, 3-fluorophenyl, 3,5-difluorophenyl, pyridin-3-yl, pyrimidin-2-yl, 4-(3'-dimethylamino-*n*-propoxy)-phenyl, and 1-methylpyrazol-4-yl.
16. The method according to Claim 13 wherein R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, phenyl, benzyl, -(CH<sub>2</sub>)<sub>2</sub>-2-thienyl, and -(CH<sub>2</sub>)<sub>2</sub>-φ.
17. The method according to Claim 13 wherein R<sup>1</sup> and R<sup>2</sup> together with the nitrogen atom bound to R<sup>2</sup> and the SO<sub>2</sub> group bound to R<sup>1</sup> are joined to form a heterocyclic group or substituted heterocyclic group.
18. The method according to Claim 13 wherein R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> substituent and the carbon bound to the R<sup>3</sup> substituent form a heterocyclic group or a substituted heterocyclic group.
19. The method according to Claim 13 wherein R<sup>3</sup> is selected from the group consisting of methyl, phenyl, benzyl, diphenylmethyl, -CH<sub>2</sub>CH<sub>2</sub>-COOH, -CH<sub>2</sub>-COOH, 2-amidoethyl, *iso*-butyl, *t*-butyl, -CH<sub>2</sub>O-benzyl and hydroxymethyl.

-- 215 --

20. The method according to Claim 13 wherein Q is -C(O)NH- or -C(S)NH-.

- 5 21. The method according to Claim 13 wherein R<sup>3</sup> is selected from the group consisting of:
- 4-[NH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-[HOOCCH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>NHBoc]benzyl, 4-[-NHC(O)CH(CH<sub>3</sub>)NHBoc]benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>φ)NHBoc]benzyl, 4-[-NHC(O)CH<sub>2</sub>NHC(O)NH-3'-  
 10 methylphenyl]benzyl, 4-[-NHC(O)CH(NHBoc)(CH<sub>2</sub>)<sub>4</sub>NHCbz]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>CH(C(O)OCH<sub>2</sub>φ)-NHCbz]benzyl, 4-φ-benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)NHBoc]benzyl,  
 4-[H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(BocHNCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-)benzyl, 4-[φCH<sub>2</sub>OCH<sub>2</sub>(BocHN)CHC(O)NH-]benzyl,  
 15 4-[CH<sub>3</sub>NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(N-methylpiperidin-4-oxy)-benzyl,  
 4-[CH<sub>3</sub>N(Boc)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[φCH<sub>2</sub>OCH<sub>2</sub>(H<sub>2</sub>N)CHC(O)NH-]benzyl, 4-[HO(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-]benzyl, 4-[φCH<sub>2</sub>O(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 4-[HO(O)C(NH<sub>2</sub>)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 20 4-[CH<sub>3</sub>(N-Boc)NCH<sub>2</sub>C(O)NH-]benzyl, 4-[CH<sub>3</sub>NHCH<sub>2</sub>C(O)NH-]benzyl,  
 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>C(O)NH-]benzyl, 4-[-O-CH(COOH)φ]benzyl,  
 4-[2-carboxylphenyl]-benzyl, 4-[2-carboxylmethylphenyl]-benzyl  
 4-[φCH<sub>2</sub>OC(O)NHCH<sub>2</sub>CH<sub>2</sub>NH-]benzyl, 4-N[-(SO<sub>2</sub>)CH<sub>3</sub>]-  
 CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]benzyl, 4-*t*-butyl-O(O)CCH<sub>2</sub>-O-benzylNH]benzyl,  
 25 4-[N,N-di(4-N,N-dimethylamino)benzyl]amino]benzyl,  
 4-(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl-CH<sub>2</sub>NH)benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1'-(4'-pyrimidinyl)-piperazinyl]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-pyrrolidinyl)-  
 benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)]-benzyl,  
 30 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-O-]benzyl, 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>O-]benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(1'-(4'-methylpiperazinyl))]-benzyl,  
 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-4-(3'-chlorophenyl)-piperazin-1-yl]-benzyl,



-- 216 --

- 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(φ)CH<sub>2</sub>CH<sub>3</sub>]-benzyl, 4-[-OCH<sub>2</sub>-3'-(N-Boc)-piperidiny]-benzyl, 4-[-O-(3-(N-Boc)-piperidiny)]benzyl, 3-[-O-(N-methylpiperidin-4-yl)]benzyl, 4-[-O-(N-methylpiperidin-4-yl)]benzyl, 4-[di-iso-propylamino-CH<sub>2</sub>CH<sub>2</sub>O]-benzyl, 4-[N-3-methylbutyl-N-trifluoromethanesulfonyl]amino]benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-morpholinyl)]-benzyl, 4-[-OCH<sub>2</sub>CH(NHBoc)CH<sub>2</sub>cyclohexyl]-benzyl, 4-[OCH<sub>2</sub>CH<sub>2</sub>-(N-piperidiny)]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(4-*m*-chlorophenyl)-piperazin-1-yl]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-homopiperidiny)]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(benzyl)<sub>2</sub>]-benzyl, 3-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)benzyl]-benzyl, 4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethyl-O]-benzyl, [cyclopentylacetylenyl]-benzyl, 4-[-C≡C-φ-4'φ]-benzyl, 4-[-C≡C-CH<sub>2</sub>-O-S(O)<sub>2</sub>-4'-CH<sub>3</sub>-φ]-benzyl, 4-[-C≡C-CH<sub>2</sub>NHC(O)NH<sub>2</sub>]-benzyl, 4-[-C≡C-CH<sub>2</sub>-O-(4'-COOCH<sub>2</sub>CH<sub>3</sub>)φ]-benzyl, 4-[-C≡C-CH(NH<sub>2</sub>)-cyclohexyl]-benzyl, 4-[-C≡C-CH<sub>2</sub>-O-phenyl]-benzyl, 4-[-C≡C-CH<sub>2</sub>OCH<sub>3</sub>]-benzyl, 4-[-C≡C-CH<sub>2</sub>-O-(4'-C(O)OC<sub>2</sub>H<sub>5</sub>)phenyl]-benzyl, 4-[-C≡C-CH<sub>2</sub>CH(C(O)OCH<sub>3</sub>)<sub>2</sub>]-benzyl, 4-[-C≡C-CH<sub>2</sub>CH(NHC(O)CH<sub>3</sub>)C(O)OH]-benzyl, 4-[-C≡C-CH<sub>2</sub>NH-(4,5-dihydro-4-oxo-5-phenyl-oxazol-2-yl)]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(N-morpholino)]-benzyl, 4-[-OCH<sub>2</sub>COOH]-benzyl, 4-[-OCH<sub>2</sub>COO-*t*-butyl]-benzyl, 4-[-N(SO<sub>2</sub>CH<sub>3</sub>)(CH<sub>2</sub>)<sub>3</sub>-N(CH<sub>3</sub>)<sub>2</sub>]-benzyl, 4-[-NHS(O)<sub>2</sub>CF<sub>3</sub>]-benzyl, 4-[-C(=NH)NH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>Cl]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)O-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)OH]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-(3-methoxypyrrol-2-yl)-piperazinyl)-benzyl, 4-[-OCH<sub>2</sub>C(O)NH<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-*t*-butyl]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-phenyl)-piperidiny]-benzyl, 4-[-NHSO<sub>2</sub>-CH=CH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>Cl]-benzyl, 4-benzylbenzyl, 4-[-OCH<sub>2</sub>C(O)piperidin-1-yl]benzyl, 4-[-OCH<sub>2</sub>C(O)N(CH(CH<sub>3</sub>)<sub>2</sub>)<sub>2</sub>]-benzyl, 4-amidinobenzyl, 4-acetamidobenzyl, 4-(N-methyl)acetamidobenzyl, 4-(NHC(O)CH<sub>2</sub>NHC(O)NH-fluorescein)benzyl, 4-(NHC(O)CH<sub>2</sub>CH(NH<sub>2</sub>)COOH, (1-toluenesulfonylimidizol-4-yl)-methyl-, [(1-*N,N*-dimethylaminosulfonyl)-imidizol-4-yl]methyl-, 4-(*N*-toluenesulfonylamino)benzyl, and 4-(*N*-methyltrifluoroacetamido)phenyl.

-- 217 --

22. The method according to Claim 13 wherein R<sup>6'</sup> is selected from the group consisting of 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), methoxy, ethoxy, *iso*-propoxy, *n*-butoxy, *t*-butoxy, cyclopentoxo, *neo*-pentoxo, 2- $\alpha$ -*iso*-propyl-4- $\beta$ -methylcyclohexoxy, 2- $\beta$ -isopropyl-4- $\beta$ -methylcyclohexoxy, 5 -NH<sub>2</sub>, benzyloxy, -NHCH<sub>2</sub>COOH, -NHCH<sub>2</sub>CH<sub>2</sub>COOH, -NH-adamantyl, -NHCH<sub>2</sub>CH<sub>2</sub>COOCH<sub>2</sub>CH<sub>3</sub>, -NHSO<sub>2</sub>-*p*-CH<sub>3</sub>- $\phi$ , -NHOR<sup>8</sup> where R<sup>8</sup> is hydrogen, methyl, *iso*-propyl or benzyl, O-(N-succinimidyl), -O-cholest-5-en-3- $\beta$ -yl, -OCH<sub>2</sub>-OC(O)C(CH<sub>3</sub>)<sub>2</sub>, -O(CH<sub>2</sub>)<sub>z</sub>NHC(O)W where z is 1 or 2 and W is selected from the group consisting of pyrid-3-yl, N-10 methylpyridyl, and N-methyl-1,4-dihydro-pyrid-3-yl, -NR''C(O)-R' where R' is aryl, heteroaryl or heterocyclic and R'' is hydrogen or -CH<sub>2</sub>C(O)OCH<sub>2</sub>CH<sub>3</sub>.

23. The method according to Claim 13 wherein said compound is selected from the group consisting of:

- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonylglycyl)amino]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(glycyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(carboxy)propionamido]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl-L-alanyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl-D-alanyl)amino]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl-D-phenylalanyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[3-(fluorescein)thiouriedo]acetamido}-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonylglycyl)amino]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[3-(3-methylphenyl)uriedo]acetamido}-L-phenylalanine

-- 218 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*α-*tert*-butoxycarbonyl-*N*ε-carbobenzyloxy-L-lysyl)amino]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[γ-(α-benzyl-*N*α-carbobenzyloxy-L-aspartyl)amino]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*α-*tert*-butoxycarbonyl-L-lysyl)amino]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[γ-(L-aspartyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(4-aminobutyramido)-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-*tert*-butoxycarbonylamino)butyramido]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-methylamino)butyramido]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4-(*N*-*tert*-butoxycarbonyl-*N*-methylamino)butyramido]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*O*-benzyl)-L-seryl)amino]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[δ-(D,L-glutamyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonylsarcosyl)amino]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N*-*tert*-butoxycarbonylsarcosyl)amino]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine ethyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(sarcosyl)amino]-L-phenylalanine
- 
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine

-- 219 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-( $\alpha$ -carboxybenzyloxy)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenyl]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(methoxycarbonyl)phenyl]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[3-(*N,N*-dimethylamino)propyl]-*N*-[trifluoromethanesulfonyl]amino}-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[4-[(*tert*-butoxycarbonyl)methoxy]benzyl]amino}-L-phenylalanine methyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N,N*-di[4-(*N,N*-dimethylamino)benzyl]amino}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl)methyl]amino}-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-*N*-methyl-L-serinyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-dimethylamino)ethoxy]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N*-ethyl-*N*-phenylamino)ethoxy]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-diisopropylamino)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-cyclohexyl-2-(*N*-*tert*-butoxycarbonylamino)propoxy]-L-phenylalanine methyl ester
- 45 *N*-(thiophene-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

-- 220 --

- N*-(5-chlorothiophene-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(*N,N*-diethylamino)ethoxy]-L-phenylalanine
- N*-(2,5-dichlorothiophene-3-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 10 *N*-(1-methylpyrazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-diethylamino)propoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(thiazole-2-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-diethylamino)propoxy]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(1-methylimidazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(2-methylthiadiazole-5-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(4-cyanobenzenesulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-(3,3-dimethyl)prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine ethyl ester

-- 221 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine methyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-(thiamorpholin-3-carbonyl)-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(cyclopentyl)ethynyl]-D,L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-[4-(phenyl)phenyl]ethynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(toluene-4-sulfonyloxy)prop-1-ynyl]-D,L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(ureido)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(1-aminocyclohex-1-yl)ethynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(phenoxy)prop-1-ynyl]-D,L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(phenoxy)prop-1-ynyl]-D,L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(methoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(methoxy)prop-1-ynyl]-D,L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[4,4-di(methoxycarbonyl)but-1-ynyl]-D,L-phenylalanine

-- 222 --

- N*-(toluene-4-sulfonyl)sarcosyl-4-[4,4-di(methoxycarbonyl)but-1-ynyl]-D,L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(4-acetamido-4-carboxybut-1-ynyl)-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-4-(4-acetamido-4-carboxybut-1-ynyl)-D,L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[(4,5-dihydro-4-oxo-5-phenyloxazol-2-yl)amino]prop-1-ynyl}-D,L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)sarcosyl-4-{3-[(4,5-dihydro-4-oxo-5-phenyloxazol-2-yl)amino]prop-1-ynyl}-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenoxy]-L-phenylalanine methyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{2-[4-(pyrimidin-2-yl)piperazin-1-yl]ethoxy}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(piperidin-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine methyl ester

-- 224 --


- N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(chloromethanesulfonyl)amino-L-phenylalanine methyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-trifluoromethanesulfonyl-*N*-isobutyl)amino-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-benzylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine methyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(aminocarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butylaminocarbonyl)methoxy]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(4-phenyl-4-hydroxypiperidin-1-yl)ethoxy]-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(piperidin-1-ylcarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 45 *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(amidino)phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(aminocarbonyl)phenylalanine



-- 224 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(chloromethanesulfonyl)amino-L-phenylalanine methyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-trifluoromethanesulfonyl)-*N*-isobutylamino-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-*N*-(vinylsulfonyl)amino-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-benzylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(benzyloxycarbonyl)methoxy]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(carboxy)methoxy]-L-phenylalanine methyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(aminocarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butylaminocarbonyl)methoxy]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(4-phenyl-4-hydroxypiperidin-1-yl)ethoxy]-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(piperidin-1-ylcarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 45 *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(amidino)phenylalanine
- N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(aminocarbonyl)phenylalanine

-- 226 --



IA

5

where

10  $\text{R}^1$  is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;


15  $\text{R}^2$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $\text{R}^1$  and  $\text{R}^2$  together with the nitrogen atom bound to  $\text{R}^2$  and the  $\text{SO}_2$  group bound to  $\text{R}^1$  can form a heterocyclic or a substituted heterocyclic group;

20  $\text{R}^3$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and, when  $\text{R}^2$  does not form a heterocyclic group with  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  together with the nitrogen atom bound to  $\text{R}^2$  and the carbon atom bound to  $\text{R}^3$  can form a heterocyclic or a substituted heterocyclic group;

$\text{R}^5$  is  $-(\text{CH}_2)_x\text{-Ar-R}^5$  where  $\text{R}^5$  is selected from the group consisting

25 (a) substituted alkylcarbonylamino with the proviso that at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, 30 carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone.

-- 226 --



IA

where

R<sup>1</sup> is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and R<sup>1</sup> and R<sup>2</sup> together with the nitrogen atom bound to R<sup>2</sup> and the SO<sub>2</sub> group bound to R<sup>1</sup> can form a heterocyclic or a substituted heterocyclic group;

R<sup>3</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and, when R<sup>2</sup> does not form a heterocyclic group with R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> and the carbon atom bound to R<sup>3</sup> can form a heterocyclic or a substituted heterocyclic group;

R<sup>5</sup> is -(CH<sub>2</sub>)<sub>x</sub>-Ar-R<sup>5'</sup> where R<sup>5'</sup> is selected from the group consisting

(a) substituted alkylcarbonylamino with the proviso that at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone,

-- 228 --

(c) aryl and heteroaryl;

(d) -NR'R' wherein each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic with the proviso that at least one of R' is substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic and with the further proviso that when R' is substituted alkyl at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl,

-- 228 --

(c) aryl and heteroaryl;

(d) -NR'R' wherein each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic with the proviso that at least one of R' is substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclic and substituted heterocyclic and with the further proviso that when R' is substituted alkyl at least one of the substituents on the substituted alkyl moiety is selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkenyl, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryloxy, substituted aryloxy, cyano, nitro, halogen, hydroxyl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl,

-- 230 --

cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(f) substituted alkenyl or substituted alkynyl with the proviso that at least one of the substituents on the substituted alkenyl/alkynyl moiety is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when substituted with substituted alkyl then at least one of the substituents on the

-- 230 --

cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(f) substituted alkenyl or substituted alkynyl with the proviso that at least one of the substituents on the substituted alkenyl/alkynyl moiety is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic with the proviso that when substituted with substituted alkyl then at least one of the substituents on the

-- 232 --

blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl;

(g) substituted aryloxy and substituted heteroaryloxy with the proviso that at least one substituent on the substituted aryloxy/heteroaryloxy is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(h) -alkoxy-saturated heterocyclic, -alkoxy-saturated substituted heterocyclic, -substituted alkoxy-heterocyclic and -substituted alkoxy-substituted saturated heterocyclic;

(i) -O-heterocyclic and -O-substituted heterocyclic;

(j) tetrazolyl;

(k) -NR-SO<sub>2</sub>-substituted alkyl where R is hydrogen, alkyl or aryl, with the proviso that at least one substituent on the alkyl moiety of the substituted alkylsulfonylamino is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(l) alkenylsulfonylamino, alkynylsulfonylamino, substituted alkenylsulfonylamino and substituted alkynylsulfonylamino;

(m) substituted alkoxy with the proviso that the substitution on the alkyl moiety of said substituted alkoxy does not include alkoxy-NR"R", unsaturated heterocyclyl, alkyloxy, aryloxy, heteroaryloxy, aryl, heteroaryl and aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-



-- 232 --

blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NR<sub>2</sub> where R is hydrogen or alkyl;

(g) substituted aryloxy and substituted heteroaryloxy with the proviso that at least one substituent on the substituted aryloxy/heteroaryloxy is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(h) -alkoxy-saturated heterocyclic, -alkoxy-saturated substituted heterocyclic, -substituted alkoxy-heterocyclic and -substituted alkoxy-substituted saturated heterocyclic;

(i) -O-heterocyclic and -O-substituted heterocyclic;

(j) tetrazolyl;

(k) -NR-SO<sub>2</sub>-substituted alkyl where R is hydrogen, alkyl or aryl, with the proviso that at least one substituent on the alkyl moiety of the substituted alkylsulfonylamino is other than halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino, alkenylamino, alkynylamino, alkylcarbonyloxy, acyl, alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkylurea;

(l) alkenylsulfonylamino, alkynylsulfonylamino, substituted alkenylsulfonylamino and substituted alkynylsulfonylamino;

(m) substituted alkoxy with the proviso that the substitution on the alkyl moiety of said substituted alkoxy does not include alkoxy-NR<sup>2</sup>"R", unsaturated heterocyclyl, alkyloxy, aryloxy, heteroaryloxy, aryl, heteroaryl and aryl/heteroaryl substituted with halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, 1,2-

-- 234 --

of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic:

(s)  $-NR'C(O)OR^9$  wherein  $R'$  is selected from the group consisting of  
 5 alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and,  $R^9$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

10 (t) -aminocarbonyl-(N-formylheterocyclyl); and

(u) -alkyl-C(O)NH-heterocyclyl and -alkyl-C(O)NH-substituted heterocyclyl,

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl,

x is an integer of from 1 to 4;

15  $R^{6'}$  is selected from the group consisting of, hydroxy, 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), amino, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, -O-(N-succinimidyl), -NH-adamantyl, -O-cholest-5-en-3- $\beta$ -yl, -NHOY where Y is hydrogen, alkyl, substituted alkyl, aryl, and substituted aryl, -NH(CH<sub>2</sub>)<sub>p</sub>COOY where p is an integer of from 1 to  
 20 8 and Y is as defined above, -OCH<sub>2</sub>NR<sup>9</sup>R<sup>10</sup> where R<sup>9</sup> is selected from the group consisting of -C(O)-aryl and -C(O)-substituted aryl and R<sup>10</sup> is selected from the group consisting of hydrogen and -CH<sub>2</sub>COOR<sup>11</sup> where R<sup>11</sup> is alkyl, and -NHSO<sub>2</sub>Z where Z is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
 25 heterocyclic and substituted heterocyclic;

Q is -C(X)NR<sup>7</sup>- wherein R<sup>7</sup> is selected from the group consisting of hydrogen and alkyl; and X is selected from the group consisting of oxygen and sulfur,

and pharmaceutically acceptable salts thereof

30 with the proviso that when R<sup>1</sup> is *p*-CH<sub>3</sub>- $\phi$ -, R<sup>6</sup> is methoxy, Q is -C(O)NH-, and R<sup>2</sup> and R<sup>3</sup> are joined to form a pyrrolidinyl group, then R<sup>5</sup> is not *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(isopropyl)]-benzyl-, *p*-

-- 234 --

of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic:

- (s) -NR'C(O)OR<sup>9</sup> wherein R' is selected from the group consisting of  
 5 alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and, R<sup>9</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;  
 10 (t) -aminocarbonyl-(N-formylheterocycl); and  
 (u) -alkyl-C(O)NH-heterocycl and -alkyl-C(O)NH-substituted heterocycl,

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl,

x is an integer of from 1 to 4;

- 15 R<sup>6'</sup> is selected from the group consisting of, hydroxy, 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), amino, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, -O-(N-succinimidyl), -NH-adamantyl, -O-cholest-5-en-3-β-yl, -NHOY where Y is hydrogen, alkyl, substituted alkyl, aryl, and substituted aryl, -NH(CH<sub>2</sub>)<sub>p</sub>COOY where p is an integer of from 1 to  
 20 8 and Y is as defined above, -OCH<sub>2</sub>NR<sup>9</sup>R<sup>10</sup> where R<sup>9</sup> is selected from the group consisting of -C(O)-aryl and -C(O)-substituted aryl and R<sup>10</sup> is selected from the group consisting of hydrogen and -CH<sub>2</sub>COOR<sup>11</sup> where R<sup>11</sup> is alkyl, and -NHSO<sub>2</sub>Z where Z is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
 25 heterocyclic and substituted heterocyclic;

Q is -C(X)NR<sup>7</sup>- wherein R<sup>7</sup> is selected from the group consisting of hydrogen and alkyl; and X is selected from the group consisting of oxygen and sulfur;

and pharmaceutically acceptable salts thereof

- 30 with the proviso that when R<sup>1</sup> is *p*-CH<sub>3</sub>-φ-, R<sup>6</sup> is methoxy, Q is -C(O)NH-, and R<sup>2</sup> and R<sup>3</sup> are joined to form a pyrrolidinyl group, then R<sup>5</sup> is not *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]-benzyl-, *p*-[-OCH<sub>2</sub>CH<sub>2</sub>N(isopropyl)<sub>2</sub>]-benzyl-, *p*-

-- 236 --

bound to R<sup>1</sup> are joined to form a heterocyclic group or substituted heterocyclic group.

30. The pharmaceutical composition according to Claim 25 wherein R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> substituent and the carbon bound to the R<sup>3</sup> substituent form a heterocyclic group or a substituted heterocyclic group.

31. The pharmaceutical composition according to Claim 25 wherein R<sup>3</sup> is selected from the group consisting of methyl, phenyl, benzyl, diphenylmethyl, -CH<sub>2</sub>CH<sub>2</sub>-COOH, -CH<sub>2</sub>-COOH, 2-amidoethyl, *iso*-butyl, *t*-butyl, -CH<sub>2</sub>O-benzyl and hydroxymethyl.

32. The pharmaceutical composition according to Claim 25 wherein Q is -C(O)NH- or -C(S)NH-.

33. The pharmaceutical composition according to Claim 25 wherein R<sup>3</sup> is selected from the group consisting of:

4-[NH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-[HOOCCH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>NHBoc]benzyl, 4-[-NHC(O)CH(CH<sub>3</sub>)NHBoc]benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>φ)NHBoc]benzyl, 4-[-NHC(O)CH<sub>2</sub>NHC(O)NH-3'-  
 methylphenyl]benzyl, 4-[-NHC(O)CH(NHBoc)(CH<sub>2</sub>)<sub>4</sub>NHCbz]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>CH(C(O)OCH<sub>2</sub>φ)-NHCbz]benzyl, 4-φ-benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)NHBoc]benzyl,  
 4-[H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(BocHNCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-)benzyl, 4-[φCH<sub>2</sub>OCH<sub>2</sub>(BocHN)CHC(O)NH-]benzyl,  
 4-[CH<sub>3</sub>NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(N-methylpiperidin-4-oxy)-benzyl,  
 4-[CH<sub>3</sub>N(Boc)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[φCH<sub>2</sub>OCH<sub>2</sub>(H<sub>2</sub>N)CHC(O)NH-]benzyl, 4-[HO(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-]benzyl, 4-[φCH<sub>2</sub>O(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 4-[HO(O)C(NH<sub>2</sub>)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 4-[CH<sub>3</sub>(N-Boc)NCH<sub>2</sub>C(O)NH-]benzyl, 4-[CH<sub>3</sub>NHCH<sub>2</sub>C(O)NH-]benzyl,

-- 236 --

bound to R<sup>1</sup> are joined to form a heterocyclic group or substituted heterocyclic group.

30. The pharmaceutical composition according to Claim 25 wherein  
 5 R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom bound to R<sup>2</sup> substituent and the carbon bound to the R<sup>3</sup> substituent form a heterocyclic group or a substituted heterocyclic group.

31. The pharmaceutical composition according to Claim 25 wherein  
 10 R<sup>3</sup> is selected from the group consisting of methyl, phenyl, benzyl, diphenylmethyl, -CH<sub>2</sub>CH<sub>2</sub>-COOH, -CH<sub>2</sub>-COOH, 2-amidoethyl, *iso*-butyl, *t*-butyl, -CH<sub>2</sub>O-benzyl and hydroxymethyl.

32. The pharmaceutical composition according to Claim 25 wherein Q  
 15 is -C(O)NH- or -C(S)NH-.

33. The pharmaceutical composition according to Claim 25 wherein  
 R<sup>5</sup> is selected from the group consisting of:  
 20 4-[NH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-[HOOCCH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>NHBoc]benzyl, 4-[-NHC(O)CH(CH<sub>3</sub>)NHBoc]benzyl,  
 4-[-NHC(O)CH(CH<sub>2</sub>φ)NHBoc]benzyl, 4-[-NHC(O)CH<sub>2</sub>NHC(O)NH-3'-  
 methylphenyl]benzyl, 4-[-NHC(O)CH(NHBoc)(CH<sub>2</sub>)<sub>4</sub>NHCbz]benzyl,  
 4-[-NHC(O)CH<sub>2</sub>CH(C(O)OCH<sub>2</sub>φ)-NHCbz]benzyl, 4-φ-benzyl,  
 25 4-[-NHC(O)CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)NHBoc]benzyl,  
 4-[H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(BocHNCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-)benzyl, 4-[φCH<sub>2</sub>OCH<sub>2</sub>(BocHN)CHC(O)NH-]benzyl,  
 4-[CH<sub>3</sub>NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl, 4-(N-methylpiperidin-4-oxy)-benzyl,  
 4-[CH<sub>3</sub>N(Boc)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
 30 4-[φCH<sub>2</sub>OCH<sub>2</sub>(H<sub>2</sub>N)CHC(O)NH-]benzyl, 4-[HO(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-  
 C(O)NH-]benzyl, 4-[φCH<sub>2</sub>O(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 4-[HO(O)C(NH<sub>2</sub>)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
 4-[CH<sub>3</sub>(*N*-Boc)NCH<sub>2</sub>C(O)NH-]benzyl, 4-[CH<sub>3</sub>NHCH<sub>2</sub>C(O)NH-]benzyl,

-- 238 --

4-[-NHS(O)<sub>2</sub>CF<sub>3</sub>]-benzyl, 4-[-C(=NH)NH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>Cl]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)O-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)OH]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-(3-methoxypyrrol-2-yl)-piperazinyl)-benzyl, 4-[-OCH<sub>2</sub>C(O)NH<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-*t*-butyl]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-phenyl)-piperidinyl]-benzyl, 4-[-NHSO<sub>2</sub>-CH=CH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>Cl]-benzyl, 4-benzyl-benzyl, 4-[-OCH<sub>2</sub>C(O)piperidin-1-yl]benzyl, 4-[-OCH<sub>2</sub>C(O)N(CH(CH<sub>3</sub>)<sub>2</sub>)<sub>2</sub>]benzyl, 4-amidinobenzyl, 4-acetamidobenzyl, 4-(*N*-methyl)acetamidobenzyl, 4-(NHC(O)CH<sub>2</sub>NHC(O)NH-fluorescein)benzyl, 4-(NHC(O)CH<sub>2</sub>CH(NH<sub>2</sub>)COOH, (1-toluenesulfonylimidizol-4-yl)-methyl-, [(1-*N,N*-dimethylaminosulfonyl)-imidizol-4-yl]methyl-, 4-(*N*-toluenesulfonyl-amino)benzyl, and 4-(*N*-methyltrifluoroacetamido)phenyl.

34. The pharmaceutical composition according to Claim 25 wherein R<sup>6'</sup> is selected from the group consisting of 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), methoxy, ethoxy, *iso*-propoxy, *n*-butoxy, *t*-butoxy, cyclopentoxy, *neo*-pentoxy, 2- $\alpha$ -*iso*-propyl-4- $\beta$ -methylcyclohexoxy, 2- $\beta$ -isopropyl-4- $\beta$ -methylcyclohexoxy, -NH<sub>2</sub>, benzyloxy, -NHCH<sub>2</sub>COOH, -NHCH<sub>2</sub>CH<sub>2</sub>COOH, -NH-adamantyl, -NHCH<sub>2</sub>CH<sub>2</sub>COOCH<sub>2</sub>CH<sub>3</sub>, -NHSO<sub>2</sub>-*p*-CH<sub>3</sub>- $\phi$ , -NHOR<sup>8</sup> where R<sup>8</sup> is hydrogen, methyl, *iso*-propyl or benzyl, O-(*N*-succinimidyl), -O-cholest-5-en-3- $\beta$ -yl, -OCH<sub>2</sub>-OC(O)C(CH<sub>3</sub>)<sub>3</sub>, -O(CH<sub>2</sub>)<sub>z</sub>NHC(O)W where *z* is 1 or 2 and W is selected from the group consisting of pyrid-3-yl, *N*-methylpyridyl, and *N*-methyl-1,4-dihydro-pyrid-3-yl, -NR''C(O)-R' where R' is aryl, heteroaryl or heterocyclic and R'' is hydrogen or -CH<sub>2</sub>C(O)OCH<sub>2</sub>CH<sub>3</sub>.

35. The pharmaceutical composition according to Claim 25 wherein said compound is selected from the group consisting of:

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)glycyl]amino]-L-phenylalanine

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(glycyl)amino]-L-phenylalanine

-- 238 --

4-[-NHS(O)<sub>2</sub>CF<sub>3</sub>]-benzyl, 4-[-C(=NH)NH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>Cl]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)O-benzyl]-benzyl, 4-[-OCH<sub>2</sub>C(O)OH]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-(3-methoxypyrrol-2-yl)-piperazinyl)-benzyl, 4-[-OCH<sub>2</sub>C(O)NH<sub>2</sub>]-benzyl, 4-[-OCH<sub>2</sub>C(O)NH-*t*-butyl]-benzyl, 4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-phenyl)-piperidinyl]-benzyl, 4-[-NHSO<sub>2</sub>-CH=CH<sub>2</sub>]-benzyl, 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>Cl]-benzyl, 4-benzyl-benzyl, 4-[-OCH<sub>2</sub>C(O)piperidin-1-yl]benzyl, 4-[-OCH<sub>2</sub>C(O)N(CH(CH<sub>3</sub>)<sub>2</sub>)<sub>2</sub>]benzyl, 4-amidinobenzyl, 4-acetamidobenzyl, 4-(*N*-methyl)acetamidobenzyl, 4-(NHC(O)CH<sub>2</sub>NHC(O)NH-fluorescein)benzyl, 4-(NHC(O)CH<sub>2</sub>CH(NH<sub>2</sub>)COOH, (1-toluenesulfonylimidazol-4-yl)-methyl-, 4-(*N,N*-dimethylaminosulfonyl)-imidazol-4-yl]methyl-, 4-(*N*-toluenesulfonylamino)benzyl, and 4-(*N*-methyltrifluoroacetamido)phenyl.

34. The pharmaceutical composition according to Claim 25 wherein R<sup>6'</sup> is selected from the group consisting of 2,4-dioxo-tetrahydrofuran-3-yl (3,4-enol), methoxy, ethoxy, *iso*-propoxy, *n*-butoxy, *t*-butoxy, cyclopentoxy, *neo*-pentoxy, 2-*α*-*iso*-propyl-4-*β*-methylcyclohexoxy, 2-*β*-isopropyl-4-*β*-methylcyclohexoxy, -NH<sub>2</sub>, benzyloxy, -NHCH<sub>2</sub>COOH, -NHCH<sub>2</sub>CH<sub>2</sub>COOH, -NH-adamantyl, -NHCH<sub>2</sub>CH<sub>2</sub>COOCH<sub>2</sub>CH<sub>3</sub>, -NHSO<sub>2</sub>-*p*-CH<sub>3</sub>-*φ*, -NHOR<sup>8</sup> where R<sup>8</sup> is hydrogen, methyl, *iso*-propyl or benzyl, O-(*N*-succinimidyl), -O-cholest-5-en-3-*β*-yl, -OCH<sub>2</sub>-OC(O)C(CH<sub>3</sub>)<sub>3</sub>, -O(CH<sub>2</sub>)<sub>z</sub>NHC(O)W where *z* is 1 or 2 and W is selected from the group consisting of pyrid-3-yl, *N*-methylpyridyl, and *N*-methyl-1,4-dihydro-pyrid-3-yl, -NR''C(O)-R' where R' is aryl, heteroaryl or heterocyclic and R'' is hydrogen or -CH<sub>2</sub>C(O)OCH<sub>2</sub>CH<sub>3</sub>.

35. The pharmaceutical composition according to Claim 25 wherein said compound is selected from the group consisting of:

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxycarbonyl)glycyl]amino]-L-phenylalanine

*N*-(toluene-4-sulfonyl)-L-prolyl-4-[(glycyl)amino]-L-phenylalanine

-- 240 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxyl-carbonylsarcosyl)amino]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N*-*tert*-butoxyl-carbonylsarcosyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine ethyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(sarcosyl)amino]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-( $\alpha$ -carboxybenzyloxy)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenyl]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(methoxycarbonyl)phenyl]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[3-(*N,N*-dimethylamino)propyl]-*N*-[trifluoromethanesulfonyl]amino}-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[4-[(*tert*-butoxycarbonyl)methoxy]benzyl]amino}-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N,N*-di[4-(*N,N*-dimethylamino)benzyl]amino}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl)methyl]amino}-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine



-- 240 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butoxyl-carbonylsarcosyl)amino]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N*-*tert*-butoxyl-carbonylsarcosyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine ethyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(sarcosyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(sarcosyl)amino]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[(*N,N*-dimethylglycyl)amino]-L-phenylalanine
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-( $\alpha$ -carboxybenzyloxy)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(carboxy)phenyl]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(methoxycarbonyl)phenyl]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{*N*-[2-(*N*-carbobenzyloxyamino)ethyl]amino}-L-phenylalanine methyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[3-(*N,N*-dimethylamino)propyl]-*N*-[trifluoromethanesulfonyl]amino}-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[4-[(*tert*-butoxycarbonyl)methoxy]benzyl]amino}-L-phenylalanine methyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N,N*-di[4-(*N,N*-dimethylamino)benzyl]amino}-L-phenylalanine
- 
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-{*N*-[(2-formyl-1,2,3,4-tetrahydroisoquinolin-3-yl)methyl]amino}-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine

-- 242 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine methyl ester
- 5 *N*-(1-methylimidazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(2-methylthiadiazole-5-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(4-cyanobenzenesulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-(3,3-dimethyl)prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine ethyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-(thiamorpholin-3-carbonyl)-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(cyclopentyl)ethynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(4-(phenyl)phenyl)ethynyl]-D,L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(toluene-4-sulfonyloxy)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(ureido)prop-1-ynyl]-D,L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(1-aminocyclohex-1-yl)ethynyl]-D,L-phenylalanine

-- 242 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(*N*-methyl-*N*-benzylamino)propoxy]-L-phenylalanine methyl ester
- 5 *N*-(1-methylimidazole-4-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(2-methylthiadiazole-5-sulfonyl)-L-prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 10 *N*-(toluene-4-sulfonyl)-L-thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(4-cyanobenzenesulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-(3,3-dimethyl)prolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine ethyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-(thiamorpholin-3-carbonyl)-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethoxy]-L-phenylalanine
- 30 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(cyclopentyl)ethynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-[4-(phenyl)phenyl]ethynyl]-D,L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(toluene-4-sulfonyloxy)prop-1-ynyl]-D,L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(ureido)prop-1-ynyl]-D,L-phenylalanine
- N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[3-(4-ethoxycarbonylphenoxy)prop-1-ynyl]-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-D,L-phenylalanyl-4-[2-(1-aminocyclohex-1-yl)ethynyl]-D,L-phenylalanine

-- 244 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine methyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(piperidin-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine methyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine methyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-[4-(3-methoxythien-2-yl)-4-hydroxypiperidin-1-yl]ethoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-3-(1-methylpiperidin-4-oxyl)-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(1-methylpiperidin-4-oxyl)-D,L-phenylalanine

-- 244 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(piperidin-1-yl)propoxy]-L-phenylalanine methyl ester
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(1-*tert*-butoxycarbonylpiperidin-3-yl)methoxy]-L-phenylalanine
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(piperidin-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-{3-[4-(3-chlorophenyl)piperazin-1-yl]propoxy}-L-phenylalanine methyl ester
- 20 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(azepan-1-yl)ethoxy]-L-phenylalanine methyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(pyrrolidin-1-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[3-(4-methylpiperazin-1-yl)propoxy]-L-phenylalanine
- 35 *N*-(toluene-4-sulfonyl)-L-prolyl-3-[2-(morpholin-4-yl)ethoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-[4-(3-methoxythien-2-yl)-4-hydroxypiperidin-1-yl]ethoxy]-L-phenylalanine methyl ester
- 40
- 
- N*-(toluene-4-sulfonyl)-L-prolyl-3-(1-methylpiperidin-4-oxy)-D,L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(1-methylpiperidin-4-oxy)-D,L-phenylalanine

-- 246 --

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(aminocarbonyl)methoxy]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butylaminocarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(4-phenyl-4-hydroxypiperidin-1-yl)ethoxy]-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(piperidin-1-ylcarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(amidino)phenylalanine
- 20 *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(aminocarbonyl)phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine isopropyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methyltrifluoroacetamido)-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine *t*-butyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-(*N*-methylpiperidinoxy)-phenylalanine *t*-butyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-L-(4-methylpiperidinoxy) phenylalanine *t*-butyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine *t*-butyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine
- 45 and pharmaceutically acceptable salts thereof as well as any of the ester compounds recited above wherein one ester is replaced with another ester

- 246 -

- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(aminocarbonyl)methoxy]-L-phenylalanine
- 5 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N*-*tert*-butylaminocarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[2-(4-phenyl-4-hydroxypiperidin-1-yl)ethoxy]-L-phenylalanine methyl ester
- 10 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(piperidin-1-ylcarbonyl)methoxy]-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- 15 *N*-(toluene-4-sulfonyl)-L-prolyl-4-[(*N,N*-diisopropylaminocarbonyl)methoxy]-L-phenylalanine methyl ester
- N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(amidino)phenylalanine
- 20 *N*-(toluene-4-sulfonyl)sarcosyl-D,L-4-(aminocarbonyl)phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine isopropyl ester
- 25 *N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methylacetamido)-L-phenylalanine
- N*-(toluene-4-sulfonyl)-L-prolyl-4-(*N*-methyltrifluoroacetamido)-L-phenylalanine methyl ester
- 30 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-4-[3-(*N,N*-dimethylamino)propoxy]-L-phenylalanine *t*-butyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-L-4-(*N*-methylpiperidinoxy)-phenylalanine *t*-butyl ester
- 35 *N*-(toluene-4-sulfonyl)-L-(5,5-dimethyl)thiaprolyl-L-(4-methylpiperidinoxy) phenylalanine *t*-butyl ester
- N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine *t*-butyl ester
- 40 *N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine
- 45 *N*-(toluene-4-sulfonyl)-L-prolyl-(4-phenyl)-L-phenylalanine

and pharmaceutically acceptable salts thereof as well as any of the ester compounds recited above wherein one ester is replaced with another ester

# INTERNATIONAL SEARCH REPORT

Internat'l Application No  
PCT/US 98/15313

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K5/062 C07K5/065 C07K5/078 A61K38/05

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 23 57 334 A (HOFFMANN LA ROCHE) 6 June 1974 see claims; example 1C ---	2,3,5
A	WO 96 22966 A (BIOGEN INC ; ADAMS STEVEN P (US); LIN KO CHUNG (US); LEE WEN CHERNG) 1 August 1996 see page 46, line 32 - page 47, line 35; claims; examples ---	1,13,25, 36,37
A	WO 95 15973 A (CYTEL CORP) 15 June 1995  see page 8, line 16 - line 26; claims; examples --- -/--	1,13,25, 36,37

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"S" document member of the same patent family

Date of the actual completion of the international search

9 December 1998

Date of mailing of the international search report

07/01/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fuhr, C



# INTERNATIONAL SEARCH REPORT

Int. No. 1st Application No.  
PCT/US 98/15313

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C07K5/062 C07K5/065 C07K5/078 A61K38/05		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. PIDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 23 57 334 A (HOFFMANN LA ROCHE) 6 June 1974 see claims; example 1C ---	2,3,5
A	WO 96 22966 A (BIOGEN INC ;ADAMS STEVEN P (US); LIN KO CHUNG (US); LEE WEN CHERNG) 1 August 1996 see page 46, line 32 - page 47, line 35; claims; examples ---	1,13,25, 36,37
A	WO 95 15973 A (CYTEL CORP) 15 June 1995 see page 8, line 16 - line 26; claims; examples --- -/--	1,13,25, 36,37
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claims or which is cited to establish the publication date of another citation or other special reason (to be specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combinations being obvious to a person skilled in the art "A" document member of the same patent family		
Date of the actual completion of the international search  9 December 1998		Date of mailing of the international search report  07/01/1999
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2260 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 881 000 rd, Fax: (+31-70) 340-3016		Authorized officer  Fuhr, C

Form PCT/ISA/210 (October 1997) (July 1998)

# INTERNATIONAL SEARCH REPORT

Int. national application No.

PCT/US 98/15313

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 36 and 37 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful international Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Indication of Application No.

PCT/US 98/15313

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2357334 A	06-06-1974	AU 6147073 A BE 807704 A FR 2207715 A JP 49082682 A NL 7314486 A ZA 7307898 A	17-04-1975 24-05-1974 21-06-1974 08-08-1974 28-05-1974 31-07-1974
WO 9622966 A	01-08-1996	AU 4911596 A BG 101841 A BR 9606778 A CA 2211181 A CN 1177343 A CZ 9702340 A EP 0805796 A FI 973087 A HU 9702461 A NO 973384 A PL 321848 A SK 98797 A	14-08-1996 30-04-1998 06-01-1998 01-08-1996 25-03-1998 18-03-1998 12-11-1997 22-09-1997 28-04-1998 19-09-1997 22-12-1997 04-02-1998
WO 9515973 A	15-06-1995	AU 693143 B AU 1215695 A CA 2177840 A CN 1142832 A EP 0737204 A JP 9509146 T NZ 277206 A US 5688913 A US 5821231 A US 5770573 A	25-06-1998 27-06-1995 15-06-1995 12-02-1997 16-10-1996 16-09-1997 24-11-1997 18-11-1997 13-10-1998 23-06-1998
WO 9216549 A	01-10-1992	AU 1354292 A	21-10-1992
DE 2655636 A	23-06-1977	JP 52113919 A US 4055636 A US 4018915 A US 4018913 A US 4055651 A US 4041156 A US 4046876 A US 4070457 A US 4036955 A US 4073914 A US 4096255 A US 4104392 A BE 849196 A BE 863092 A CA 1102316 A CA 1137469 A CA 1140917 A CH 630349 A CH 630348 A DD 128681 A DK 550576 A, B. FI 763528 A, B. FR 2334349 A FR 2378004 A GR 63103 A	24-09-1977 25-10-1977 19-04-1977 19-04-1977 25-10-1977 09-08-1977 06-09-1977 24-01-1978 19-07-1977 14-02-1978 20-06-1978 01-08-1978 08-06-1977 19-07-1978 02-06-1981 14-12-1982 08-02-1983 15-06-1982 15-06-1982 07-12-1977 10-06-1977 10-06-1977 08-07-1977 18-08-1978 05-09-1979

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat. Application No.

PCT/US 98/15313

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2655636 A		LU 76347 A	24-06-1977
		LU 78911 A	28-09-1978
		NL 7613660 A, B	13-06-1977
		PT 65934 B	13-06-1978
		AT 367750 B	26-07-1982
		AT 205379 A	15-12-1981
		AT 356304 B	25-04-1980
		AT 905176 A	15-09-1979
		AU 515473 B	09-04-1981
		AU 2035076 A	15-06-1978
		AU 527063 B	17-02-1983
		AU 3757978 A	03-01-1980
		CS 228113 B	16-04-1984
		CS 228103 B	14-05-1984
		GB 1538207 A	10-01-1979
		GB 1538206 A	10-01-1979
		JP 52097934 A	17-08-1977
		SE 438680 B	29-04-1985
		SE 7613799 A	10-06-1977
		SU 965353 A	07-10-1982
		US 4097591 A	27-06-1978
		ZA 7607200 A	27-09-1978
		ZA 7803673 A	25-10-1978
		US 4069318 A	17-01-1978
		US 4071621 A	31-01-1978